CONTENTS

NUMBER 1, FEBRUARY, 1918

1
9
17
27
43
63
69
81
89
103
133
147
147
147 159
147 159
147 159

CONTENTS

XVIII. The Effect of Lactic Acid on the Respiratory Center. By Seymour	209
J. Cohen	221
XX. Investigations on the Composition of Oil of Chenopodium and the Anthelmintic Value of Some of its Components. By Maurice C. Hall and	231
Number 4, May, 1918	
XXII. Observations on the Effect of Intravenous Injections of Some Sodium Salts, with Special Reference to the Supposed Toxicity of Sodium Phosphate. By Isidor Greenwald	281 303
XXIV. The Treatment of Amoebic Dysentery with Chaparro Armagosa (Castela Nicholsoni of the Family Simarubaceae). By Andrew Watson Sellards and Monroe Anderson McIver	
Number 5, June, 1918	
XXV. Distribution of Gold in Animal Tissues. Studies on the Biochemistry and Chemotherapy of Tuberculosis. XVII. By Lydia M. DeWitt, Sidney M. Cadwell and Gladys Leavell	
XXVII. On the Relation between the Chemical Structure of the Opium Alkaloids and their Physiological Action on Smooth Muscle with a Pharmacological and Therapeutic Study of Some Benzyl Esters. I. On the Relation of the Chemical Structure of the Opium Alkaloids to their Action on Smooth Muscle. By David I. Macht	
Number 6, July, 1918	
XXVIII. Ou the Relation between the Chemical Structure of the Opium Alkaloids and their Physiological Action on Smooth Muscle with a Pharmachological and Therapeutic Study of Some Benzyl Esters. II. A Pharmacological and Therapeutic Study of Some Benzyl Esters. By David I. Macht	

ILLUSTRATIONS

•	
Experiment of April 20 (Fig. 1)	65
of April 27, 1917 (Fig. 2)	66
—— of April, 1917 (Fig. 3)	66
Perfusion with 1:7:9-trimethyl-2:8-dioxypurin (Fig. 1)	83
—— with caffein (Fig. 2)	84
	108
Ventricular tracing of perfused rabbit's heart during auricular fibrillation	
from electrical stimulation (Fig. 2)	111
tracing of perfused rabbit's heart (Fig. 3)	112
— tracing of rabbit's heart perfused for thirty minutes (Fig. 4)	115
Tracings of the ventricle and auricle of rabbit's heart (Fig. 5)	
Ventricular tracing in perfused heart under strophanthin (Fig. 6)	
Tracing of ventricle and blood pressure in cat (Fig. 7)	
Rabbit's heart perfused with Ringer containing strophanthin (Fig. 8)	
(Fig. 1)	
(Fig. 2)	
(Fig. 1)	
(Fig. 2)	
(Fig. 3)	
(Fig. 4) folderfacing	
Folder	
Successive microphotographs of the same scale (Fig. 1)	
Schematic median section through the thermostat reservoir (Fig. 2)	
Warming stage (Fig. 3)	
Dipping device (Fig. 4)	
Blood pressure and respiratory tracing (Fig. 1)	
— pressure and respiratory tracing (Fig. 2)	
— pressure and respiratory tracing (Fig. 3)facing	
— pressure and respiratory tracing (Fig. 4)	
Intravenous injection of a solution of benzyl alcohol one per cent in physio-	
logical NaCl (Fig. 1)	269
Manchurian scorpion (Fig. 1)	
Venom glands (Fig. 2)	
— glands (Fig. 3)	
Pig's ureter (Fig. 1)	
— ureter (Fig. 2)	
— ureter (Fig. 3)	
— ureter (Fig. 4)	
— ureter (Fig. 5)	
ureter (Fig. 6)	
— ureter (Fig. 7)	
— ureter (Fig. 8)	392
4-000 (- 6-0/····································	JU-

ILLUSTRATIONS

Rabbit's vas deferens (Fig. 9)	393
vas deferens (Fig. 10)	394
Isolated gall bladder strip of dog (Fig. 11)	395
— gall bladder of dog (Fig. 12)	
Strips of dog's gall bladder (Fig. 13)	395
Dog's gall bladder in situ (Fig. 14)	
Intestine of cat (Fig. 15)	
— of cat (Fig. 16)	
Jejunum of cat (Fig. 17)	
Ileum of cat (Fig. 18)	
Action of papaverin on internal iliac artery of pig (Fig. 19)	
Experiment performed on a cat (Fig. 20)	
Uterus of cat, non-pregnant (Fig. 21)	
— of rat (Fig. 22)	
of cat, non-pregnant (Fig. 23).	
Pig's ureter (Fig. 27)	
— ureter (Fig. 29)	
— ureter (Fig. 31)	
—— ureter (Fig. 32)	
— ureter (Fig. 35)	
ureter (Fig. 36)	407
— ureter (Fig. 37)	
Rabbit's bladder (Fig. 38)	
Loop of jejunum, guinea pig (Fig. 39)	
Strip of gall bladder of cat (Fig. 40)	
Vas deferens of rat (Fig. 41)	
Half of horn of virgin uterus of cat (Fig. 42)	
Three rings of uterine artery of pig (Fig. 43)	
rings of uterine artery of pig (Fig. 44)	
Strip of pregnant uterus of cat (Fig. 50)	
Pig's ureter (Fig. 51)	
Ileum of cat (Fig. 52)	
Intestinal loop of rabbit's jejunum (Fig. 1)	
Experiment. Isolated uterine horn of rat (Fig. 2)	
Ureteral ring preparation (pig) (Fig. 3)	
ring preparation (pig) (Fig. 4)	425
— ring preparation (pig) (Fig. 5)	425
Experiment. Dog, 12 kilo, paraldehyde anesthesia (Fig. 6)	
February 14, 1918. Rabbit, 2.4 kilo, paraldehyde anesthesia (Fig. 7)	
— March 22, 1918. Dog, 6 kilo, paraldehyde anesthesia (Fig. 8)	
Perfusion of isolated heart of cat (Fig. 9)	429
Experiment. January 30, 1918. Dog, 4 kilo, paraldehyde anesthesia (Fig.	
10)	430
Nerve-muscle preparation of gastrocnemius of Rana maculata (Fig. 11)	
	448
Contraction of a frog's gastrocnemius in Ringer's solution containing 1:1500	
Mexican extract (Fig. 1)	463

ILLUSTRATIONS

Contraction of a frog's gastrocnemius in Ringer's solution containing 1: 100 Chinese extract (Fig. 2)	463
— curve of a frog's gastrocnemius after ten minutes in 1:1000 Mexican	
extract (Fig. 3)	464
Blood pressure of a cat of 1.8 kgm. body weight injected with the extract of	
0.36 gram Chinese scorpion sting intravenously (Fig. 4)	469
Action of Mexican extract (1:1200) upon frog's heart perfused with Ringer's	
solution (Fig. 5)	471
of Chinese extract (1:100) upon frog's heart perfused with Ringer's solu-	
tion (Fig. 6)	471
Tracing of a cat's heart under perfusion of Locke's solution with blood	
(Fig. 7)	472
of a cat's heart under perfusion of Locke's solution with blood (Fig. 8)	473
Action of Chinese extract (1:300) on ring of frog's stomach (pyloric half) in	
Ringer's solution (Fig. 9)	476
—— of Mexican extract (1:2500) on strip of kitten's stomach (circular fibers)	
in Locke's solution mixed with blood (Fig. 10)	477
— of Mexican extract (1:5000) on a loop of small intestine of rabbit in	
Tyrode's solution with blood (Fig. 11)	477
— of Chinese extract (1:150) on a loop of scall intestine of rabbit in Ty-	
rode's solution mixed with blood (Fig. 12)	478
— of Mexican extract (1:750) on surviving uterus of guinea-pig (non-	
pregnant) in Locke's solution (Fig. 13)	479
— of Mexican extract (1:1000) on surviving pregnant uterus of rabbit in	
Locke's solution (Fig. 14)	479
— of Mexican extract (1:1500) on circular strip of urinary bladder of cat in	
	480
— of Chinese extract (1:100) on circular strip of urinary bladder of rabbit in	
Ringer's solution (Fig. 16)	480
—— of Chinese extract (1:150) on ring of pig's ureter in Ringer's solution with	
/0· /· · · · · · · · · · · · · · · · · ·	481
of Mexican extract (1:350) on ring of pig's ureter in Ringer's solution	
with urine (Fig. 18)	481