## CONTENTS

## Number 1, March, 1920

<ul> <li>I. The Physiological Action of the Fumes of Iodine. By A. B. Luckhardt, F. C. Koch, W. F. Schroeder, and A. H. Weiland</li></ul>	1 23 29 75 77 81
Number 2, April, 1920	
<ul> <li>VII. An Apparatus for the Exposure of Skin or Mucous Membrane to the Vapor of Toxic Substances, with Observations on Dichlorethylsulphide. By J. A. E. Eyster and Mary E. Maver.</li> <li>VIII. On Optical Isomers. V. The Tropeines. By A. R. Cushny.</li> <li>IX. The Local Anesthetic Properties of Phenyl Methyl Carbinol. By Axel M. Hjort and Charles E. Kaufmann.</li> <li>X. Effect of Atropine on Chloroform Hyperglycemia. By Ellison L. Ross</li> <li>XI. The Effect of Some Antipyretics on the Acuity of Hearing. By D. I. Macht, J. Greenberg and S. Isaacs</li> <li>XII. Comments on the Clotting Efficiency of Thromboplastic Agents. By Frederic Fenger.</li> <li>XIII. Iodine Absorption from the Human Skin. By Norman C. Wetzel and Torald Sollmann.</li> <li>XIV. Drug Perfusion of the Medulla of the Turtle. II. Aconitine, Morphine, Cocaine, Quinine. By A. D. Bush.</li> </ul>	105 129 135 149 167 169
Number 3, May, 1920	
XV. The Reaction of the Cat's Uterus to Strophanthus and Calcium. By Fred Ransom.  XVI. The Point of Attack of Certain Drugs Acting on the Periphery. I. Action on the Bladder. By Charles W. Edmunds and George B. Roth XVII. The Point of Attack of Certain Drugs Acting on the Periphery. II. Action on the Retractor Penis Muscle of the Dog. By Charles W. Edmunds.  XVIII. The Influence of Heavy Metals on the Isolated Frog Heart. By William Salant and Helene Connet.	189 201
XIX. Scientific Proceedings of the American Society for Pharmacology and Experimental Therapeutics.	

## CONTENTS

## Number 4, June, 1920

XX. A Study of the Anurias Occurring in Normal Animals During the Use of the General Anesthetics. By William de B. MacNider	
XXII. IV. The Antagonistic actions of Epinephrin and potassium Chloride on the Tonus and Tonus Waves in the Excised Terrapin Auricles. By Charles M. Gruber.	
XXIII. Further Pharmacologic Studies on Arsphenamine. By Maurice I.	 279
XXIV. Perfusion of the Medulla of the Turtle. III. Epinephrin. By A.	297
XXV. The Mechanism of the Toxic Action of Cyanogen Chloride. By C. I.	301
XXVI. Tyramine as a Morphine Antagonist. By Henry G. Barbour and Lloyd L. Maurer	305
XXVII. Blood Volume and Blood Volume Methods. By Paul D. Lamson and Takeyoshi Nagayama	331
of the Pituitary Gland and on Preliminary Experiments with the Pressor Constituent. By John J. Abel and T. Nagayama	
Number 5, July, 1920	101
XXX. The Action of Chloral on the Pupil. By E. G. Hyatt, Hugh McGuigan and F. A. Rettig.	415
XXXI. The Effect of High Temperature Upon the Action and Toxicity of Digitalis. By Arthur D. Hirschfelder, J. Bicek, F. J. Kucera and W. Hanson	197
XXXII. Continuous Blood Pressure Tracings in Man: An Apparatus. By A. C. Kolls	
XXXIII. An Indirect Method for the Determination of Blood Pressure in the Unanesthetized Dog. By A. C. Kolls	
XXXIV. Quantitative Studies in Chemotherapy. I. The Trypanocidal Action of Antimony Compounds. By Carl Voegtlin and Homer W.	- 40
Smith	<b>4</b> 53
Action of Arsenic Compounds. By Carl Voegtlin and Homer W. Smith	475

Apparatus for liberation of fumes of iodine (Fig. 1)	2
Photographs of sections of the thyroid gland magnified fifty times (Fig. 2)	10
Photographic reproductions of sections of the thyroid glands of the same	
animal (dog 22) (Fig. 3)	10
reproductions of sections of the thyroid gland magnified fifty times	
(Fig. 4)	13
Effect of the intrabracheal administration of the fumes of iodine (Tracing 1)	
facing	16
— of the intrabracheal administration of the fumes of iodine (Tracing 2).	
facing	16
of pilocarpine, atropine and adrenaline upon the tonus waves in the	
terrapin heart (Fig. 1)	<b>25</b>
— of pilocarpine, atropine and adrenaline upon the tonus waves in the	
terrapin heart (Fig. 2)	26
of pilocarpine, atropine and adrenaline upon the tonus waves in the	
terrapin heart (Fig. 3)	26
Saline injection after hemorrhage, compared with the effect of gum acacia	
(Fig. 1)	34
Effect of hypertonic saline after hemorrhage (Fig. 2)	36
Dilution of the blood (Fig. 3)	37
Muscle injury (Fig. 4)	41
Histamine shock. Effect of gum saline (Fig. 5)	45
Effect of intravenous injections in the spinal cat (Fig. 6)	61
A non-clotting blood pressure apparatus (Fig. 1)	<b>76</b>
Apparatus for exposure of skin to toxic vapors (Fig. 1)	97
— for exposure of skin toxic vapors (Fig. 2)	97
for exposure of skin to toxic vapors (Fig. 3)	98
On optical isomers (Fig. 1)	
optical isomers (Fig. 2)	
— optical isomers (Fig. 3)	
optical isomers (Fig. 4)	
optical isomers (Fig. 5)	116
— optical isomers (Fig. 6)	117
— optical isomers (Fig. 7)	
optical isomers (Fig. 8)	
— optical isomers (Fig. 9)	
— optical isomers (Fig. 10)	
Turtle 2. Aconitine, crystalline, 0.005 per cent (Fig. 1)	
2. Morphine sulphate, 0.05 per cent (Fig. 2)	
3. Cocaine sulphate, 0.05 per cent (Fig. 3)	176
6. Quinine sulphate, 0.01 per cent (Fig. 4)	178

Reaction of uterus to strophanthus and calcium (Fig. 1)	182
— uterus to strophanthus and calcium (Fig. 2)	183
— uterus to strophanthus and calcium (Fig. 3)	
uterus to strophanthus and calcium (Fig. 4)	185
— uterus to strophanthus and calcium (Fig. 5)	186
Tracing from bladder of cat (Fig. 1)	
— from bladder of male cat (Fig. 2)	203
Tracings from bladder of cat (Fig. 3)	204
Isolated retractor muscle (Fig. 4)	
Morphine upon the isolated retractor muscle (Fig. 5)	211
Nicotine upon a relatively inactive retractor muscle (Fig. 6)	211
Frog heart perfused for one minute with 1:10,000 copper sulphate in Ringer	
solution minus sodium bicarbonate (Fig. 1)	219
Perfusion of frog heart with 1:1,000 iron citrate in Ringer's solution for one-	
half minute (Fig. 2)	221
with 1:10,000 nickel acetate in Ringer's solution for one minute (Fig. 3)	223
Frog heart perfused for three minutes with 1:10,000 cobalt chloride in Ring-	
er's solution (Fig. 4)	225
Zinc malate, 1:50,000, in Ringer's solution minus bicarbonate (Fig. 5)	
Effect of 1:1,000,000 cadmium acetate in Ringer's solution minus sodium	
bicarbonate when perfused through the frog heart for one minute (Fig. 6)	229
Perfusion with 1:100,000 cadmium acetate in Ringer solution minus sodium	
bicarbonate (Fig. 7)	229
Gradual weakening of heart action by 1:1000 uranium acetate when perfused	
	230
Effect of continuous injection of 4 per cent saligenin in 0.9 per cent NaCl	
upon respiration and blood pressure (Fig. 1A)	266
— of continuous intravenous injection of 4 per cent saligenin solution in a	-00
• • • •	267
Effects of intermittent injections of saligenin intravenously in a dog weigh-	201
	വര
ing 10 kgm. (Figs. 2A, 2B)	
Chrysemys cinera (Fig. 1)	
—, 9 kilos, morphine and chloretone (Fig. 2)	
, 9 kilos, morphine and chloretone (Fig. 2)	
, 25 kilos, morphine and chloretone (Fig. 3), 10 kilos, morphine and chloretone (Fig. 4)	
, 10.3 kilos, ether (Fig. 5)	200
—, 5.5 kilos, ether (Fig. 6)	
Respiratory apparatus for rat (Fig. 1)	
Rat 6, 180 grams (Fig. 2)	911
Effect on respiratory volume of different rats of various doses of tyramine	212
(Fig. 3)	010 010
Effects of large doses of tyramine upon the respiratory volume (Fig. 4)	
of morphine and tyramine injected together in rat 19, 135 grams (Fig. 5)	
of morphine and tyramine injected together in rat 28, 200 grams (Fig. 6)	219

Comparison of effects on respiratory volume of morphine alone with those	
of a simultaneous injection of morphine on one side and tyramine on the	
other side of a rat (Fig. 7)	
Cat, 3.3 kgm., ether (Fig. 1)	
, 2.75 kgm., urethane anesthesia (Fig. 2)	354
Large cat, ether	356
Dog, 5.6 kgm., ether anesthesia (Fig. 4)	
Cat, 3.3 kgm., ether anesthesia being the same animal used for the experi-	
ment illustrated in figure 1 (Fig. 5)	
Effect of boiling an infundibular extract (Armour's Pituitary Liquid) with	
acid (Fig. 6)	
- of heating infundibular extracts (Armour's Pituitary Liquid) with	
acids (Fig. 7)	
Young dog, 5.4 kgm., anesthetic, chloretone (Fig. 8)	
Guinea-pig's uterus (Fig. 9)	
Virgin guinea-pig's uterus (Fig. 10)	
— guinea-pig's uterus (Fig. 11).	
Action of 2 drops of a diluted infundibular extract (Armour) on the virgin	
guinea-pig's uterus (Fig. 12)	
Large cat, ether (Fig. 13)	
Dog, 5.5 kgm., chloretone in oil intraperitoneally (Fig. 15)	
Young dog, 6 kgm., anesthetic not noted at time (Fig. 16)	
dog, 4.9 kgm., chloretone in oil intraperitoneally (Fig. 17)	
— dog, 3.95 kgm., anesthetic not noted (Fig. 18)	
Dog, 6.9 kgm., ether (Fig. 19)	
Young dog, 4.2 kgm.; anesthetic, chlorotone intraperitoneally (Fig. 20)	
Dog, 7.9 kgm., ether (Fig. 21)	
Young dog, 5.55 kgm., ether (Fig. 1)	
Animal as in figure 1 (Fig. 2)	
Comparative action on the virgin guinea-pig's uterus of 0.37 per cent solu-	
tion of the unaltered proteoses (Fig. 3)	
Dog, ♂, 9.2 kgm. (Fig. 4)	
Animal as in figure 3 (Fig. 5)	
Young dog, 3.2 kgm., ether (Fig. 6)	410
Same animal as in figure 6 (Fig. 7)	
Effects of chloroform extracts after hydrolysis of the proteoses on the virgin	
guinea-pig's uterus (Fig. 8)	411
Cat, Q, 2.63 kgm., ether (Fig. 9)	
Effect of equal quantities of the alcohol extract of the three proteoses on	
the virgin guinea-pig's uterus (Fig. 10)	414
Diagram of apparatus for continuous blood-pressure tracings (Fig. 1)	435
Apparatus for continuous blood pressure tracings in man (Fig. 2)	436
Electromagnetic valve—actual size (Fig. 3)	
Continuous blood pressure tracing (Fig. 4)	
— blood pressure tracing (Fig. 5)	
Cross section of dog's thigh (lower diagram shows the relation of cuff to	
leg) (Fig. 1)	445

Drawing of the cuff (Fig. 2)	446
Sphymograph (approximately actual size) (Fig. 3)	447
The sphygmomanometer (Fig. 4)	
Record of a determination on a normal unanesthetized dog (Fig. 5)	
Comparison of direct and indirect methods on the anesthetized animal	
(Fig. 6)	450
Quantitative studies in chemotherapy. I (Fig. 1)	459
studies in chemotherapy. I. (Fig. 2)	459
- studies in chemotherapy. I. (Fig. 3)	
studies in chemotherapy. I. (Fig. 4)	<b>46</b> 0
studies in chemotherapy. I. (Fig. 5)	461
studies in chemotherapy. I. (Fig. 6)	461
studies in chemotherapy. I. (Fig. 7)	462
studies in chemotherapy. I. (Fig. 8)	465
studies in chemotherapy. I. (Fig. 9)	466
studies in chemotherapy. I. (Fig. 10)	467
Quantitative studies in chemotherapy. II. (Chart 1)	476
studies in chemotherapy. II. (Chart 2)	480
studies in chemotherapy. II. (Chart 3)	481
studies in chemotherapy. II. (Chart 4)	482
studies in chemotherapy. II. (Chart 5)	488
studies in chemotherapy. II. (Chart 6)	489