Per. V.26 1925-26

650

Downloaded from jpet.aspetjournals.org at ASPET Journals on April 17, 2024

## CONTENTS

## Number 1, August, 1925

I. Slowing and Block in the Heart Under Digitalis in Animals. By V. R. Cushy and K. Y. Yu	1
II. Effects of Quinidine on the Auricular Irritability and Conduction in the	1
	19
III. Studies of Intoxication. III. The Action of Ethylene. By Bessie M.	
Davidson	27
•••	33
V. Studies of Intoxication. V. The Action of Ethyl Chloride. By Bessie M.	-
	37
VI. Studies of Intoxication. VI. The Action of Methyl Ether. By Bessie	
	43
VII. The Excretion of Morphin into the Stomach. By Robert A. Hatcher and David Davis	49
VIII. The Salicylates. XV. Liberation of Salicyl from and Excretion of	70
	61
IX. The Salicylates. XVI. Liberation of Salicyl from and Excretion of	
Methyl Salicylate, with a Note on the Irregular Toxicity of the Ester in	
Man. By P. J. Hanzlik and N. E. Presho	71
Number 2, September, 1925	
X. The Effect of Ephedrine on Experimental Shock and Hemorrhage. By	
K. K. Chen	83
XI. A Preliminary Paper Concerning the Toxic Effect of Certain Alcoholic	
Beverages for the Kidney of Normal and Naturally Nephropathic Dogs.	0=
By Wm. deB. MacNider	97
Davidson	05
XIII. Studies of Intoxication. VIII. The Influence of Oxygen. By Bessie	•
M. Davidson	11
XIV. An Experimental Study of Certain Benzyl Compounds with Especial	
Attention to the Importance of Substitution in the Benzyl Nucleus and	00
the Significance of the Side Chains. Herman H. Jensen	23
Hamilton Crawford	71
XVI. II. The Action of Sparteine Sulphate on Experimental Fibrillation of	_
the Auricles. By J. Hamilton Crawford	Q1

## Number 3, October, 1925

XVII. A Proposed Standardized Method for the Therapeutic Study of Compounds in Experimental Rabbit Syphilis. By G. E. Wakerlin, W. F. Lorenz, and A. S. Loevenhart	
tive Relation to Weight-Changes, and to Vitamine Storage, in Adult Pigeons. By J. D. Pilcher and Torald Sollmann	215 227 233
Number 4, November, 1925	
XXIV. On the Inactivation of Atropine Sulphate by Rabbit Serum. By Jean La Barre	259
XXV. Physiological Action of Furane. By Elizabeth M. Koch and Meyer H. Cahan	
XXVI. The Effect of Sodium Arsenite on the Blood Sugar Concentration of the Rabbit and Dog. By H. B. van Dyke	287
By E. G. Holmes	
Number 5, December, 1925	
•	
XXIX Acute Cocaine Poisoning, its Prophylaxis and Treatment in Laboratory Animals. By A. L. Tatum, A. J. Atkinson and K. H. Collins XXX. On the Active Principles of the Pituitary Extract. By H. H. Knaus. XXXI. The Action of Certain Drugs and Ions on the Rat's Uterus. By H. H.	
Knaus and A. J. Clark	
Clark, H. H. Knaus and A. S. Parkes.  XXXIII. A Comparative Study of Hypnotics of the Barbituric Acid Series.  Pro Corl Nickey, John A. Hinging and Honor C. Spread.	
By Carl Nielsen, John A. Higgins and Henry C. Spruth	
Number 6, January, 1926	
, ,	
XXXV. Chlorocodon Whiteii: Its Constituents and Their Pharmacological Actions. By Walter J. Dilling	

CONTENTS v

XXXVI. The Influence of Histamine on the Intestinal Secretion of the Dog.	
By W. Koskowski	13
XXXVII. Tannin Occurring in the Kino Eucalyptus Calophylla. By	
Alexander McGookin and I. M. Heilbron 42	21
XXXVIII. Bacterial Chemotherapy with Special Reference to Mercury	
Dyestuffs. By George W. Raiziss, M. Severac and John C. Moetsch 44	17
XXXIX. Chemotherapy of Bacterial Infections. I. Action of Acriflavine	
Gentian-Violet and Mercurochrome in Experimental Bacterial In-	
fections. By Ernest Linwood Walker and Marian A. Sweeney 46	31
XL. Morphine Miosis. By V. E. Henderson and R. W. Graham 46	39
XLI. A Study of the Narcotic Action of Propylene. By J. T. Halsey,	
Chapman Reynolds and W. A. Prout	79

## ILLUSTRATIONS

missions of the ventricle from partial A-V block (Fig. 1)	6
Half-rhythm at a later stage than figure 1 (Fig. 2)	7
Marked A-V block, the ventricle beating once to the auricle's 4 to 7 beats	•
(Fig. 3)	8
sinus slowing with no A-V block in the first half of the tracing (Fig. 4)	9
Partial block, with spontaneous ventricular beat interpolated during failure	9
	10
of conduction (Fig. 5)	10
Repeated extrasystoles in the ventricle during pauses in the auricle from	12
inhibition (Fig. 6)	12
The auricle beating slowly from inhibition due to scillaren, while the ventricle	10
has assumed more rapid spontaneous rhythm (Fig. 7)	13
Absence of fibrillation when maximal faradic stimuli are applied to the right	00
auricle of the terrapin (Fig. 1)	20
Response of both auricles to interrupted stimuli (Fig. 2A)	22
— of both auricles to interrupted stimuli (Fig. 2B)	23
Depression of irritability after quinidine (Fig. 3)	24
Graph showing reaction times with choice under different concentrations of	
ethylene (Fig. 1)	29
showing reaction times with choice under different concentrations of	
ethyl chloride (Fig. 1)	<b>4</b> 0
showing reaction times with choice during successive administrations of	
2.5 per cent ethyl chloride (Fig. 2)	41
showing reaction times with choice under different concentrations of	
methyl ether (Fig. 1)	46
Liberation of salicyl from methyl salicylate in "buffer" mixtures at 38°C.	
at the end of one hour, six and twenty-four hours with and without	
pancreatine (Fig. 1)	72
Effect of ephedrine in histamine shock (Fig. 1)	85
— of ephedrine in peptone shock (Fig. 2)	86
— of ephedrine in anaphylactic shock (Fig. 3)	87
— of ephedrine in surgical and traumatic shock (Fig. 4)	88
— of ephedrine in hemorrhage (Fig. 5)	89
— of ephedrine on peripheral circulation in histamine shock (Fig. 6)	90
— of ephedrine on the heart in histamine shock (Fig. 7)	91
Graph showing simple reaction times after cessation of the inhalation of	
•	106
showing the effect of 0.5 gram of caffeine on reaction times with choice	
(Fig. 2)	107

Graph showing reaction times with choice under 2.5 per cent ethyl chloride	
(Fig. 3)	108
—— showing reaction times with choice (Fig. 4)	109
showing reaction times with choice (Fig. 1)	114
showing the effect of 40 per cent nitrous oxide + 12 per cent oxygen fol-	
lowed by 40 per cent nitrous oxide + 28 per cent oxygen on reaction times	
with choice (Fig. 2)	116
—— showing the effect of 33 per cent acetylene $+$ 20 per cent oxygen, and 33	
per cent acetylene + 33 per cent oxygen on reaction times with choice	
(Fig. 3)	117
ethyl chloride in air and 3.36 per cent ethyl chloride in oxygen (Fig. 4)	120
Blood pressure and respiration. Dialcohol p-cresol. Rabbit—ether anes-	
thesia (Fig. 1)	136
pressure and respiration. Alcohol salicylaldehyde. Rabbit—ether	
anesthesia (Fig. 2)	137
Antispasmodic action. Alcohol p-oxy-benzaldehyde. Intestinal segments—	
rabbit	146
Diagram (Fig. 5)	
Myocardiograms from the auricle and ventricle and blood pressure tracing	100
(Fig. 1)	183
from the auricle and ventricle and blood pressure tracing (Fig. 2)	184
Healing power of neoarsphenamine in experimental rabbit syphilis (Fig. 1)	189
— power of sodium salt of 3, 3', 4, 4'-tetraamino-arsenobenzene tetra-	100
methylene sulphinic acid in experimental rabbit syphilis (Fig. 2)	190
Average percentage weight of pigeons by groups according to yeast extract	200
doses; polished rice diet (Fig. 1)	205
Percentage weight loss or gain at end of twenty-one days of yeast extract	
administration, arranged by groups, according to dosage (Fig. 2)	206
Average percentage weight of pigeons by groups during the vitamine and	
avitamine periods; polished rice diet throughout (Fig. 3)	207
— percentage weight of pigeons by groups during vitamine and avitamine	20.
periods; polished rice diet throughout (Fig. 4)	208
Mesenteric vein ring (sheep), showing rhythmic movements (Fig. 1)	
— vein ring (sheep) (Fig. 2)	218
vein rings (sheep) (Figs. 3 and 4)	
— vein rings (ox) (Figs. 5 and 6).	
	222
— vein ring (ox) (Fig. 1)	228
Small intestine (rabbit) (Fig. 2)	
Intrapulmonary bronchus ring (sheep) (Fig. 3)	
Bladder (rabbit) (Fig. 4)	
Spleen (rabbit) (Fig. 5)	
Chart showing distribution of deaths, within four weeks, following adrenal-	
ectomy (Fig. 1)	247
	248

Chart showing the effect of 0.4 mgm. pilocarpine nitrate alone on normal	
intestine; the effect of same dose of pilocarpine nitrate on pieces of	
intestine treated previously with 0.01 mgm. atropine; and the effect of	
same dose of pilocarpine after 0.008 mgm. atropine (Fig. 1)	264
- showing average tonic variations after addition of 0.4 mgm. of pilocar-	
pine nitrate to Ringer solution in which were suspended pieces of intestine	
treated previously by different quantities of extracted atropine (Fig. 2).	265
showing tonic variation after addition of 0.4 mgm. pilocarpine nitrate to	
Ringer solution in which are suspended intestinal pieces treated previ-	
ously by different quantities of atropine solution extracted from serum	
(Fig. 3)	26 <b>6</b>
of the relation of inactivation of atropine with the duration of contact	
with serum (Fig. 4)	272
Abscissa, temperature to which the mixture serum-atropine was exposed for	
	272
Cat, weighed 3.5 kgm., intravenous injection of extract equal to 1 mgm. of	
moist gland (Fig. 1)	338
, weight 2.5 kgm. (Fig. 2)	340
	342
—— cat, weight 3 kgm., intramuscular injection of extract equal to 1 mgm.	
(0.2 cc. fluid) of moist gland (Fig. 6)	
Cat, weight 3.5 kgm. (Figs. 7 and 8)	
The action of adrenalin on the rat's uterus (Fig. 1)	348
action of adrenalin and pituitary extract on conduction in the rat's	
uterus (Fig. 2)	
action of pituitary extract on the rat's uterus (Fig. 3)	
Movements of same uterus (1) in situ and (2) isolated (Fig. 1)	
— of the same uterus (1) isolated and (2) in situ (Fig. 2)	362
Diagram showing method of recording contractions of uterus from three leads	
simultaneously (Fig. 3)	
Movements of same uterus in situ and isolated (Fig. 4)	
	368
Results of a comparative study on hypnotics of the barbituric acid series	001
(Charts 1 and 2)	
Effect produced by stimulation of right vagus before and after eserine (Fig. 1)	
Sections from an adrenaline curve before section of vagi and eserine (Fig. 2)	
from adrenaline curve after section of vagi nerves (Fig. 3)	389
Adrenaline curve after section of vagi and atropine (Fig. 4)	
Inhibition produced by 1 cc. 1:10,000 adrenaline given intravenously (Fig. 5).  —— produced by 1 cc. 1:10,000 adrenaline plus vagus stimulation (Fig. 6)	
	390 391
— produced by adrenaline (Fig. 7)	
Chlorocodon Whiteii. Action of volatile oil (1:3000) on rabbit intestine (Fig.	
1)	
Action of chlorocodon glucoside 0.1 per cent on frog heart (Fig. 2)	
— of chlorocodon glucoside on frog heart (Fig. 3)	
of chlorocodon glucoside 1 per cent on frog's hypothossus (Fig. 4)	