Per. V.26 1925-26

650

Downloaded from jpet.aspetjournals.org at ASPET Journals on April 19, 2024

## CONTENTS

#### Number 1, August, 1925

I. Slowing and Block in the Heart Under Digitalis in Animals. By V. R. Cushy and K. Y. Yu	1
II. Effects of Quinidine on the Auricular Irritability and Conduction in the	1
	19
III. Studies of Intoxication. III. The Action of Ethylene. By Bessie M.	
Davidson	27
•••	33
V. Studies of Intoxication. V. The Action of Ethyl Chloride. By Bessie M.	-
	37
VI. Studies of Intoxication. VI. The Action of Methyl Ether. By Bessie	
	43
VII. The Excretion of Morphin into the Stomach. By Robert A. Hatcher and David Davis	49
VIII. The Salicylates. XV. Liberation of Salicyl from and Excretion of	70
	61
IX. The Salicylates. XVI. Liberation of Salicyl from and Excretion of	
Methyl Salicylate, with a Note on the Irregular Toxicity of the Ester in	
Man. By P. J. Hanzlik and N. E. Presho	71
Number 2, September, 1925	
X. The Effect of Ephedrine on Experimental Shock and Hemorrhage. By	
K. K. Chen	83
XI. A Preliminary Paper Concerning the Toxic Effect of Certain Alcoholic	
Beverages for the Kidney of Normal and Naturally Nephropathic Dogs.	0=
By Wm. deB. MacNider	97
Davidson	05
XIII. Studies of Intoxication. VIII. The Influence of Oxygen. By Bessie	•
M. Davidson	11
XIV. An Experimental Study of Certain Benzyl Compounds with Especial	
Attention to the Importance of Substitution in the Benzyl Nucleus and	00
the Significance of the Side Chains. Herman H. Jensen	23
Hamilton Crawford	71
XVI. II. The Action of Sparteine Sulphate on Experimental Fibrillation of	_
the Auricles. By J. Hamilton Crawford	Q1

## Number 3, October, 1925

XVII. A Proposed Standardized Method for the Therapeutic Study of Compounds in Experimental Rabbit Syphilis. By G. E. Wakerlin, W. F. Lorenz, and A. S. Loevenhart	
<ul> <li>XX. The Pharmacology of the Isolated Vein Ring. By K. J. Franklin</li></ul>	
XXIII. The Influence of the Adrenals on the Toxicity of Morphine. By J. M. Rogoff and Jos. DeNecker	
Number 4, November, 1925	
XXIV. On the Inactivation of Atropine Sulphate by Rabbit Serum. By	
Jean La Barre	
H. Cahan	
XXVII. A Contribution to the Study of the Metabolism of Salicylic Acid.  By E. G. Holmes	
XXVIII. Skin Absorption of Certain Gases. By D. C. Walton and M. G. Witherspoon	j
Number 5, December, 1925	
XXIX Acute Cocaine Poisoning, its Prophylaxis and Treatment in Labora-	
tory Animals. By A. L. Tatum, A. J. Atkinson and K. H. Collins 325  XXX. On the Active Principles of the Pituitary Extract. By H. H. Knaus. 337  XXXI. The Action of Certain Drugs and Ions on the Rat's Uterus. By H. H.	
Knaus and A. J. Clark	,
Clark, H. H. Knaus and A. S. Parkes	)
By Carl Nielsen, John A. Higgins and Henry C. Spruth	
Number 6, January, 1926	
, ,	
XXXV. Chlorocodon Whiteii: Its Constituents and Their Pharmacological Actions. By Walter J. Dilling	,

#### CONTENTS v

# ILLUSTRATIONS

missions of the ventricle from partial A-V block (Fig. 1)	6
Half-rhythm at a later stage than figure 1 (Fig. 2)	7
Marked A-V block, the ventricle beating once to the auricle's 4 to 7 beats	•
(Fig. 3)	8
	9
Partial block, with spontaneous ventricular beat interpolated during failure	
of conduction (Fig. 5)	10
Repeated extrasystoles in the ventricle during pauses in the auricle from	
inhibition (Fig. 6)	12
The auricle beating slowly from inhibition due to scillaren, while the ventricle	
has assumed more rapid spontaneous rhythm (Fig. 7)	13
Absence of fibrillation when maximal faradic stimuli are applied to the right	
auricle of the terrapin (Fig. 1)	20
Response of both auricles to interrupted stimuli (Fig. 2A)	22
— of both auricles to interrupted stimuli (Fig. 2B)	23
Depression of irritability after quinidine (Fig. 3)	24
Graph showing reaction times with choice under different concentrations of	
ethylene (Fig. 1)	29
showing reaction times with choice under different concentrations of	
ethyl chloride (Fig. 1)	40
showing reaction times with choice during successive administrations of	
2.5 per cent ethyl chloride (Fig. 2)	41
showing reaction times with choice under different concentrations of	
methyl ether (Fig. 1)	46
Liberation of salicyl from methyl salicylate in "buffer" mixtures at 38°C.	
at the end of one hour, six and twenty-four hours with and without	
pancreatine (Fig. 1)	72
Effect of ephedrine in histamine shock (Fig. 1)	85
— of ephedrine in peptone shock (Fig. 2)	86
— of ephedrine in anaphylactic shock (Fig. 3)	87
of ephedrine in surgical and traumatic shock (Fig. 4)	88
— of ephedrine in hemorrhage (Fig. 5)	89
of ephedrine on peripheral circulation in histamine shock (Fig. 6)	90
— of ephedrine on the heart in histamine shock (Fig. 7)	91
Graph showing simple reaction times after cessation of the inhalation of	
pure nitrous oxide to unconsciousness (Fig. 1)	106
(Fig. 2)	107

Graph showing reaction times with choice under 2.5 per cent ethyl chloride	
	108
showing reaction times with choice (Fig. 4)	109
showing reaction times with choice (Fig. 1)	114
- showing the effect of 40 per cent nitrous oxide + 12 per cent oxygen fol-	
lowed by 40 per cent nitrous oxide + 28 per cent oxygen on reaction times	
with choice (Fig. 2)	116
showing the effect of 33 per cent acetylene + 20 per cent oxygen, and 33	
per cent acetylene + 33 per cent oxygen on reaction times with choice	
(Fig. 3)	117
ethyl chloride in air and 3.36 per cent ethyl chloride in oxygen (Fig. 4)	120
Blood pressure and respiration. Dialcohol p-cresol. Rabbit—ether anes-	
thesia (Fig. 1)	136
— pressure and respiration. Alcohol salicylaldehyde. Rabbit—ether	
anesthesia (Fig. 2)	137
Antispasmodic action. Alcohol p-oxy-benzaldehyde. Intestinal segments—	
	146
	147
Diagram (Fig. 5)	160
Myocardiograms from the auricle and ventricle and blood pressure tracing	100
	183
from the auricle and ventricle and blood pressure tracing (Fig. 2)	184
Healing power of neoarsphenamine in experimental rabbit syphilis (Fig. 1)	189
— power of sodium salt of 3, 3', 4, 4'-tetraamino-arsenobenzene tetra-	200
methylene sulphinic acid in experimental rabbit syphilis (Fig. 2)	190
Average percentage weight of pigeons by groups according to yeast extract	
	205
Percentage weight loss or gain at end of twenty-one days of yeast extract	
administration, arranged by groups, according to dosage (Fig. 2)	206
Average percentage weight of pigeons by groups during the vitamine and	
	207
— percentage weight of pigeons by groups during vitamine and avitamine	
	208
	217
	218
	219
	221
vein rings (sheep) (Figs. 7, 8, 9, 10, 11 and 12)	222
— vein ring (ox) (Fig. 1)	228
Small intestine (rabbit) (Fig. 2)	
ntrapulmonary bronchus ring (sheep) (Fig. 3)	
Bladder (rabbit) (Fig. 4)	
	231
Chart showing distribution of deaths, within four weeks, following adrenal-	
<u> </u>	247
	248

Chart showing the effect of 0.4 mgm. pilocarpine nitrate alone on normal	
intestine; the effect of same dose of pilocarpine nitrate on pieces of	
intestine treated previously with 0.01 mgm. atropine; and the effect of	004
same dose of pilocarpine after 0.008 mgm. atropine (Fig. 1)	204
- showing average tonic variations after addition of 0.4 mgm. of pilocar-	
pine nitrate to Ringer solution in which were suspended pieces of intestine	265
treated previously by different quantities of extracted atropine (Fig. 2).  — showing tonic variation after addition of 0.4 mgm. pilocarpine nitrate to	200
Ringer solution in which are suspended intestinal pieces treated previ-	
ously by different quantities of atropine solution extracted from serum	
(Fig. 3)	26 <b>6</b>
of the relation of inactivation of atropine with the duration of contact	20 <b>0</b>
with serum (Fig. 4)	272
Abscissa, temperature to which the mixture serum-atropine was exposed for	
• = = = = = = = = = = = = = = = = = = =	272
Cat, weighed 3.5 kgm., intravenous injection of extract equal to 1 mgm. of	
moist gland (Fig. 1)	338
, weight 2.5 kgm. (Fig. 2)	
Virgin cat, weight 2 kgm. (Figs. 3, 4 and 5)	
cat, weight 3 kgm., intramuscular injection of extract equal to 1 mgm.	
(0.2 cc. fluid) of moist gland (Fig. 6)	344
Cat, weight 3.5 kgm. (Figs. 7 and 8)	
The action of adrenalin on the rat's uterus (Fig. 1)	
uterus (Fig. 2)	350
action of pituitary extract on the rat's uterus (Fig. 3)	
Movements of same uterus (1) in situ and (2) isolated (Fig. 1)	
— of the same uterus (1) isolated and (2) in situ (Fig. 2)	
Diagram showing method of recording contractions of uterus from three leads	
simultaneously (Fig. 3)	364
Movements of same uterus in situ and isolated (Fig. 4)	365
Records of the movements of three different isolated uteri (Fig. 5)	368
Results of a comparative study on hypnotics of the barbituric acid series	
(Charts 1 and 2)	381
Effect produced by stimulation of right vagus before and after eserine (Fig. 1)	
Sections from an adrenaline curve before section of vagi and eserine (Fig. 2)	
from adrenaline curve after section of vagi nerves (Fig. 3)	
Adrenaline curve after section of vagi and atropine (Fig. 4)	
Inhibition produced by 1 cc. 1:10,000 adrenaline given intravenously (Fig. 5).	
produced by 1 cc. 1:10,000 adrenaline plus vagus stimulation (Fig. 6)	
— produced by adrenaline (Fig. 7)	
—— produced by adrenaline plus vagus stimulation (Fig. 8)	
Chlorocodon Whiteii. Action of volatile oil (1:3000) on rabbit intestine (Fig.	
1)	
Action of chlorocodon glucoside 0.1 per cent on frog heart (Fig. 2)	
of chlorocodon glucoside on frog heart (Fig. 3)	
— of chlorocodon glucoside 1 per cent on frog's hyoglossus (Fig. 4)	409