CONTENTS

Number 1, February, 1926

· · · · · · · · · · · · · · · · · · ·	
Experiments with Anesthetic Gases. By W. Easson Brown and Velyien	
E. Henderson	1
of Its Action. By Melvin Dresbach and Kenneth C. Waddell	9
I. The Effects of General Anesthesia on the Muscular Activity of the	
Gastro-Intestinal Tract. A Study of Ether, Chloroform, Ethylene and Nitrous-Oxide. By G. H. Miller	41
7. The Acute Toxicity of Ephedrine. By K. K. Chen	61
The Effect of Repeated Administration of Ephedrine. By K. K. Chen	61
I. The Effect of Ephedrine on Digestive Secretions. By K. K. Chen	87
Number 2, March, 1926	
II. Comparative Studies of Propylene, Ethylene, Nitrous Oxide and Ether.	
By Chapman Reynolds	93
Studies of Acute Anilin Poisoning. By A. G. Young, C. W. Muehlberger,	
and W. J. Meek	101
7. Toxicological Studies of Anilin and Anilin Compounds. II. Hemato-	
logical Studies of Anilin Poisoning. By A. G. Young	125
logical and Hematological Studies of Acetanilid Poisoning. By A. G.	
Young and J. A. Wilson	133
I. Effects of Carminative Volatile Oils on the Muscular Activity of the	140
Stomach and Colon. By O. H. Plant and G. H. Miller	149
tives. V. Alkyl-, Hydroxy-, and Keto-Acids. By Ralph C. Corley	
with William C. Rose	165
Number 3, April, 1926	
III. Persistence of Action of Physostigmin and the Atropin-Physostigmin	
Antagonism in Animals and in Man. By Soma Weiss	181
IV. Isoamyl Ethyl Barbituric Acid (Amytal), its Use as an Intravenous Anesthetic. By Irvine H. Page and Pol Coryllos	120
V. Prevention of the Edema of Paraphenylenediamine by Drugs Acting on	100
the Adrenals. By M. L. Tainter	201
VI. The Effect of Iodides on the Nitrogen Partition. By G. P. Grabfield	231
and A. M. Prentiss	201
Experimental Therapeutics	235
VIII. Arthur Robertson Cushny and Pharmacology	265

Number 4, May, 1926

· · ·	
XIX. Observations on the Action of Digitalis on the Frog Heart and its Modification by Quinidine. By McKeen Cattell	287
•	299
Vagus of the Fowl. By O. S. Gibbs	319 327
XXIII. XVII. The Effect of Adrenalin and other Vaso-Motor Drugs upon the Cerebral Blood Vessels. By Charles M. Gruber and Samuel J.	
Roberts	
Numbers 5 and 6, June, 1926	
	355
XXVI. Effects of Morphine and Some Other Opium Alkaloids on the Muscular Activity of the Alimentary Canal. I. Action on the Small Intestine in Unanesthetized Dogs and Man. By O. H. Plant and G. H. Miller	361
XXVII. The Toxicity of Mercuric Salicylate for Rabbits. By G. E. Wakerlin and A. S. Loevenhart	385
· · · · · · · · · · · · · · · · · · ·	389
XXIX. Plexus-free Preparations of the Small Intestine. A Study of their Rhythmicity and of their Response to Drugs. By Herbert S. Gasser XXX. Lymph Alterations Following Arsenic Injections. By W. F. Petersen	395
	411
By Eleanor M. Scarborough	42 1
XXXII. The Pharmacological Action of Sparteine and Some Related Alkaloids. I. Sparteine and Oxysparteine. By Reginald St. A. Heathcote	431
	449
XXXIV. Biological Significance of Cystine and Glutathione. I. On the Mechanism of the Cyanide Action. By Carl Voegtlin, J. M. Johnson and	
H. A. Dyer	467

ILLUSTRATIONS

Ether anesthesia (Fig. 1)	46
Chloroform anesthesia (Fig. 2)	48
Ethylene anesthesia (Fig. 3)	49
Nitrous-oxide anesthesia (Fig. 4)	51
Effects of anesthesia on a denervated Thiry-Vella loop of ileum (Fig. 5)	53
Comparative magnitudes of M.L.D. of ephedrine sulphate in rabbits by dif-	
ferent methods of administration (Fig. 1)	67
Comparative magnitudes of M.L.D. of ephedrine sulphate in different species	
of animals by intravenous injection (Fig. 2)	67
Effect on respiration and circulation of an injection of a M.L.D. of ephedrin	
sulphate (Fig. 3)	71
Fibrillation of rabbit's heart by a large dose of ephedrine sulphate with	
Langendorff-Locke's method of perfusion (Fig. 4)	72
The change of electrocardiogram after the M.L.D. intravenously in a rabbit	
(Fig. 5)	74
Effect of ephedrine on blood pressure after its repeated administration	
(Fig. 1)	83
Effect of ephedrine on Pavlov pouch preparation (no. 3) in comparison with	
that of a meal (Fig. 1)	90
Effect of intravenous injection of 0.1 gram anilin per kilo in female dog	
(Fig. 1)	104
Effect of intravenous injection of 0.1 gram anilin per kilo body weight in	
male dog following the section of vagi (Fig. 2)	105
Effect of intravenous injection of 0.1 gram anilin per kilo into a male dog fol-	
lowing section of the vagi and intravenous administration of 3 mgm. of	
	106
Myocardiograph record showing the terminal effect of intravenous injection	
of anilin in a dog (Fig. 4)	107
Tonograph tracing showing effect of intravenous injection of anilin in a	
dog (Fig. 5)	110
Experiment 3 after third injection of 0.1 gram of anilin in a 10 per cent solu-	
tion of the acetate (Fig. 6)	111
Experiment 3, lead 2 (Fig. 7)	113
Experiment 4. First effects after intravenous injection of 0.1 gram per kilo	
of anilin in the form of acetate (Fig. 8)	116
Experiment 9. Gradual development of right bundle branch block after	
anilin (Fig. 9)	117
Experiment 4. After third injection. Complete heart block (Fig. 10)	118
Effect produced by injecting intravenously 24 cc. of an 0.8 per cent solution	
of acetanilid in 2 per cent acetic acid; before and after paralyzing the	
vagi with atropine (Fig. 1)	137

Effect of injecting intravenously 10 cc. of a 10 per cent solution of anilin	
HCl (Fig. 2)	138
Same dog as figure 1, showing terminal stage of the experiment (Fig. 3) Injection of 25 cc. of 10 per cent anilin. Terminal effects of injection (Figs.	139
4 (a) and (b))	140
Normal E. K. G. after injection of 24 cc. of 35 per cent alcohol, and after	140
injection of 24 cc. of 5 per cent acetanilid in 35 per cent alcohol (Figs.	
5 (a) and (b))	142
Later effects, auricles quiescent (Fig. 5 (c))	143
Auricles recovered (Fig. 5 (d))	
Experiment K-12 (Fig. 1)	155
Experiment K-21 (Fig. 2)	
Experiment K-18 (Fig. 3)	
Experiment K-31 (Fig. 4)	
Experiment K-36 (Fig. 5)	
Experiment K-41 (Fig. 6)	
Experiment K-45 (Fig. 7)	
Increased output of epinephrine as indicated by marked dilatation of sen-	
sitized pupil in cat 356 in which the edema of paraphenylenediamine	
was prevented by nicotine (Fig. 1)	218
Increased output of epinephrine as indicated by marked dilatation of sen-	
sitized pupil in rabbit 327 in which the edema of paraphenylenediamine	
was prevented by picrotoxin (Fig. 2)	223
Increased output of epinephrine as indicated by marked dilatation of sen-	
sitized pupil in cat 357 in which edema of paraphenylenediamine was	
prevented by strychnine (Fig. 3)	225
	263
Graph showing tracing obtained by perfusing heart with a 1:50 dilution of	
	289
Graph showing record of ventricular contractions to illustrate the gradual	
diminution of the alternate beat under digitalis perfusion (Fig. 2)	290
Graph showing record similar to figure 1, except that the ventricle was per-	
	292
Graph showing typical tracing of developing "block" in a heart perfused	
simultaneously with quinidine and digitalis (Fig. 4)	294
Graph showing blood pressure changes on stimulation of splanchnic nerve	
(Fig. 1)	303
Graph showing blood pressure changes on stimulation of splanchnic nerve	
(Fig. 2)	304
Graph showing tracings of contraction of the vagina on hypogastric nerve	
20111111111111111111111111111111111111	307
Graph showing tracings of the contraction of the vagina on hypogastric	
nerve stimulation in a decapitated and curarized cat (Fig. 4)	308
Graph showing tracings of changes in the volume of the nasal cavity on	
stimulation of the cervical sympathetic (Fig. 5)	312
Graph showing tracings of changes in the volume of the nasal cavity on	
stimulation of the cervical sympathetic in the cat anesthetized by par-	
aldehyde (Fig. 6)	313

Graph showing blood pressure changes on stimulation of superior mesenteric	
nerves (Fig. 7)	
Graph showing effects of vagus stimulation on the heart and blood pressure	
	321
Graph showing rise in blood pressure caused by an injection of 1 mgm. of	
	321
Graph showing the primary and secondary rise of blood pressure caused by	
an injection of physostigmine (Fig. 3)	321
Graph showing the effect of vagus stimulation, after an injection of phy-	
sostigmine, on the heart rate (Fig. 4)	321
Graph showing the effect of an injection of atropine on a heart slowed by	
physostigmine and vagus stimulation (Fig. 5)	
	322
Graph showing the negative results of an injection of pilocarpine on the	
heart and blood pressure after an injection of physostigmine (Fig. 7)	
Graph showing a continuation of figure 7) (Fig. 8)	324
Graph showing rabbit heart perfused with oxygenated Locke's solution	
modified by the addition of blood 2 per cent, at a pressure of 72 mm.	
of mercury, at a temperature of 37.5°C. (Fig. 1)	
Graph showing same heart as used in figure 1 (Fig. 2)	330
Graph, same as figure 2, except that the heart of another animal was used	990
(Fig. 3)	
Dog 20 kgm. ether anesthesia (Fig. 4). Graphs showing records of a dog's brain perfused through the basilar artery	991
with carotids ligated (Figs. 1-9)	246
Graph showing one cc. saturated solution of "luminal" in blood-Ringer's	340
solution (Fig. 1)	350
Graph showing use of Ringer's solution pH 7.6 modified by the addition of	000
20 per cent defibrinated autogenous blood (Fig. 2)	351
Graph showing use of Ringer's solution pH 7.6 (Fig. 3)	
Graph showing use of Ringer's solution pH 7.6 modified by addition of 20	001
	352
Graph showing tracing of dog IA22, weight 9 kgm., six hours after injection	
(Fig. 1)	368
Graph showing tracing of dog IA5, weight 20 kgm., twenty minutes after	
	368
Graph showing tracing of dog IA6, weight 23 kgm., during first hour after	
	369
Graph showing tracing of dog IA22, twelve hours after injection (Fig. 4)	370
Graph showing tracing of dog 1A6, weight 23 kgm., fifteen minutes after	
injecting 0.2 mgm. atropine sulphate per kilogram (Fig. 5)	374
Graph showing tracing of dog 1A11, weight 17 kgm. (Fig. 6)	
Graph showing tracing of Case A. Ileostomy (Fig. 7)	377
Case C. Scrotal hernia (Fig. 8)	379
Graph showing tracing of Dog 1A8, weight 18 kgm. (Fig. 9)	
Graph showing tracing of rabbit fundus (Fig. 1)	391
Graph showing tracing of a preparation of the trigone from the bladder of	
rabbit (Fig. 2)	391

ILLUSTRATIONS

Graph showing tracing of bladder of rabbit (Fig. 3)	
Plexus-free preparations of small intestine (Fig. 1)	
Plexus-free preparations of small intestine (Figs. 2 and 3)	400 401
Plexus-free preparations of small intestine (Figs. 5 and 6)	
Plexus-free preparations of small intestine (Figs. 7 and 8)	
Diagram comparing the average body weight of 8 control rats with that of	700
8 rats under separate and combined thyroid and morphine administra-	
tion (Fig. 1).	425
Diagram comparing the average temperature of 8 control rats, with that of	
8 rats under separate and combined thyroid and morphine administra-	
tion (Fig. 2)	426
Tracing to show action of sparteine, 1:5,000, on isolated rabbit intestine;	
prolonged period of increased tone and amplitude followed by final	
depression (Fig. 1)	437
Tracing to show action of oxysparteine, 1:1000, on isolated rabbit intes-	
\-	439
Tracing to show action of sparteine, 1:10,000, on isolated toad heart; im-	
mediate diminution of amplitude and of rate, the amplitude recovers	
gradually to nearly the original size but the slowing is progressive	441
(Fig. 3)	
Tracing from isolated toad heart (Fig. 5)	
Tracing to show action of oxysparteine, 1:20,000 on isolated toad heart; im-	770
mediate reduction of amplitude and soon cessation of beat (Fig. 6)	443
Effect of periplocin on blood-pressure and heart-rate. (Rabbit under ure-	110
thane) (Fig. 1)	455
	456
Effect of periplocin on isolated perfused frog heart (Tracing 2)	458
Effect of periplocin on isolated rabbit heart (Tracing 3)	459
Isolated jejunum of cat suspended in oxygenated Ringer-Locke solution	
(Tracing 4)	
Significance of cystine and glutathione (Chart 1)	47 8