Downloaded from jpet.aspetjournals.org at ASPET Journals on April 20, 2024

## CONTENTS

## Number 1, November, 1926

I. On the Influence of Asphyxia upon the Action of Convulsant Dyes and upon Their Entrance Into the Substance of the Central Nervous System. II. By Hans C. Syz.  II. Effects of Ephedrine on the Perfused Frog Heart and Blood Vessels. By O. W. Barlow and Torald Sollmann.  III. Studies on Strychnin. By Grace Newman.  IV. The Action of Bismuth on the Circulatory System. By George A. Masson.  V. One of the Factors Governing the Relaxation of Non-striated Muscle (Intestine) by Commercial Pituitary Extracts. By Charles M. Gruber VI. The Influence of Hydrazine and Its Derivatives on Metabolism. I. The Effect of Substitution in the Hydrazine Molecule upon the Hypoglycemic Action of Hydrazine. By Seiichi Izume and Howard B. Lewis.  VII. A Note on the Lack of Anaphylactoid Changes in the Lungs of Guinea Pigs After the Intravenous Administration of Mercurochrome-220 Soluble. By J. E. Sanner and J. H. Hill.	1 21 31 39 73 87
Number 2, December, 1926	
VIII. Studies in Chemotherapy. By Myer Coplans and Arthur G. Green. IX. The Toxicity of Bismuth. By George A. Masson	
ments of Intestine, Uterus and Ureters. By Charles M. Gruber XI. The Actions of Tyramine on the Circulation and Smooth Muscle. By M.	149
L. TainterXII. The Choleretic Action of Tolysin (Ethyl Ester of Para Methyl Phenylcinchonic Acid) in Cholecystography. A Preliminary Report. By R. G.	
Spurling and E. E. Hartman	185
Number 3, January, 1927	
XIII. The Antagonism of the Pressor Action of Tyramine by Cocaine. By M. L. Tainter and D. K. Chang	
XIV. Ephedrine Hyperglycemia in Dogs and Rabbits. By J. Allen Wilson : XV. Studies on the Cardiodynamic Actions of Drugs. I. The Application of Optical Methods of Pressure Registration in the Study of Cardiac Stimu-	209
lants and Depressants. By Carl J. WiggersXVI. Studies on the Cardiodynamic Actions of Drugs. II. The Mechanism	217
of Cardiac Stimulation by Epinephrin. By Carl J. Wiggers	233

CVII. Studies on the Cardiodynamic Actions of Drugs. III. The Mechanism of Cardiac Stimulation by Digitalis and g-Strophanthin. By Carl J. Wiggers and Barbara Stimson	271
Number 4, February, 1927	
CX. The Pharmacology of Ceanothus Americanus. I. Preliminary Studies: Hemodynamics and the Effects on Coagulation. By James T. Groot.  CXI. On the Standardisation of Digitalis by the Cat Unit Method. By A. McFarlane and G. A. Masson  CXII. Ethylene Glycol—a Pharmacological Study. By Irvine H. Page  CXIII. Some Experimental Observations on the Cardiac Effects of Acetanilid, Caffeine and its Citrate. By George B. Roth  CXIV. II. Studies on Quinin. By Soma Weiss and Robert A. Hatcher  CXV. III. Studies on Quinidin. By Soma Weiss and Robert A. Hatcher  CXVI. IV. Studies on Quinin. By Robert A. Hatcher and Harry Gold	293 313 321 327 335
Number 5, March, 1927	
CXVII. The Effect of Atropine on the Gastro-Intestinal Hemorrhage Produced by Witte's Peptone. By Erwin G. Gross	361 389
Number 6, April, 1927	
XXXI. The Toxic Action of Acetaldehyde on the Organs of Vertebrata. By J. V. Supniewski	429
with Hydrochloric Acid and Pepsin. By Herbert V. Dobson  KXXIV. A Quantitative Comparison and Toxicological Study of Ephedrine	
and Epinephrin. By J. Ernest Nadler	489

## ILLUSTRATIONS

ILLUSTRATIONS	
Chart representing for each concentration studied the cardiac changes of amplitude in millimeters, rate changes in beats per minute and output in per cent per minute change as well as changes in percentile rate flow	
through the frog vessels (Fig. 1)	23
Diagram representing the typical cardiac response to ephedrine 1:10 <sup>3</sup> when perfused into the vena cava (Fig. 2a)	25
ephedrine to a normal perfusate (Fig. 2b)	25
centration of ephedrine (Fig. 3a)	26
showing the response of the heart in figure 3a to the same concentration of ephedrine during electrical stimulation of the accelerator nerve	
(Fig. 3b)	<b>26</b>
Auricle (upper) and ventricle (lower) from a cat in an advanced stage of bismuth poisoning (Fig. 1)	44
Tracing from a cat's heart after 3.56 mgm. bismuth metal per kilogram intravenously (Fig. 2)	45
Myocardiographs from the auricle (upper) and ventricle (lower), of a dog	30
after 3.1 mgm. bismuth metal per kilogram intravenously in two doses (Fig. 3)	46
Apparatus for detecting the excitability of the heart muscle after small doses	10
of bismuth (Fig. 4)	50
after small doses of bismuth (Fig. 5)	51
Electrocardiograms (lead 2) from a dog before and at one minute's intervals after 1.08 mgm. bismuth metal per kilogram (Fig. 6)	<b>54</b>
Tracings from auricle (upper), ventricle (lower) and blood pressure from a cat 30 seconds after 3.2 mgm. bismuth metal have been injected quickly	
into a vein (Fig. 7)	59
Relaxation of non-striated muscle (Figs. 1 and 2)	77
— of non-striated muscle (Fig. 3)	78
— of non-striated muscle (Fig. 4)	79
— of non-striated muscle (Fig. 5)	80
— of non-striated muscle (Fig. 6)	81
— of non-striated muscle (Fig. 7)	82
Pig 1. Ether anesthesia only; lungs normal (Fig. 1)	97
15. Acute anaphylactic shock (Fig. 2)	97
5. Five milligrams per kilogram mercurochrome intravenously; lungs	
normal (Fig. 3)	98
Blood pressure tracing from the carotid artery of a cat (Fig. 1)	142
Excised longitudinal segments of rabbit's intestine bathed in oxygenated,	
Tyrode's solution pH 7.6 at a temperature of 37.5°C. (Fig. 1)	151

Segments taken from another animal (Fig. 2)	152
Excised cat uterus, bathed in warm 37.5°C. oxygenated Locke's solution pH 7.6 to 7.8 (Fig. 3)	154
— segments of dogs uterus bathed in warm 37.5°C, oxygenated Locke's	101
solution pH 7.4 modified by the addition of autogenous defibrinated	
blood (Fig. 4)	155
longitudinal strips of pigs' ureters, bathed in warm 37.5°C. Locke's Solu-	
tion pH 7.6 to 7.8 (Fig. 5)	156
—— longitudinal segments of pigs' ureters bathed in warm 37.5°C. Locke's	
solution pH 7.6 to 7.8 (Fig. 6)	157
Same as figure 6, except that the segments were taken from another animal	
(Fig. 7)	158
Excised longitudinal segments of a cat's uterus bathed in oxygenated Ringer's	
solution pH 7.8 to 8.0 at a temperature of 38.5°C. (Fig. 8)	160
Failure of nicotine tartrate to prevent the pressor action of tyramine. Cat	1.07
3.5 kgm. (Fig. 1)	107
sympathetic vasoconstrictions by ergot. Cat 2.1 kgm. (Fig. 2)	160
Pressor action of tyramine produced during paralysis of the sympathetic	105
vasoconstrictors by ergotoxine. Dog 6.3 kgm. (Fig. 3)	160
Stimulation of excised rabbit duodenum (longitudinal strip) by tyramine	100
(Fig. 4)	175
— of excised rabbit colon by tyramine during paralysis of parasympathetic	
endings (Fig. 5)	176
of excised, ergotoxinized, pregnant, rabbit uterus by tyramine (during	
paralysis of sympathetic augmentors) (Fig. 6)	
Two-hour film of tolysin cholecystogram (Fig. 1)	
Six-hour film from same patient (Fig. 2)	
Two-hour film of tolysin cholecystogram (Fig. 3)	
Six-hour film from same patient (Fig. 4)	190
Pressor action of tyramine abolished, and of epinephrine increased, by co-	104
caine in a rabbit (2.18 kgm.) (Fig. 1)	194
in a cat (2.65 kgm.) (Fig. 2)	104
Antagonistic action of cocaine on the changes in blood pressure, heart and	134
peripheral organ volume caused by tyramine contrasted with its sen-	
sitizing action on the effects of epinephrine in a cat (1.9 kgm) (Fig. 3).	199
- action of cocaine on the changes in blood pressure, heart and peripheral	
organ volume caused by tyramine contrasted with its sensitizing action	
on the effects of epinephrine in a dog (6.3 kgm.) (Fig. 4)	199
Ephedrine hyperglycemia in dogs and rabbits (Fig. 1)	
— hyperglycemia in dogs and rabbits (Fig. 2)	212
Four transcribed left intraventricular pressure curves showing the effects of a	
slight aortic compression (Fig. 1)	220
Three transcribed curves of left intraventricular pressure to illustrate the	
influence of changes in venous pressure, relaxation gradient and duration	001
of systole on the duration of the inflow phase (Fig. 2)	221

## ILLUSTRATIONS

Simultaneous aortic (upper) and left intraventricular pressure curves showing	
three successive beats of different vigor (Fig. 3)	225
Cardiodynamic actions of drugs (Fig. 4)	227
actions of drugs (Fig. 5)	227
Three segments of records from experiment C361, showing the influence of	
epinephrin on the pressure curves from the aorta (upper) and the left	
ventricle (middle), together with electrogram from the apex of the left	
ventricle (Fig. 1)	235
Segments of optically recorded records from experiment C373, separating the	
primary and secondary effects of epinephrin on the heart (Fig. 2)	236
Two series of superimposed tracings showing effect of epinephrin on right	
(lower), and left (upper) intraventricular pressures (Fig. 3)	238
Two records of right (lower) and left (upper) intraventricular pressures from	
experiment C348, showing, especially, opposite effects on initial tension	
when aortic pressure is allowed to rise (Fig. 4)	241
Diagram of heart upon which are located points of lead in these experiments	
(Fig. 5)	245
Five segments of records showing volume curves of ventricles (V) and mean	
blood pressure (B. P.) (Fig. 1)	253
Four segments of records of aortic pressure (upper curve) and left ventricular	
pressure (middle) from experiment 356 (Fig. 2)	257
— superimposed and transcribed curves of left ventricular pressure from	
experiment 358 (Fig. 3)	262
— superimposed and transcribed curves of left ventricular pressure from	
experiment 373, showing relations of relaxation gradient to initial pres-	
sure and effect of controlling arterial resistance (Fig. 4)	263
Pharmacology of ceanothus americanus (Tracing 1)	
— of ceanothus americanus (Tracing 2)	
— of ceanothus americanus (Tracing 3)	
— of ceanothus americanus (Tracing 4)	278
— of ceanothus americanus (Tracing 5)	278
— of ceanothus americanus (Tracing 6)	
Graph representing the group frequency of the results in table 2 (Fig. 1)	
— representing the group frequencies of the experiments in tables 2 and 4	
(Fig. 3)	305
Ethylene glycol (Fig. 1)	
— glycol (Fig. 2)	
—— glycol (Fig. 3)	
Effect of 0.05 per cent caffeine in buffered Locke-Ringer's solution on the per-	
fused frog's heart (Fig. 1)	323
— of 0.05 per cent caffeine citrate in buffered Locke-Ringer's solution on	
perfused frog's heart (Fig. 2)	323
— of 0.05 per cent sodium citrate in buffered Locke-Ringer's solution on	
	324
Curves of light absorption of two screens used to induce violet fatigue	
(Fig. 1)	372

Curve of light absorption of colored substance excreted after santonin	
Curves of light absorption of urines secreted during the first six hours (con-	390
secutively numbered) and (broken line) of urine secreted between twenty- third and twenty-fourth hours after the administration of 0.5 gram san-	
tonin (Fig. 2)	391
Change in light absorption curve of mixture of equal parts of alcoholic potash and 2 per cent santonin in alcohol (Fig. 3)	393
Curves of reaction velocity taken at λ 520μμ of equal volumes of alcoholic	บขบ
potash and (A) 1 per cent santonin in alcohol; (B) 2 per cent santonin in alcohol (Fig. 4)	394
alcohol (Fig. 4)	394
ing the seventh hour after the administration of 0.5 gram santonin	
(Fig. 5)	395
sium santoninate; (B) 0.5 gram powdered santonin; (C) 0.2 gram	
powdered santonin; (D) 0.2 gram crystalline santonin; given by the	000
mouth (Fig. 6)	396
after the administration of 1.0 gram crystalline santonin by the mouth	
(Fig. 7)	398
0.1 gram powdered santonin given by the mouth (Fig. 8)	399
Respiration curve blood pressure curve, and urine flow of a dog injected sub- cutaneously with acetaldehyde (Fig. 1)	431
Contraction curve of an isolated guinea pig's rectum in a solution of acetal-	491
	<b>433</b>
Heart beat of an isolated frog's heart perfused with acetaldehyde solution (Fig. 3)	434
Tetanus curves of a gastrocnemius of the frog in 0.8 per cent NaCl and acetal-	
dehyde solution (Fig. 4)	435 443
Cat, anesthetized with urethane. Record of cardiometer and blood pres-	
(B·-/·································	444 445
	445 445
Relation between acidity, dilution and peptic activity for gastron (Fig. 1).	471
	473
~- bob~ /9. =/	