#### CONTENTS

#### NUMBER 1, MAY, 1930

I. The Action of the "Parotid" Gland Secretion of Bufo regularis. By David	-
Epstein and J. W. C. Gunn.	1
II. Studies on the Colloid Chemistry of Antisepsis and Chemotherapy. III. The Ultramicroscopic Examination of Neoarsphenamine and of Certain Antiseptics, and Their Effects upon Protein Solutions. By Arthur D. Hirschfelder and Harold N. Wright	13
III. Studies on the Colloid Chemistry of Antisepsis and Chemotherapy. IV. The Duplication in Vitro of the "Interference Phenomenon" in Combi- nation Chemotherapy. By Harold N. Wright and Arthur D. Hirsch- felder.	39
<ul> <li>IV. On the Pharmacological Action of Various Aromatic-Aliphatic Lactones.</li> <li>II. Studies on Chemical Constitution and Pharmacological Action. By</li> <li>W. F. von Oettingen.</li> </ul>	59
V. Pharmacological Studies of Twenty-Three Isomeric Octyl Alcohols. By David I. Macht and Harriet P. Leach	71
VI. Experimental Studies on Heart Tonics. II. The Application of Bio- metric Methods to Digitalis Standardization. By William Nyiri and Louis DuBois	99
VII. Experimental Studies on Heart Tonics. III. The Relationships of Calcium Ions, Hydrogen Ions and Digitalis. By William Nyiri and Louis DuBois	
NUMBER 2, JUNE, 1930	
VIII. Some Ob (vations on the Suitability of Amytal as an Anesthetic for Laborate) inimals. By R. C. Garry	120
IX. Continu X. An Appa	137
	141
XI. The innuence of Drugs on the Transmission of Arsenic into the Aqueous Humor. By A. C. Krause, A. M. Yudkin, M. A. Stevens, W. W. Bunnell	
	153
XII. A Comparison of the Premedication Values of Several Barbituric Acid	

- XII. A Comparison of the Premedication Values of Several Barbituric Acid Derivatives in Relation to Nitrous Oxide Anesthesia. By M. S. Stormont, I. Lampe and O. W. Barlow.
  165
  XIII. Studies on the Duration of Action of Drugs. I. Analgesics and

iii

# CONTENTS

XVI. The Absorption of Drugs from the Right Ventricular Cavity. By Carl	
J. Wiggers	209
XVII. The Estimation of Digitalis by Pigeon-Emesis and Other Methods.	
By J. H. Burn	221
XVIII. Detoxification of Nicotine by Ultraviolet Rays. By A. J. Pacini and	
Hugh McGuigan	241

## NUMBER 3, JULY, 1930

<ul> <li>XIX. The Scientific Proceedings of the American Society for Pharmacology and Experimental Therapeutics. Twenty-First Annual Meeting, Held at Chicago, Ill., March 26, 27, 28, 29, 1930.</li> <li>XX. II. Observations on the Effect of Pitressin on Blood Pressure, Pulse Rate and Respiration in Dogs. By Charles M. Gruber and William B. Kountz</li> </ul>	
XXI. Ephedrine on the Ureter. By George B. Roth	
XXII. The Action of Alcohol, Caffeine, and Tobacco, on the Cardiac Output	301
, , , ,	010
(and Its Related Functions) of Normal Man. By Arthur Grollman	
XXIII. The Testing of Local Anesthetics. By T. H. Rider	329
XXIV. Pharmacological Note on Carbo Ligni and Carbo Animalis. By	
David I. Macht.	343
XXV. Biological and Chemical Studies of the Relationship between Arsenic	
and Crystalline Glutathione. By Sanford M. Rosenthal and Carl	
• •	947
Voegtlin.	341
XXVI. Studies on Crystalline Insulin. XI. Does Insulin Cause an Initial	
Hyperglycemia? By E. M. K. Geiling and A. M. De Lawder	369
NUMBER 4, AUGUST, 1930	
XXVII. The Action of an Alkaloidal Product from the Leaf of Solanum	

XXVII. The Action of an Alkaloidal Product from the Leaf of Solanum	
Pseudocapsicum, L. By J. M. Watt, H. L. Heimann and E. Meltzer	387
XXVIII. The Prevention of Acute Intoxication from Local Anesthetics. By	
P. K. Knoefel, R. P. Herwick and A. S. Loevenhart.	397
XXIX. Chemotherapy of Quinoline Compounds. Part I. A Preliminary	
Report on the Action of Certain Quinoline Compounds on Paramoecia.	
By Upendranath Brahmachari, Tarapada Bhattacharyya, Radhakrishna	
Banerjea and Bibhuti Bhusan Maity.	413
XXX. Idiosyncrasy to Quinine, Cinchonidine and Ethylhydrocupreine and	
Other Levorotatory Alkaloids of the Cinchona Series: Further Chemical	
Delimitation of the Idiosyncrasy; Alteration in Sensitiveness. By W.	
T. Dawson and Francis A. Garbade	417
XXXI. Studies in Cancer Chemotherapy. IX. The Reaction of the Blood	
in Cancer. By Fritz Bischoff, M. Louisa Long and Elsie Hill	425
XXXII. Some Observations on the Effect of Pitressin upon the Cardiovascu-	
lar System. By Charles M. Gruber and William B. Kountz	435
XXXIII. IV. The Effect of Epinephrine upon the Rate of Contraction and	
upon the Conduction Time of Peristalsis and Antiperistalsis in Excised	
Ureters. By Charles M. Gruber	<b>44</b> 9
XXXIV. Local Anesthetics Derived from Dialkylamino Propandiols. I.	
Phenylurethans. By T. H. Rider	457
XXXV. A Study of the Innervation of the Pylorus of the Terrapin. By	
Nolan L. Kaltreider	469

•

# iv

.

Perfusion of the isolated heart of the frog with a 1:8000 solution of Transvaal toad secretion (Fig. 1)	3
The effect of an intravenous injection of toad secretion (3 mgm. per kilogram) in the cat (Fig. 2)	5
	7
	8
Drawing showing the appearance of a cholesterol suspension and of egg albu- min, viewed with the ultramicroscope (Fig. 1)	17
Aggregation of egg albumin by antiseptics, viewed with the ultramicroscope	20
Effect of changes in H-ion concentration upon the flocculation of neoarsphe-	
	27
Neoarsphenamine and its reaction with egg albumin (Fig. 4)	28
Effects of neoarsphenamine and mercurochrome on rabbit's blood plasma	
(Fig. 5)	31
Studies on the colloid chemistry of antisepsis and chemotherapy (Fig. 1)	47
on the colloid chemistry of antisepsis and chemotherapy (Fig. 2)	<b>4</b> 8
on the colloid chemistry of antisepsis and chemotherapy (Fig. 3)	50
on the colloid chemistry of antisepsis and chemotherapy (Fig. 4)	52
Ichthyometer (Fig. 1)	78
Ichthyogram (Fig. 2)	79
Helicogram. Effect of octyl alcohol XXI (Fig. 3)	83
Cat, 2.6 kgm. Ether anesthesia (Fig. 4)	86
Vas deferens of rat. Effect of octyl alcohol I (Fig. 5)	88
	89
	116
Toxic effect of 0.002 per cent CaCl <sub>2</sub> , subsequent to administration of 0.002 mg.	117
Hastening and reinforcing effect of toxic digitan action by subsequent admin-	
istration of calcium (Fig. 3) 1	118
Almost immediate toxic effect of 0.08 cc. Digitan intravenously subsequent to	
0.2 per cent CaCl <sub>2</sub> in 0.65 per cent NaCl solution intravenously (Fig. 4) 1	110
Typical digitalis poisoning in slightly acid menstruum (Fig. 5)	
Male cat (Fig. 1)	
Perfusion of frog's heart with Greene's frog-heart cannula (Fig. 3)	
Female cat (Fig. 4)	134
B, capillary tube enclosed in C, straw to which is attached F, the pen (Fig. 1).	
A, reservoir with, leading from it, B, capillary tube, enclosed in C, the straw,	.00
which is nipped in $D$ , a small half-cylinder of metal, chilled at $E$ , the	
fulcrum, to fit the bearing-cones (Fig. 2)	130
Turorum, to no me bearing-conce (Fig. 2)	100

v

A, the float proper, held central by $B$ , the manometer tube cover, $C$ , the ink reservoir with $D$ , the capillary leading from it to $E$ , the pen held in $F$ ,	
celluloid (Fig. 3)	
Armlet for use with the Thompson sphygmanograph (Fig. 1)	
Wristlet for use with the Thompson sphygmanograph (Fig. 2) Photograph of the manometer and portable case (Fig. 3)	
Typical human blood pressure tracing recorded by the sphygmanograph	140
(Fig. 4)	146
Rise of blood pressure in human subject resulting from subcutaneous injection	
(five minutes previous to beginning of record of ephedrine hydrochloride).	1 477
Time intervals of five minutes (Fig. 5) Fall of blood pressure in human subject during inhalation of amyl nitrite	14/
	147
(Fig. 6) A comparison of the premedication values of several barbituric acid deriva-	14/
tives in relation to nitrous oxide anesthesia (Fig. 1)	100
comparison of the premedication values of several barbituric acid deriv-	100
atives in relation to nitrous oxide anesthesia (Fig. 2)	167
Persistence of mydriasis after graded intravenous injections of epinephrine	107
in three cats. Large doses (0.01 to 0.5 mgm.) (Fig. 1)	190
	109
cats. Small doses (0.00006 to 0.0125 mgm.) (Fig. 2)	100
	150
left common carotid artery (Fig. 3)	101
Comparison of the difference in the duration of mydriasis in the right pupil	191
after a given dose of epinephrine injected into the femoral vein $(A)$ and	
into the left common carotid artery (B). Same cat as in figure 3 (Fig. 4).	192
Persistence of mydriasis after graded intraocular injections of epinephrine	102
(Fig. 5)	193
of mydriasis after graded intravenous injections of atropine (Fig. 6)	196
of mydriasis after graded intraocular injections of atropine (Fig. 7)	197
Diagram showing arrangement of perfusion apparatus. Relative size of	
heart and manometer not drawn to scale (Fig. 1)	211
Optical pressure curves from left ventricle showing depressing effect of KCl	
injection limited to right ventricle. $B$ , 15 seconds; $C$ , 30 seconds; $D$ , 45	
seconds and $E$ , 2 minutes after injection. Time $\frac{1}{2}$ second (Fig. 2)	213
Curves same as figure 2, showing stimulating effects of CaCl <sub>2</sub> . B, 1 minute;	
C, 1.5 minutes; $D$ , 4.5 minutes after injection (Fig. 3)	213
same as figure 2, showing stimulating effects of 0.5 cc. of a 1:10,000	
epinephrin solution. $B$ , 15 seconds; $C$ , 30 seconds; $D$ , 1 minute; and $E$ ,	
2.5 minutes after injection (Fig. 4)	213
same as figure 2, showing stimulating and toxic effects of strophanthin	
(1 mgm.). B, 1 minute; C, 3 minutes; D, 4 minutes; E, 9.5 minutes; F,	
12 minutes; G, 26.5 minutes; H, 27 minutes; I, 27.5 minutes after adminis-	
tration (Fig. 5)	213
Optical pressure curves from left ventricle illustrating depressant effects of	
two doses of 1:200 chloroform. A and D, controls (Fig. 6)	216
Curves same as in figure 6, showing depressant effects of 3 cc., 1:200 chloral	010
hydrate solution. A, control; D, E and F, recovery (Fig. 7)	216

Curves same as in figure 6, showing depressant effects of 2 cc., 1:10,000 ephedrine. A, control (Fig. 8)	216
same as in figure 6, showing depressant effect of 2 cc., 1 per cent quinidine	
sulphate solution (Fig. 9)	
	228
estimation of digitalis by pigeon-emesis and other methods (Fig. 2)	228
estimation of digitalis by pigeon-emesis and other methods (Fig. 3)	
estimation of digitalis by pigeon-emesis and other methods (Fig. 4)	231
Graph illustrating the normal variations in blood pressure in unanesthetized	
dogs (Fig. 1)	280
Curve plotted from the data obtained from a 9-kgm. dog under chloretone	
anesthesia (Fig. 2)	282
obtained from a 6-kgm. dog. Local anesthesia. Upper record is that of	
the respiration with, tambour and pneumograph, the middle record the	
blood pressure with membrane manometer and the lower record the time	
interval in fifteen seconds (Fig. 3)	283
plotted from the data obtained from a non-anesthetized 6.5-kgm. dog	
(Fig. 4)	285
9.5-kgm. dog. Local anesthesia. Top record the blood pressure with mem-	
brane manometer, and bottom record the time interval in fifteen seconds	
(Fig. 5)	286
Curve plotted from the data obtained from an experiment upon an unanesthe-	
tized 9.5-kgm. dog (Fig. 6)	288
6-kgm. dog. Local anesthesia, both vagi cut four hours earlier. Top record	
the blood pressure with membrane manometer, bottom record the time	
interval in fifteen seconds (Fig. 7)	289
Curve plotted from the data obtained from an experiment upon an 8-kgm.	
dog. Both vagi cut two hours previously. Local anesthesia (Fig. 8)	290
14-kgm. dog under ether anesthesia. Both adrenal glands excised. Top	
curve that of the blood pressure with a membrane manometer, bottom	
curve the time interval in fifteen seconds (Fig. 9)	292
12-kgm. dog. Local anesthesia. Both carotid arteries had been tied off	
under ether anesthesia three hours before this record was made. Top	
curve the blood pressure with membrane manometer and bottom record	004
the time in fifteen-second intervals (Fig. 10)	294
Effect of 10 mgm. of ephedrine sulphate (Lilly & Co.) in 100 cc. of alkaline	
Locke-Ringer on the excised dogs' ureter (Fig. 1)	305
of 20 mgm. of ephedrine hydrochloride (Read) on the excised ureter of	007
dog (Fig. 2).	305
of 80 additional milligrams of ephedrine hydrochloride (Read) on the	200
same segment of dog's ureter which was used to obtain figure 2 (Fig. 3)	306
of 20 mgm. of ephedrine hydrochloride (Read) in 100 cc. of alkaline	907
Locke-Ringer, on a quiescent segment of dogs' ureter (Fig. 4)	307
of 0.1 mgm. of epinephrine (Parke Davis & Co.) on the same segment	
which was used for obtaining figure 4, the Locke-Ringer having been changed seven minutes previously (Fig. 5)	200
	909
	308
pigs' ureter (Fig. 6)	909

•

Prevention of the local inflammatory reaction from N/100 arsenoxide when	
N/10 glutathione is simultaneously injected (right ear) (Fig. 1)	356
Biological and chemical studies of the relationship between arsenic and	
crystalline glutathione (Chart 1)	360
and chemical studies of the relationship between arsenic and crystalline	
glutathione (Chart 2)	360
and chemical studies of the relationship between arsenic and crystalline	
glutathione (Chart 3)	361
glutathione (Chart 4)	361
Blood sugar effects of insulin preparations at various stages of purification	
(Fig. 1).	375
	379
Cats. Electrocardiogram. Lead II. Time $\frac{1}{5}$ and $\frac{1}{25}$ second (Fig. 1)	
The action of an alkaloidal product from the leaf of Solanum pseudocapsicum,	991
	200
L. (Fig. 4)	
Cat—to show intermittent stoppage of the heart (Fig. 5)	
Record I. 8.46 kgm. dog. Amytal anesthesia (Fig. 1).	406
Some observations on the effect of pitressin upon the cardiovascular system	
(Fig. 1).	441
observations on the effect of pitressin upon the cardiovascular system	
(Fig. 2)	441
observations on the effect of pitressin upon the cardiovascular system	
(Fig. 3)	443
observations on the effect of pitressin upon the cardiovascular system	
(Fig. 4)	444
	452
Excised pig ureter kept in ice cold Locke's solution 24 hours. Bladder half of	
ureter (Fig. 2)	453
Showing the stimulating effect of pilocarpine, the inhibiting action of ephe-	
drine and epinephrin and the antagonistic action of atropine on the	
rhythmic contractions of the pylorus (Fig. 1)	473
Illustrating the effect of ephedrine on the tonus of the pylorus and lack of	
stimulation by pilocarpine, following atropine (Fig. 2)	473
Showing the characteristic rhythmic contractions with no alteration of the	
base-line, in response to pilocarpine on the pyloric sphincter (Fig. 3)	475
the typical response of the pyloric sphincter to epinephrin, when not	
previously stimulated by drugs (Fig. 4)	475
no effect of adrenalin or ephedrine on the rhythmic contractions and on	
the tonus of the pylorus (Fig. 5)	475
Illustrating the stimulating effect of epinephrin and ephedrine on the tonus	
	475
Showing the typical effect of ephedrine on quiescent, relaxed sphincters not	
previously stimulated by drugs (Fig. 7)	475
contractions and tonus of the small intestine, initiated by pilocarpine	
	475
<u> </u>	

viii

.