

**INVOLVEMENT OF Na⁺-Ca²⁺ EXCHANGER IN INTRACELLULAR Ca²⁺
INCREASE AND NEURONAL INJURY INDUCED BY POLYCHLORINATED
BIPHENYLS IN HUMAN NEUROBLASTOMA SH-SY5Y CELLS**

SIMONA MAGI, PASQUALINA CASTALDO, GIUSEPPINA CARRIERI, ANTONELLA
SCORZIELLO, GIANFRANCO DI RENZO AND SALVATORE AMOROSO

Unit of Pharmacology, Department of Neuroscience, School of Medicine, University
“Politecnica delle Marche”, Ancona, Italy (SM, PC, GC, SA); and Unit of Pharmacology,
Department of Neuroscience, School of Medicine, University of Naples “Federico II”,
Naples, Italy (AS, GDR)

Running title:

Na⁺-Ca²⁺-exchanger and PCBs Ca²⁺ increase.

Corresponding author: Prof. Salvatore Amoroso, Unit of Pharmacology, Department of Neuroscience, School of Medicine, University “Politecnica delle Marche”, Via Tronto 10/A, 60020 Torrette, Ancona, Italy

Phone +39-071-2206176. Fax +39-071-2206178.

E-mail : s.amoroso@univpm.it

Number of text pages: 25

Number of tables: 1

Number of figures: 6

Number of references: 41

Number of words in the *Abstract*: 163

Number of words in the *Introduction*: 363

Number of words in the *Discussion*: 897

Abbreviations used: A1254, Aroclor 1254; AS1, NCX1 antisense; AS3, NCX3 antisense; KB-R7943, 2-[2-[4-(4-nitrobenzyloxy) phenyl] ethyl] isothiourea methanesulphonate; [Ca²⁺]_i, intracellular calcium concentration; DMSO, dimethylsulphoxide; Fura-2AM, fura-2 acetoxymethyl ester; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; IP3, inositol 1,4,5-triphosphate; MTT, 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyltetrazolium bromide; NCX, sodium calcium exchanger; NMDA, N-methyl-D-aspartate; ODNs, oligodeoxynucleotides; PBS, phosphate buffered saline; PCBs, polychlorinated biphenyls; VSCCs, voltage-sensitive Ca²⁺ channels.

Abstract

In SH-SY5Y, a human neuroblastoma cell line, Aroclor 1254 (A1254) induced a dose-dependent (10-50 $\mu\text{g/ml}$) intracellular calcium concentration ($[\text{Ca}^{2+}]_i$) increase. Two rather specific sodium calcium exchanger (NCX) inhibitors like bepridil (10 μM) and 2-[2-[4-(4-nitrobenzyloxy) phenyl] ethyl] isothiourea methanesulphonate (KB-R7943) (10 μM), reduced A1254-induced $[\text{Ca}^{2+}]_i$ increase. 24 hours exposure to 30 $\mu\text{g/ml}$ A1254 caused a remarkable SH-SY5Y neuroblastoma cell damage. Interestingly, both bepridil and KB-R7943 counteracted A1254-induced neuronal injury. These results indicate that NCX contributes to $[\text{Ca}^{2+}]_i$ increase and neuronal injury induced by A1254. RT-PCR experiments revealed in SH-SY5Y neuroblastoma cells, expression of NCX1 and NCX3 isoforms. In order to investigate which isoform was involved in $[\text{Ca}^{2+}]_i$ increase and neuronal damage induced by A1254, we used specific antisense oligodeoxynucleotides (ODNs) to reduce NCX1 or NCX3 protein expression. The results showed that only NCX1 ODN reduced $[\text{Ca}^{2+}]_i$ increase and neuronal injury induced by A1254. In conclusion, these results indicate that, NCX1 may participate to $[\text{Ca}^{2+}]_i$ increase and neurotoxicity evoked by A1254 in SH-SY5Y neuroblastoma cells.

Introduction

Polychlorinated biphenyls (PCBs) are a class of persistent pollutants present in the environment, and there is increasing evidence that exposure to PCBs can cause neurotoxicity (Hwang et al., 2001; Mariussen et al., 2002; Howard et al., 2003; Sanchez-Alonso et al., 2003, 2004; Kang et al., 2004; Lee and Opanashuk, 2004). A number of potential mechanisms have been proposed to explain PCBs neurotoxicity i.e. alterations in neurotransmitter levels (Bemis and Seegal, 1999), perturbation in intracellular second messenger systems (Kodavanti and Tilson, 1997; Yang and Kodavanti, 2001) and elevation of intracellular Ca^{2+} concentrations ($[\text{Ca}^{2+}]_i$) (Sharma et al., 2000). This last effect has been attributed to a Ca^{2+} influx (Mundy et al., 1999; Inglefield and Shafer, 2000), or to the inhibition of Ca^{2+} -ATPase activity at the synaptic level and Ca^{2+} sequestration by mitochondria and microsomes (Kodavanti et al., 1993). In addition it has been reported that PCBs can also perturb Ca^{2+} homeostasis by acting on ryanodine (Wong and Pessah, 1996; Mundy et al., 1999) and inositol 1,4,5-triphosphate (IP3) receptor-sensitive Ca^{2+} release (Inglefield et al., 2001).

The Na^+ - Ca^{2+} exchanger (NCX) is a plasmamembrane exchanger mainly involved in the maintenance of cytosolic Ca^{2+} homeostasis (Sanchez-Armass and Blaustein, 1987). This antiporter couples the uphill extrusion of Ca^{2+} to the entrance of Na^+ into the cell (forward mode) and down its electrochemical gradient. However, this mechanism can also operate as a Na^+ efflux- Ca^{2+} influx pathway, depending not only on membrane potential (Snelling and Nicholls, 1985), but also on the intracellular concentrations of Na^+ and Ca^{2+} (Amoroso et al., 1997, 2000; Pignataro et al., 2004a, 2004b).

In light of the role played by NCX in the maintenance of $[\text{Ca}^{2+}]_i$ homeostasis, the aim of the present study was to investigate the role played by NCX in PCBs-induced $[\text{Ca}^{2+}]_i$ perturbation and neuronal injury. For this purpose SH-SY5Y cells were exposed to Aroclor 1254 (A1254), a commercial mixture of PCBs, and Fura-2AM monitored $[\text{Ca}^{2+}]_i$ and 3-[4,5-

dimethylthiazol-2-yl]-2,5 diphenyltetrazolium bromide (MTT) detected cell injury, were evaluated in the presence or in the absence of NCX inhibitors like bepridil (Amoroso et al., 2000; Pignataro et al., 2004a), KB-R7943 (Iwamoto and Shigekawa, 1998) and specific NCX antisense oligodeoxynucleotides (ODNs) (Pignataro et al., 2004b).

Materials and Methods

Cell culture

The SH-SY5Y cell line (purchased from the American Type Culture Collection) was cultured as monolayer in polystyrene dishes (100 mm diameter) and grown in RPMI 1640 medium (Gibco, Grand Island, NY USA) containing 10% heat inactivated fetal bovine serum (Gibco, Grand Island, NY USA), 1% of 200 mM L-glutamine (Gibco Grand Island, NY USA), 1% of sodium pyruvate 100 mM (Gibco Grand Island, NY USA), 100 IU/ml penicillin (Gibco, Grand Island, NY USA), and 100 µg/ml streptomycin (Gibco, Grand Island, NY USA). Cells were grown in a humidified incubator at 37°C in a 5% CO₂ atmosphere. The medium was changed every 2 days. Each experiment was performed with cells (passage 40-60) in multiple well flow dishes.

Cytosolic Ca²⁺ concentration measurements

Just before the experiment, 2x10⁶ SH-SY5Y cells were detached, then centrifuged and resuspended in 1 ml of a medium whose composition was in mM: NaCl 140, KCl 5, CaCl₂ 1, MgCl₂ 0.5, glucose 5.5, Hepes 10; pH was adjusted to 7.4 with 1 M Tris (standard buffer). The cells were then incubated with 5 µM Fura-2AM (Calbiochem, San Diego, CA, USA) for 1 hour at 30°C. After the loading period, the medium was diluted with 2 volumes of the same balanced salt solution, incubated at 37°C, and then washed twice before the experiment was performed (Amoroso et al., 2000). [Ca²⁺]_i values were measured in a 2 ml suspension of SH-SY5Y cells at 37°C in a quartz cuvette equipped with a magnetic stirrer bar. Fura-2 fluorescence was monitored in a Perkin Elmer model 55 LS spectrophotofluorimeter. The excitation wavelengths were at 340 and 380 nm (bandpass 5 nm) with emission at 509 nm (bandpass 5 nm). [Ca²⁺]_i values were determined according to the equation of Grynkiewicz et al., (1985).

Determination of cell viability evaluated as mitochondrial activity

Cell viability evaluated as mitochondrial activity was quantified by measuring dehydrogenase activity retained in the cultured cells, using the MTT (Sigma-Aldrich, Milan, Italy) assay (Mossman, 1983; Amoroso et al., 1999). The assay is based on the ability of living cells to convert dissolved MTT into insoluble formazan. Therefore, the amount of formazan produced is proportional to the number of living cells. Briefly SH-SY5Y cells (3×10^5) were incubated in 2 ml MTT solution (0.5 mg/ml) for 1 hour in a humidified 5% CO₂ incubator at 37°C. Thereafter the medium was removed and cells were washed twice with PBS. Then, 1 ml of dimethylsulphoxide (DMSO) was added to the cells to solubilize the formazan. The absorbance was read at 540 nm in a plate reader (Victor² 1420 Multilabel Counter, Perkin Elmer). Data are expressed as amount of formazan produced. In control cultures the same volume of A1254 vehicle was added. The osmotic pressure, measured by Autostat Osmometer 6030 (Arkay, KDK corporation, Japan) in media containing standard buffer or A1254 (30 µg/ml) or its vehicle, was identical (302 mOsm). When experiments with oligos were performed, A1254 was added 5 hours after the transfection.

RT-PCR analysis of mRNA expression of NCX isoforms in SH-SY5Y cells

Total RNA was extracted from SH-SY5Y cells by using Trizol reagent (Invitrogen, Milan, Italy). To avoid contamination with genomic DNA, the extracted RNA was treated with 10U/µl of RNAase-free DNAase I (Stratagene, Milan, Italy) for 1 hour at 37°C. The purity and integrity of RNA was checked by denaturing agarose gel electrophoresis. Two micrograms of total RNA were reverse-transcribed in the presence of oligo(dt), by using SuperScript III reverse transcriptase (Invitrogen, Milan, Italy), according to the manufacturer protocol. The retrotranscribed cDNAs (2 µl) were then amplified in a MJ Research PTC 2000 Peltier Thermal Cycler (CELBIO, Milan, Italy) by using the primers shown in Table 1, previously described by Quednau et al., (1997) and Papa et al., (2003). Each 50 µl reaction

containing 1.25 U of Taq DNA Polymerase (Eppendorf, Milan, Italy) and 10 pmol of each primer, was amplified (30 cycles) by the following procedure: 94°C for 1 min, 55°C for 1 min and 72°C for 1 min. The amplification products were visualized on agarose (2%) gel electrophoresis, loading approximately half (25 μ l) of each reaction volume per lane.

Oligos Transfection

One day after plating, SH-SY5Y cells were transfected with the appropriate phosphorothioate ODNs (Table 1) previously described by Pignataro et al., (2004b), by using lipofectamine 2000 according to the manufacturer protocol (Invitrogen, Milan, Italy). Briefly, oligos (5 μ M) were dissolved in lipofectamine (DNA/lipofectamine 2000 1:1), and then incubated at room temperature for 30 min before the addition to the culture medium. Control dishes received lipofectamine only. In order to evaluate cell transfection efficiency, ODNs were mixed with a plasmid encoding the enhanced green fluorescent protein as marker (Castaldo et al., 2004). Efficiency transfection was \cong 90% (data not shown).

Drugs and Chemicals

All the chemicals were of analytical grade and were purchased from Sigma (Milan, Italy). A1254 solution in isooctane (Lot Number M-841B, Analyte Lot NT01022, stock solution 1000 \pm 5 μ g/ml) was purchased from Ultra Scientific, North Kingstown, RI; bepridil was obtained from Sigma Italy, KB-R7943 was obtained from Tocris UK. Bepridil was dissolved in a mixture of acetone and water (stock solution 10 mM), whereas KB-R7943 was dissolved in a mixture of DMSO and water (stock solution 10 mM).

Statistics

Data were analyzed by one-way ANOVA followed by Dunnett's test.

Results

Effect of different concentrations of A1254 on $[Ca^{2+}]_i$ in the presence and in the absence of extracellular Ca^{2+} in SH-SY5Y cells.

When SH-SY5Y cells were exposed to different concentrations (10, 30 and 50 $\mu\text{g/ml}$) of A1254, a dose-dependent increase of $[Ca^{2+}]_i$ occurred. This increase started almost immediately and lasted for the whole considered time. The elevation of $[Ca^{2+}]_i$ induced by A1254 (30 $\mu\text{g/ml}$) was completely abolished by removal of extracellular Ca^{2+} (Fig.1).

Effect of bepridil and the isothiourea derivative KB-R7943, two rather specific inhibitors of NCX, on the $[Ca^{2+}]_i$ elevation induced by A1254.

Bepridil and the isothiourea derivative KB-R7943, added to the incubation medium at the concentration of 10 μM , known to inhibit NCX activity (Iwamoto and Shigekawa, 1998; Amoroso et al., 2000; Annunziato et al., 2004; Pignataro et al., 2004a), did not modify basal $[Ca^{2+}]_i$, whereas they counteracted $[Ca^{2+}]_i$ increase induced by A1254 (30 $\mu\text{g/ml}$) (Fig.2).

Effect of bepridil and KB-R7943 on A1254-induced neuronal injury.

When SH-SY5Y cells were exposed to A1254 (30 $\mu\text{g/ml}$) for 24 hours, a significant reduction of cell survival, evaluated as mitochondrial activity by means of MTT method, occurred. Bepridil and KB-R7943, at the concentration (10 μM) which had reduced $[Ca^{2+}]_i$, rises triggered by A1254, were able to prevent neuronal injury evoked by PCB mixture (Fig. 3).

RT-PCR analysis of SH-SY5Y mRNA expression of NCX1, NCX2 and NCX3.

Regarding the NCX family, three dominant genes coding for three different NCX1 (Nicoll et al., 1990), NCX2 (Li et al., 1994) and NCX3 (Nicoll et al., 1996) proteins have been identified in several mammal tissues, including central nervous system. In order to characterize the pattern of the expression of the three NCX isoforms in SH-SY5Y cells, RT-PCR analysis of NCX1, NCX2 and NCX3 mRNA expression was performed. As shown in

Fig. 4, only mRNA expression of NCX1 and NCX3 was detected in SH-SY5Y cell line, (Fig.4).

RT-PCR analysis of NCX1 and NCX3 mRNA expression in SH-SY5Y transfected with the respective sense and antisense ODNs.

As shown in Fig 5, the transfection with NCX1 and NCX3 antisenses caused a marked reduction ($\cong 80\%$) of NCX1 and NCX3 mRNA expression, whereas the transfection with the NCX1 and NCX3 senses failed to change mRNA levels of the two isoforms.

Effect of NCX1 and NCX3 antisense ODNs on $[Ca^{2+}]_i$ increase and neuronal injury elicited by A1254.

In SH-SY5Y cells transfected with NCX1 or NCX3 antisense ODNs the basal $[Ca^{2+}]_i$ were similar to those of untransfected cells. When the effect of NCX1 and NCX3 antisense ODNs on $[Ca^{2+}]_i$ increase evoked by A1254 was considered, a reduction ($\cong 60\%$) of $[Ca^{2+}]_i$ elevation induced by the PCB mixture occurred in cells transfected with NCX1 antisense ODN, whereas in cells transfected with NCX3 antisense ODN, A1254 caused the same increase of $[Ca^{2+}]_i$ as that detected in untransfected cells (Fig. 6A). Interestingly, in cells transfected with NCX1 ODN antisense, inhibition of $[Ca^{2+}]_i$ increase evoked by A1254 was coupled to a marked reduction of neuronal injury (Fig. 6B), while in NCX3 antisense ODN transfected SH-SY5Y cells, the extent of neuronal injury induced by A1254 was not modified exactly as $[Ca^{2+}]_i$. NCX1 and NCX3 sense ODNs did not modify both $[Ca^{2+}]_i$ increase and neuronal injury induced by A1254 (data not shown).

Discussion

The results of present study showed that in SH-SY5Y neuroblastoma cells the PCB mixture A1254 induces a dose-dependent increase of $[Ca^{2+}]_i$ that is dependent upon extracellular Ca^{2+} ions. In fact the removal of extracellular Ca^{2+} completely abolished A1254-induced $[Ca^{2+}]_i$ increase. These results are in agreement with previous studies performed in cultured cerebellar granule cells by Mundy et al., (1999). Regarding the mechanism by which A1254 induced an influx of Ca^{2+} , it has been suggested that extracellular Ca^{2+} entry occurs via L-type voltage sensitive calcium channels (VSCCs), since A1254-evoked intracellular Ca^{2+} oscillations was abolished by addition of nifedipine (Inglefield and Shafer, 2000). We now report that NCX exchanger may be involved in Ca^{2+} entry induced by the PBC mixture. In fact both bepridil, a rather specific inhibitor of NCX (Amoroso et al., 2000; Annunziato et al., 2004; Pignataro et al., 2004a) and KB-R7943, which inhibits NCX only when it is operating in the reverse mode (Iwamoto and Shigekawa, 1998), significantly counteracted $[Ca^{2+}]_i$ increase induced by A1254. However, bepridil has been reported also to block Ca^{2+} (Kaczorowski et al., 1989) and Na^+ channels (Kaczorowski et al., 1989), and KB-R7943, besides its peculiar NCX blocking properties, also exerts an inhibitory effect on several other ionic transport mechanisms, such L-type VSCCs and receptor operated ion channels, like NMDA (Matsuda et al., 2001; Annunziato et al., 2004). Therefore the possibility exists that the inhibition produced by these drugs of $[Ca^{2+}]_i$ increase induced by A1254, might be due to the blockade of Na^+ and/or Ca^{2+} channels and/or NMDA receptor activation. On the other hand this possibility seems to be supported by the results of Inglefield and Shafer (2000) showing that A1254-induced Ca^{2+} oscillations in developing neocortical cells can be blocked by the VSCC antagonist nifedipine, by the voltage sensitive Na^+ channel antagonist, tetrodotoxin (Almers and Levinson, 1975; Weiser and Wilson, 2002) and by ionotropic glutamate receptor antagonists. In order to clarify whether NCX is effectively involved in

A1254-induced $[Ca^{2+}]_i$ increase, firstly the pattern of the expression of the three known NCX isoforms NCX1, NCX2 and NCX3 (Annunziato et al., 2004) was characterized in SH-SY5Y cells by RT-PCR analysis. Then, the effect of A1254 on $[Ca^{2+}]_i$ increase was investigated in SH-SY5Y transfected with the antisense NCX1 or NCX3 ODNs, since only these two isoforms were detected. The results of these experiments showed that, a significant reduction ($\cong 60\%$) of $[Ca^{2+}]_i$ elevation induced by the PCB mixture occurred in cells transfected with antisense NCX1 ODN. These findings demonstrate that the NCX is effectively involved in $[Ca^{2+}]_i$ elevation induced by PCB mixture, as suggested by results obtained with bepridil and KB-R7943. In addition, it is noteworthy that between the two NCX isoforms detected in SH-SY5Y cells, only NCX1 seems to be involved in this phenomenon. In fact, no inhibition of A1254-induced $[Ca^{2+}]_i$ increase was found in cells transfected with antisense NCX3 ODN. If the inhibition of NCX reduces A1254-induced Ca^{2+} influx, this implies that NCX is operating in the reverse mode (Ca^{2+} entry- Na^+ efflux pathway). This mode of operation could be explained by the fact that A1254 induces a membrane depolarization, as suggested by Inglefield and Shafer, (2000), with a consequent opening of Na^+ sensitive voltage channels. This event leads to an increase of intracellular Na^+ ion concentration, which in turn may force the NCX to work in the reverse mode (Baker and McNaughton, 1976; DiPolo, 1979; Sanchez-Armass and Blaustein, 1987; Annunziato et al., 2004). This chain of events seems to be supported by the results showing that tetrodotoxin, a well known blocker of fast activated Na^+ channels (Almers and Levinson, 1975; Weiser and Wilson, 2002) prevents Ca^{2+} oscillations evoked by A1254 in developing cortical neurons (Inglefield and Shafer, 2000). Thus, the hypothesis that, in SH-SY5Y cells, A1254 induces a membrane depolarization with a consequent intracellular increase of Na^+ ions which in turn activates NCX to operate in the reverse mode, seems reasonable.

However, the inhibition of NCX did not fully counteract $[Ca^{2+}]_i$ elevation induced by the PCB mixture. These results suggest that other mechanisms may be also involved in A1254-elicited $[Ca^{2+}]_i$ increase, like extracellular Ca^{2+} influx through L-type VSCCs (Inglefield and Shafer, 2000); inhibition of Ca^{2+} -ATPase activity and Ca^{2+} sequestration by mitochondria and microsomes (Kodavanti et al., 1993) and mobilization of intracellular Ca^{2+} stores (Inglefield et al., 2001).

Another finding of the present paper that deserves to be discussed is that the inhibition of A1254-elicited $[Ca^{2+}]_i$ increase mediated by NCX is able to protect SH-SY5Y cells from the injury induced by the exposure to the PCB mixture. In fact, both bepridil and KB-R7943, at the same concentrations that inhibited $[Ca^{2+}]_i$ increase, caused a significant reduction of cell injury elicited by A1254 and in cells transfected with antisense NCX1 ODNs, the extent of cell damage induced by the PCB mixture was significantly lower than that observed in naïve cells. On the other hand it is well known that an increase of $[Ca^{2+}]_i$ can trigger intracellular pathways leading to cell death (Kristian and Siesjo, 1998; Kang et al., 2002, 2004). It is noteworthy that cell protection from A1254 elicited injury, obtained by the inhibition of NCX, was not complete, exactly as the reduction of A1254- elicited $[Ca^{2+}]_i$ increase.

In conclusion, the results of the present paper seem to suggest that NCX may participate to $[Ca^{2+}]_i$ increase induced by A1254 exposure and that its inhibition may contribute to protect cells from the injury evoked by the PCB mixture.

Acknowledgements

The authors thank Gerardo Galeazzi, Franco Pettinari and Carlo Alfredo Violet for their invaluable technical assistance.

References

Almers W and Levinson SR (1975) Tetrodotoxin binding to normal depolarized frog muscle and the conductance of a single sodium channel. *J Physiol* **247**:483-509.

Amoroso S, De Maio M, Russo GM, Catalano A, Bassi A, Montagnani S, Di Renzo G and Annunziato L (1997) Pharmacological evidence that the activation of the Na⁺-Ca²⁺ exchanger protects C6 glioma cells during chemical hypoxia. *Br J Pharmacol* **121**:303-309.

Amoroso S, Gioielli A, Cataldi M, Di Renzo G and Annunziato L (1999) In the neuronal cell line SH-SY5Y, oxidative stress-induced free radical overproduction causes cell death without any participation of intracellular Ca²⁺ increase. *Biochim Biophys Acta* **1452**:151-160.

Amoroso S, Tortiglione A, Secondo A, Catalano A, Montagnani S, Di Renzo G and Annunziato L (2000) Sodium nitroprusside prevents chemical hypoxia-induced cell death through iron ions stimulating the activity of the Na⁺-Ca²⁺ exchanger in C6 glioma cells. *J Neurochem* **74**:1505-1513.

Annunziato L, Pignataro G and Di Renzo GF (2004) Pharmacology of brain Na⁺-Ca²⁺ exchanger: from molecular biology to therapeutic perspectives. *Pharmacol Rev* **56**:633-654.

Baker PF and McNaughton PA (1976) Kinetics and energetics of calcium efflux from intact squid giant axons. *J Physiol* **259**:103-144.

Bemis JC and Seegal RF (1999) Polychlorinated biphenyls and methylmercury act synergistically to reduce rat brain dopamine content in vitro. *Environ Health Perspect* **107**:879-885.

Castaldo P, Stefanoni P, Miceli F, Coppola G, Miraglia del Giudice E, Bellini G, Pascotto A, Trudell JR, Harrison NL, Annunziato L and Tagliatela M (2004) A novel hyperekplexia-causing mutation in the pre-transmembrane segment 1 of the human glycine receptor alpha subunit reduces membrane expression and impairs gating by agonists. *J Biol Chem* **279**:25598-25604.

DiPolo R (1979) Calcium influx in internally dialyzed squid giant axons. *J Gen Physiol* **73**:91-113.

Grynkiewicz G, Poenie M and Tsien RY (1985) A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J Biol Chem* **260**:3440-3450.

Howard AS, Fitzpatrick R, Pessah I, Kostyniak P and Lein PJ (2003) Polychlorinated biphenyls induce caspase-dependent cell death in cultured embryonic rat hippocampal but not cortical neurons via activation of the ryanodine receptor. *Toxicol Appl Pharmacol* **190**:72-86.

Hwang SG, Lee HC, Lee DW, Kim YS, Joo WH, Cho YK and Moon JY (2001) Induction of apoptotic cell death by a p53-independent pathway in neuronal SK-N-MC cells after treatment with 2,2',5,5'-tetrachlorobiphenyl. *Toxicology* **165**:179-188.

Inglefield JR and Shafer TJ (2000) Polychlorinated biphenyl-stimulation of Ca^{2+} oscillations in developing neocortical cells: a role for excitatory transmitters and L-type voltage-sensitive Ca^{2+} channels. *J Pharmacol Exp Ther* **295**:105-113.

Inglefield JR, Mundy WR and Shafer TJ (2001) Inositol 1,4,5-triphosphate receptor-sensitive Ca^{2+} release, store-operated Ca^{2+} entry, and cAMP responsive element binding protein phosphorylation in developing cortical cells following exposure to polychlorinated biphenyls. *J Pharmacol Exp Ther* **297**:762-773.

Iwamoto T and Shigekawa M (1998) Differential inhibition of $\text{Na}^{+}\text{-Ca}^{2+}$ exchanger isoforms by divalent cations and isothiourea derivative. *Am J Physiol* **275**:C423-C430.

Kaczorowski GJ, Slaughter RS, King VF and Garcia ML (1989) Inhibitors of sodium-calcium exchange: identification and development of probes of transport activity. *Biochim Biophys Acta* **988**:287-302.

Kang JH, Jeong W, Park Y, Lee SY, Chung MW, Lim HK, Park IS, Choi KH, Chung SY, Kim DS, Park CS, Hwang O, Kim J (2002) Aroclor 1254-induced cytotoxicity in catecholaminergic CATH.a cells related to the inhibition of NO production. *Toxicology* **177**:157-166.

Kang JH, Park IS, Oh WY, Lim HK, Wang SY, Lee SY, Choi KH, Kim JI, Jung SY, Suh CK and Kim DS (2004) Inhibition of aroclor 1254-induced depletion of stored calcium prevents the cell death in catecholaminergic cells. *Toxicology* **200**:93-101.

Kodavanti PR, Shin DS, Tilson HA. and Harry GJ (1993) Comparative effects of two polychlorinated biphenyl congeners on calcium homeostasis in rat cerebellar granule cells. *Toxicol Appl Pharmacol* **123**:97-106.

Kodavanti PR and Tilson HA (1997) Structure-activity relationships of potentially neurotoxic PCB congeners in the rat. *Neurotoxicology* **18**:425-441.

Kristian T and Siesjo BK (1998) Calcium in ischemic cell death. *Stroke* **29**:705-718.

Lee DW and Opanashuk LA (2004) Polychlorinated biphenyl mixture aroclor 1254-induced oxidative stress plays a role in dopaminergic cell injury. *Neurotoxicology* **25**:925-939.

Li Z, Matsuoka S, Hryshko LV, Nicoll DA, Bersohn MM, Burke EP, Lifton RP and Philipson KD (1994) Cloning of the NCX2 isoform of the plasma membrane Na⁺-Ca²⁺ exchanger. *J Biol Chem* **269**:17434-17439.

Mariussen E, Myhre O, Reistad T and Fonnum F (2002) The polychlorinated biphenyl mixture aroclor 1254 induces death of rat cerebellar granule cells: the involvement of the N-methyl-D-aspartate receptor and reactive oxygen species. *Toxicol Appl Pharmacol* **179**:137-144.

Matsuda T, Arakawa N, Takuma K, Kishida Y, Kawasaki Y, Sakaue M, Takahashi K, Takahashi T, Suzuki T, Ota T, Hamano-Takahashi A, Onishi M, Tanaka Y, Kameo K and Baba A (2001) SEA0400, a novel and selective inhibitor of the Na⁺-Ca²⁺ exchanger,

attenuates reperfusion injury in the in vitro and in vivo cerebral ischemic models. *J Pharmacol Exp Ther* **298**:249-256.

Mossman T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol. Methods* **65**:55-63.

Mundy WR, Shafer TJ, Tilson HA and Kodavanti PR (1999) Extracellular calcium is required for the polychlorinated biphenyl-induced increase of intracellular free calcium levels in cerebellar granule cell culture. *Toxicology* **136**:27-39.

Nicoll DA, Longoni S and Philipson KD (1990) Molecular cloning and functional expression of the cardiac sarcolemmal Na⁺-Ca²⁺ exchanger. *Science* **250**:562-565.

Nicoll DA, Quednau BD, Qui Z, Xia YR, Lusic AJ and Philipson KD (1996) Cloning of a third mammalian Na⁺-Ca²⁺ exchanger, NCX3. *J Biol Chem* **271**:24914-24921.

Papa M, Canitano A, Boscia F, Castaldo P, Sellitti S, Porzig H, Taglialatela M and Annunziato L (2003) Differential expression of the Na⁺-Ca²⁺ exchanger transcripts and proteins in rat brain regions. *J Comp Neurol* **461**:31-48.

Pignataro G., Tortiglione A., Scorziello A., Giaccio L., Secondo A., Severino B., Santagada V., Caliendo G., Amoroso S., Di Renzo G. and Annunziato L (2004a) Evidence for a protective role played by the Na⁺/Ca²⁺ exchanger in cerebral ischemia induced by middle cerebral artery occlusion in male rats. *Neuropharmacology* **46**, 439-448.

Pignataro G, Gala R, Cuomo O, Tortiglione A, Giaccio L, Castaldo P, Sirabella R, Matrone C, Canitano A, Amoroso S, Di Renzo G and Annunziato L (2004b) Two sodium/calcium exchanger gene products, NCX1 and NCX3, play a major role in the development of permanent focal cerebral ischemia. *Stroke* **35**:2566-2570.

Quednau BD, Nicoll DA and Philipson KD (1997) Tissue specificity and alternative splicing of the Na⁺-Ca²⁺ exchanger isoforms NCX1, NCX2, and NCX3 in rat. *Am J Physiol* **272**:C1250-C1261.

Sanchez-Alonso JA, Lopez-Aparicio P, Recio MN and Perez-Albarsanz MA (2003) Apoptosis-mediated neurotoxic potential of a planar (PCB 77) and a nonplanar (PCB 153) polychlorinated biphenyl congeners in neuronal cell cultures. *Toxicol Lett* **144**:337-349.

Sanchez-Alonso JA, Lopez-Aparicio P, Recio MN and Perez-Albarsanz MA (2004) Polychlorinated biphenyl mixtures (Aroclors) induce apoptosis via Bcl-2, Bax and caspase-3 proteins in neuronal cell cultures. *Toxicol Lett* **153**:311-326.

Sanchez-Armass S and Blaustein MP (1987) Role of sodium-calcium exchange in regulation of intracellular calcium in nerve terminals. *Am J Physiol* **252**:C595-C603.

Sharma R, Derr-Yellin EC, House DE and Kodavanti PR (2000) Age-dependent effects of Aroclor 1254R on calcium uptake by subcellular organelles in selected brain regions of rats. *Toxicology* **156**:13-25.

Snelling R and Nicholls D (1985) Calcium efflux and cycling across the synaptosomal plasma membrane. *Biochem J* **226**:225-231.

Weiser T and Wilson N (2002) Inhibition of tetrodotoxin (TTX)-resistant and TTX-sensitive neuronal Na⁺ channels by the secretolytic ambroxol. *Mol Pharmacol* **62**:433-438.

Wong PW and Pessah IN (1996) Ortho-substituted polychlorinated biphenyls alter calcium regulation by a ryanodine receptor-mediated mechanism: structural specificity toward skeletal- and cardiac-type microsomal calcium release channels. *Mol Pharmacol* **49**:740-751.

Yang JH and Kodavanti PR (2001) Possible molecular targets of halogenated aromatic hydrocarbons in neuronal cells. *Biochem Biophys Res Commun* **280**:1372-1377.

Footnotes:

This work was supported by MIUR (PRIN 2003) to S. A. and GF Di Renzo

Send reprint request to: Prof. Salvatore Amoroso, Unit of Pharmacology, Department of Neuroscience, School of Medicine, University "Politecnica delle Marche", Via Tronto 10/A, 60020 Torrette di Ancona, Ancona, Italy

Phone +39-071-2206176. Fax +39-071-2206178.

E-mail : s.amoroso@univpm.it

Legends for figures

Fig. 1. Effect of different concentrations of A1254 on $[Ca^{2+}]_i$ in the presence and in the absence of extracellular Ca^{2+} ions in SH-SY5Y cells.

SH-SY5Y cells were exposed to different concentrations (10, 30 and 50 $\mu\text{g/ml}$) of A1254, for 31 min. Each point represents the mean \pm SE (bars) of 4 values. In the experiments performed in the absence of extracellular Ca^{2+} ions, A1254 was added at the concentration of 30 $\mu\text{g/ml}$. The arrow indicates the minute at which A1254 was added. Inset shows a typical calibration trace in the presence of A1254 (30 $\mu\text{g/ml}$). $[Ca^{2+}]_i$ values were measured in sample containing 2×10^6 cells/2 ml.

Fig. 2. Effect of bepridil and the isothiourea derivative KB-R7943 on the $[Ca^{2+}]_i$ elevation induced by A1254. SH-SY5Y cells were exposed to a standard medium or to A1254 in the presence or in the absence of 10 μM bepridil or 10 μM KB-R7943. Each point represents the mean \pm SE (bars) of 4 values. The arrows indicate the minute at which NCX inhibitors and A1254 were added. $[Ca^{2+}]_i$ values were measured in sample containing 2×10^6 cells/2 ml.

Fig. 3. Effect of bepridil and KB-R7943 on A1254-induced neuronal injury. SH-SY5Y cells were exposed to A1254 (30 $\mu\text{g/ml}$) for 24 hrs in the presence or in the absence of 10 μM bepridil or 10 μM KB-R7943 and then assessed in their ability to produce formazan. Each column represents the mean \pm SE (bars) of 8 values.

F= 14.35 (one-way ANOVA)

*denotes statistical significance ($p < 0.001$) vs all other experimental groups

**denotes statistical significance ($p < 0.05$) vs control group

Fig. 4. RT-PCR analysis of SH-SY5Y mRNA expression of NCX1, NCX2 and NCX3 isoforms. In order to characterize the pattern of the expression of the three NCX isoforms in SH-SY5Y cells, RT-PCR analysis of NCX1, NCX2 and NCX3 mRNA expression was performed. Each lane shows PCR product obtained by using NCX1, NCX2 and NCX3 isoform primers.

Fig. 5. RT-PCR analysis of NCX1 and NCX3 mRNA expression in SH-SY5Y cells transfected with respective sense and antisense ODNs. Upper part of the figure shows RT-PCR of NCX1 and NCX3 products in SH-SY5Y transfected with the respective sense and antisense ODNs. Normalization of results was ensured by running parallel RT-PCR with glyceraldehyde-3-phosphate dehydrogenase (GAPDH) primers. Bottom part of the figure indicates optical density values representing the Mean \pm SE of three independent experiments.

Fig. 6. Effect of antisense NCX1 or NCX3 ODNs transfection on $[Ca^{2+}]_i$ increase and neuronal injury elicited by A1254.

A. SH-SY5Y naïve and antisense NCX1 or NCX3 ODNs transfected cells were exposed to A1254 (30 μ g/ml). Each point represents the mean \pm SE (bars) of 4 values. The arrow indicates the minute at which A1254 was added. $[Ca^{2+}]_i$ values were measured in sample containing 2×10^6 cell/2 ml.

B. SH-SY5Y naïve and antisense NCX1 or NCX3 ODNs transfected cells were exposed to A1254 (30 μ g/ml) and assessed in their ability to reduce MTT. Each column represents mean \pm SE (bars) of 8 values .

F = 14.38 (one-way ANOVA)

*p<0.001 vs control and AS1+A1254 groups.

** $p < 0.05$ vs control group; $p < 0.001$ vs A1254 group.

TABLE 1

NCX1, NCX2, NCX3 and GAPDH primers used in RT PCR reactions and chimeric phosphorothioated antisense (AS1, AS3) and sense (S1, S3) ODNs used to reduce the NCX isoforms expression. Bold indicates phosphorothioated bases.

GenBank accession number	Sequence (5' to 3')	Base pairs (bp)	Using
NCX1 X68191	sense: cccaagcttaatggagagaccaccaagac reverse: cgcgatccttgaagctggtctgtctcc	503	RT-PCR
NCX2 U08141	sense: cccaagcttgctgtgggcgatgctca reverse: cgcgatccgacctcgaggcgacagttc	528	RT-PCR
NCX3 U53420	sense: cccaagcttctggaagaggggatgaccc reverse: cgcgatccgttagggtgttcaccaata	516	RT-PCR
GAPDH	sense: ccatggagaaggctgggg reverse: caaagttgcatggatgacc	195	RT-PCR
NCX1	AS1: tgagactccaattggt		Transfection
NCX1	S1: aacaattggaagtctca		Transfection
NCX3	AS3: gccatacacaagag		Transfection
NCX3	S3: ctcttgtgatggc		Transfection

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6A




Fig. 6B

