

JPET #197483

Title Page:

Title: AT1 receptor antagonism is proangiogenic in the brain: BDNF a novel mediator

Authors: Ahmed Alhusban, Pharm.D., Anna Kozak, MS, Adviyee Ergul, MD, PhD,
Susan C Fagan*, Pharm.D.

Affiliation: Program in Clinical and Experimental Therapeutics- College of Pharmacy,
University of Georgia and Charlie Norwood VA Medical Center (AA, AK, AE, SCF),
Departments of Physiology (AE) and Neurology (SCF) , Georgia Health Sciences
University, Augusta, GA

JPET #197483

Running tile page:

Running title: BDNF mediates ARBs angiogenesis

Correspondence to:

Susan C Fagan, Pharm.D.

Jowdy Professor / Assistant Dean

University of Georgia College of Pharmacy

HM-1200; 1120 15th St.

Augusta, GA 30912

sfagan@georgiahealth.edu

Number of text pages: 30

Number of figures: 10

Number of supplemental figures: 4

References: 40

Number of words:

Abstract: 247

Introduction: 376

Discussion: 1810

JPET #197483

Abbreviations: AngII, angiotensin II; ARBs, angiotensin II receptor blockers; BDNF, brain derived neurotrophic factor; VEGF, vascular endothelial growth factor; cand, candesartan; GSK-3 β , Glycogen Synthase Kinase-3 β ; SHR, spontaneously hypertensive rats; hCMECs, human cerebromicrovascular endothelial cells. PD, PD-123319 (S-(+)-1-[4-(Dimethylamino)-3-methylphenyl)methyl]-5-(diphenylacetyl)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carboxylic acid di(trifluoroacetate) salt hydrate); K252a, (9S,10R,12R)-2,3,9,10,11,12-Hexahydro-10-hydroxy-9-methyl-1-oxo-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-l][1,6]benzodiazocine-10-carboxylic acid methyl ester; CGP-42112A, N α -Nicotinoyl-Tyr-(N α -Cbz-Arg)-Lys-His-Pro-Ile; GSK-3 β , Glycogen Synthase Kinase-3 β ; SHR, spontaneously hypertensive rats; hCMECs, human cerebromicrovascular endothelial cells.

Abstract

Candesartan is an angiotensin II type 1 receptor blocker (ARB) that has been shown to limit ischemic stroke and improve stroke outcome. In experimental stroke, candesartan induces a proangiogenic effect that is partly due to vascular endothelial growth factor (VEGF). Brain derived neurotrophic factor (BDNF) is a member of the neurotrophin family that has been reported to have angiogenic effects and play an important role in recovery after stroke. The purpose of this investigation was to determine the role of BDNF in the proangiogenic effect of candesartan in the brain under hypertensive conditions. Accordingly, spontaneously hypertensive rats were treated with candesartan and brain tissues were collected for quantification of BDNF expression. In addition, human cerebromicrovascular endothelial cells were treated with either low dose (1fM) or high dose (1μM) angiotensin II alone or in combination with candesartan (0.16μM) to assess the effect of candesartan treatment and BDNF involvement in the behavior of endothelial cells. Candesartan significantly increased the expression of BDNF in the SHR ($p < 0.05$). In addition, candesartan reversed the antiangiogenic effect of the 1μM dose of AngII ($p = 0.0001$). The observed effects of candesartan were ablated by neutralizing the effects of BDNF. Treatment with the AT2 antagonist PD-123319 significantly reduced tube-like formation in endothelial cells. AT2 stimulation induced the BDNF expression and migration ($p < 0.05$). In conclusion candesartan exerts a proangiogenic effect on brain microvascular endothelial cells treated with angiotensin II. This response is due to increased BDNF expression and is mediated through stimulation of the AT2 receptor.

JPET #197483

Introduction:

Angiotensin II type 1 receptor blockers (ARBs) have been shown to be vascular protective and seem to have a particularly robust effect in reducing the incidence of cerebrovascular events (Björn Dahlöf et al., 2002; Engelhorn et al., 2004; Kozak et al., 2009). Acutely, ARBs have been shown to improve outcome in experimental stroke (Fagan et al., 2006) and the long term functional benefit was accompanied by an augmented proangiogenic state (Kozak et al., 2009). This angiogenic effect was only partially attributed to an increase in VEGF expression (Kozak et al., 2009). Interestingly, the angiogenic response of candesartan was maintained even in non-stroked rats (Kozak et al., 2009). Subsequently, it was demonstrated that candesartan increased the expression of a number of genes for proangiogenic growth factors, including brain derived neurotrophic factor (BDNF), following experimental cerebral ischemia (Guan et al., 2011). BDNF is a member of the neurotrophin family that is expressed in a number of cell types and has been shown to have potent neurogenic, neuroprotective and angiogenic effects (Caporali and Emanuelli, 2009; Greenberg et al., 2009). The proangiogenic effects of ARBs are hotly debated, however, (Schieffer et al., 1994; Herr et al., 2008; Kozak et al., 2009) and may be tissue and situation dependent (Willis et al., 2011). In the brain, it is unclear whether the effects of ARBs are due to blood pressure lowering or a direct effect of candesartan on endothelial cells.

Glycogen synthase kinase- 3 β (GSK-3 β) is a serine threonine kinase that plays a key role in gene expression regulation (Wada, 2009; Hur and Zhou, 2010). Recently, Li *et al.* demonstrated the involvement of GSK-3 β in the recovery after CNS ischemic insults (Li et al., 2011). They demonstrated the involvement of GSK inhibition in regulating

JPET #197483

neural stem cells-endothelial cells cross talk (Li et al., 2011). This cross talk was found to be mediated through soluble growth factors like BDNF (Madri, 2009).

The purpose of this investigation was to determine whether candesartan-mediated blood pressure lowering increased BDNF protein expression in the brain in vivo and whether BDNF is involved in the proangiogenic effect of candesartan in vitro. In addition the involvement of GSK-3 β in candesartan mediated effects was assessed.

Materials and methods:

Animals:

The experimental protocol was approved by the Institutional Animal Care and Use Committee (IACUC) of the Charlie Norwood Veterans Affairs Medical Center (09-04-008). Male spontaneously hypertensive rats (SHRs) (280-300g, n=4-6 per group) were subjected to middle cerebral artery occlusion (MCAO) sham surgery and randomized to receive a single IV dose of candesartan 0.3mg/kg, hydralazine 1mg/kg, or saline. In addition, male wistar rats (280-300g, n=4 per group) were subjected to the same surgery and randomized to receive either a single IV dose of candesartan 1mg/kg or saline. Twenty four hours later the rats were euthanized and the brains were harvested and flash frozen in liquid nitrogen.

Western blotting:

To assess BDNF expression, the right and left hemispheres were separated and processed as described previously (Guan et al., 2011) and the blots probed with

JPET #197483

antiBDNF (1:250, abcam) and β -Actin (1:10000, sigma). For the in vitro experiments; human cerebrovascular endothelial cells (hCMECs) were cultured to confluence and serum starved for 10 hours followed by incubations with either 1fM or 1 μ M angiotensin II (AngII). After 6 hours, candesartan 0.16 μ M was added to the cells and incubated for 16 hours and then homogenized and processed for immunoblotting. To assess the involvement of AT2 receptor in BDNF expression HCMECs were serum starved for 16 hours and pretreated with PD-123319(0.1 μ M) 30 minutes before being incubated with AngII (1fM or 1 μ M) for 6 hours. Candesartan or vehicle were introduced in the media for 10 hours. To further confirm AT2 involvement, cells were serum starved for 16 hours followed by treatment with either the AT2 agonist CGP-121141A (0.1 μ M) or vehicle for 16 hours. The expression of AT1, AT2 and the phosphorylation status of GSK-3 β were assessed using the same above treatment paradigm. Blots were probed with mouse monoclonal AT1 antibody (Abcam; 1:1000); rabbit monoclonal AT2 antibody (Abcam; 1:1000); p-S9GSK-3 β (cell signaling; 1:1000) and total GSK-3 β (cell signaling; 1:1000) Protein expression was quantified as the relative optic density of the protein band normalized to actin using NIH-image J software.

Cell culture:

hCMECs were provided as a generous gift from Dr. J. Zastre (UGA College of Pharmacy). hCMECs were cultured in minimum essential media (ATCC) supplemented with EGM-2 SingleQuot Kit Suppl. & Growth Factors (Lonza) and 10% FBS (Atlanta Biologicals) and p30-34 were used in the experiments.

JPET #197483

Treatments:

Candesartan was provided as a generous gift from Astra-Zeneca. Hydralazine was purchased from Sigma-Aldrich (St. Louis, MO) and was reconstituted with 0.9% normal saline. AngII was purchased from Sigma-Aldrich (St. Louis, MO) and was reconstituted and diluted to the desired concentration using serum free media. BDNF neutralization was achieved using 100 nM K252a (Trk receptor inhibitor) dissolved in 25% DMSO both purchased from Sigma-Aldrich (St. Louis, MO), 10ng/ml anti-BDNF neutralizing antibody (Abcam; Cambridge, MA) and 0.4ug/ml TrkB-Fc (soluble BDNF receptor chimera. R&D systems; Minneapolis, MN). The involvement of AT2 receptor was assessed using the AT2 antagonist PD-123319 (0.1 μ M) and the AT2 receptor agonist CGP-42112A (0.1 μ M) both purchased from Sigma-Aldrich (Sigma-Aldrich; St. Louis, MO). All the inhibitors were added to the media 30 minutes before AngII treatment.

Dose and time study:

hCMECs were cultured until confluence and serum starved for 10 hours . Cells were incubated with six different concentrations of AngII (0-1 μ M) for 2, 6and8 hours and the cells were homogenized and processed for immunoblotting. BDNF expression was quantified as the relative density of the BDNF band to the corresponding β -actin or GAPDH bands. The calculated relative density was normalized to the relative density of the control band, and reported as fold change.

Proliferation assay:

JPET #197483

Proliferation was assessed using BrdU incorporation (Cell Proliferation ELISA, BrdU (colorimetric); Roche Applied Science) according to the manufacturer recommendations. Briefly, 5000 hCMECs were seeded into each well of a 96 well plate and left to attach for 24 hours. Cells were serum starved for 10 hours and then treated with AngII (1fM or 1μM) for 6 hours. Following 6 hours, cells were treated with different combinations of candesartan, antiBDNF, TrkB-Fc, K252a, DMSO or IgG and incubated for 18 hours. The cells were then labeled with BrdU for 4 hours and then processed to quantify BrdU incorporation.

Angiogenesis Assays

Cell migration:

Wound recovery assay was used to assess cell migration where hCMECs were cultured in a 12 well plate to confluence and then serum starved for 10 hours followed by 6 hours of AngII (1fM or 1μM) treatment. A wound was introduced in the monolayer of endothelial cells and the cells treated with AngII (1fM or 1μM) with different combinations of candesartan, antiBDNF, TrkB-Fc, K252a, DMSO or IgG. In some experiments, cells were pretreated with PD-123319 (0.1μM) 30 minutes before AngII treatment. In another set of studies cells were treated with CGP-42112A (0.1μM), candesartan (0.16μM) or their combination. Scratch recovery was assessed by taking images of the scratch at baseline and at 16 hours after scratch introduction and the width of the scratch was measured at both time points using NIH image J software. The percentage wound recovery was calculated as the percent decrease in scratch width at

JPET #197483

16 hours and the data was presented as percentage scratch recovery as compared to the control.

Tube formation:

2×10^4 hCMECs were suspended in serum free media and mixed with matrigel (BD Biosciences; San Jose, CA) in a 60:30 ratio and plated in a 96 well plate and then treated with different combinations of AngII (1fM or 1 μ M, sigma), candesartan, antiBDNF, TrkB-Fc, K252a, DMSO, IgG ,or PD-123319. Tube like structure formation was assessed using a digital camera attached to an Olympus microscope. Three images from each well were photographed at 24 hours and the number of tube like structures was quantified.

Statistical analysis:

All experiments were repeated three times in triplicate and data was quantified in a blinded manner. Statistical significance was detected by one-way ANOVA for in vitro data followed by post-hoc Tukey test. Unpaired t-test was used to determine the significance of BDNF expression in the right and left brain hemispheres of candesartan treated SHRs as compared to the corresponding hemisphere in saline treated animals. Statistical analyses were performed using GraphPad prism software (5.1). $P < 0.05$ was considered significant.

Results:

Candesartan increases the expression of BDNF in SHR brain:

JPET #197483

BDNF has been shown to exert a beneficial effect in a variety of CNS pathologies (Greenberg et al., 2009); the direct use of BDNF in therapy is limited by its pharmacokinetic profile (Greenberg et al., 2009). A plausible alternative approach is to use either synthetic BDNF mimetics or agents that can induce BDNF expression in the brain. Treatment with a single dose of candesartan (0.3mg/kg) dramatically reduced the blood pressure from 150 mmHg at baseline to 120 mmHg after treatment (Figure 1A). Candesartan treatment significantly increased the expression of BDNF in both right and left hemisphere of SHR animals 24 hours after sham surgery (Figure 1B). The effect of candesartan on BDNF expression was maintained in wistar rats (supplemental data S3). In contrast, hydralazine treatment did not affect BDNF expression in SHRs (supplemental data S4).

Angiotensin II modulates the expression of BDNF in hCMECs.

Angiotensin II has been reported to affect the expression of BDNF in the adrenals (Szekeres et al., 2010) and brain (Chan et al., 2010); but its effect on BDNF expression in endothelial cells has never been reported. In hCMECs treatment with AngII was found to induce a dose and time-dependent modulation of BDNF expression that was maximal at 6 hours after incubation with either AngII 1fM or 1µM (Figure 2A). These two concentrations were used in the following experiments.

Candesartan increases BDNF expression in hCMECs:

After establishing the effect of ARBs on the expression of BDNF in brain tissue in vivo, the effect of candesartan on the expression of BDNF in AngII treated cells in vitro was investigated. Candesartan significantly increased the expression of BDNF in hCMECs

JPET #197483

treated with AngII 1fM in comparison to AngII 1fM alone and in hCMECs treated with both concentrations of AngII as compared to the control as assessed after 16 hours of incubation with candesartan (Figure 2B).

Candesartan has a dose dependent modulatory effect on the angiogenic potential of hCMECs.

Data from clinical studies in stroke presented conflicting results on whether the hypotensive effect of ARBs is an essential requirement for its reported benefit (Schrader et al., 2003; Schrader et al., 2005). Consequently, we were interested in assessing whether a therapeutically relevant concentration of candesartan would have an in vitro effect in hCMECs in the absence of AngII. We calculated the amount of candesartan that can give a concentration similar to that achieved in patients receiving the drug (0.16 μ M). The angiogenic effect of a range of candesartan concentrations, including the concentration under consideration, was assessed using an in vitro matrigel tube formation assay (Figure 2C). The observed response was further confirmed using the wound recovery migration assay (Figure 2B). Candesartan induced a dose dependent, bell-shaped modulatory effect on the tube formation rate in hCMECs. The 0.16 μ M candesartan concentration (therapeutically relevant) significantly increased the tube formation rate, whereas the other concentrations did not affect the rate of tube formation (Figure 2C). A similar effect was observed in the wound recovery migration assay, except for an increased migration rate in hCMECs treated with candesartan 1.6 μ M concentration (Figure 2D). This differential effect of the 1.6 μ M candesartan concentration might be attributed to dose dependent effects of candesartan on the different processes involved in migration and tube formation.

JPET #197483

Candesartan modulates the proliferation of hCMECs:

Angiotensin II has been reported to induce the proliferation of cells in vitro (Kou et al., 2007; Herr et al., 2008). This proliferative effect has been shown to be blocked with ARBs (Kou et al., 2007; Herr et al., 2008); but the reported concentration of ARBs in these studies was supratherapeutic (Herr et al., 2008). Consequently, we attempted to assess the effect of the therapeutically relevant concentration on the proliferative effect of AngII in hCMECs. Both low and high concentrations of AngII significantly increased the proliferation of hCMECs (Figures 3A and D). Treatment with candesartan maintained and further enhanced this proliferative effect (Figures 3A and D).

Candesartan modulates the migration rate of hCMECs:

Similar to its proliferative effect, AngII has been reported to affect the migration of cells in vitro (Buharalioglu et al., 2011). Since we have found a proliferative effect of candesartan on AngII treated hCMECs, we were interested in assessing the effect of this dose of candesartan on two critical steps of angiogenesis: migration and tube formation in AngII treated hCMECs. We observed a significant increase in the migration of hCMECs in response to treatment with AngII 1 fM (Figure 4A). Interestingly, the higher AngII concentration did not have an effect on the migration of hCMECs (Figure 4D) in our model. While candesartan maintained the increased hCMECs migration in the low dose AngII group (Figure 4A), candesartan increased hCMECs migration in the high dose AngII dose treated cells (Figure 4D). Cells migration was not affected by K252a, DMSO or IgG (Figures 4C and F).

Candesartan has a proangiogenic effect in hCMECs:

JPET #197483

The pro-angiogenic effect of AngII has been reported previously (Hu et al., 2007; Buharalioglu et al., 2011). In hCMECs there was a significantly increased rate of tube formation in response to AngII 1fM treatment (Figure 5A) and a reduction in tube formation in AngII 1μM treated cells (Figure 5D). Interestingly, candesartan maintained AngII 1fM induced tube formation and reversed the antiangiogenic effect of AngII 1μM (Figure 5A and D).

BDNF mediates the effects of candesartan on hCMECs:

Our in vivo results demonstrated the ability of candesartan to increase the expression of BDNF in SHR's brain. Previously, BDNF has been demonstrated to have an angiogenic effect which shifted our interest to assess whether BDNF is involved in the observed effects of candesartan. Upon neutralizing BDNF using three different methods of BDNF neutralization, the observed effects of candesartan were consistently ablated (Figures 3B and E; 4B and E, 5B and E); a finding that identifies BDNF as an essential mediator of candesartan effects in hCMECs. Interestingly, the angiogenic effect of AngII 1fM was significantly inhibited upon BDNF neutralization (Figure 5C), which suggests the involvement of BDNF in AngII induced angiogenesis, which is confirmed by the robust angiogenic response to BDNF treatment (Figures 5C and F).

AT2 receptor mediates the angiogenic response to angiotensin II in hCMECs.

Candesartan mediates its effects through AT1 blockade(Engelhorn et al., 2004; Willis et al., 2011), which induces an unopposed stimulation of AT2 receptor(Hashikawa-Hobara et al., 2012). Findings in this investigation suggest a possible involvement of AT2 receptor in mediating candesartan effects.

JPET #197483

AT2 blockade using PD-123319 significantly inhibited the angiogenic response induced by either angiotensin II or the combination of angiotensin II and candesartan (Figure 6 A and B). In addition the AT2 agonist, CGP-24112A, significantly increased the migration of hCMECs. This migratory response was not affected by the concomitant treatment with candesartan (Figure 6C). These findings demonstrate an essential role of AT2 in the angiogenic process of hCMECs.

Candesartan induced BDNF expression is mediated through the AT2 receptor.

Findings in this study demonstrated the ability of candesartan to increase the expression of BDNF both in vivo and in vitro; in addition the in vitro effects of candesartan were shown to be mediated through the AT2 receptor. Accordingly, the involvement of AT2 receptor in BDNF expression was evaluated. As has been demonstrated earlier candesartan significantly increased the expression of BDNF in AngII treated cells (Figure 2B, 7A). Pretreatment with the AT2 antagonist PD-123319 significantly inhibited candesartan induced BDNF expression, which suggests the involvement of AT2 receptor. To confirm this findings hCMECs were treated with the AT2 agonist CGP-42112A (0.1 μ M) and the expression of BDNF was assessed. CGP-42112A significantly increased t expression of BDNF in hCMECs by about one fold as compared to vehicle treated cells (Figure 7B).

AT1 antagonism regulates the expression of AT1 in an AT2 dependent manner.

Candesartan treatment significantly increased the expression of AT1 receptor in cells treated with high dose of AngII as compared to both control and AngII alone. This

JPET #197483

response was ablated when the AT2 receptor was blocked using PD-123319 (Figure 7B). Additionally, the AT2 agonist CGP-24112A induced a significant fourfold increase in AT1 receptor expression in hCMECs (supplementary data). In cells treated with low dose AngII, candesartan significantly increased the expression of AT1 compared to control alone (Figure 7A). The expression of AT2 was not changed under the different treatments used (Figure 7 C and D).

AT1 antagonism modulates the activity of GSK-3 β in an AT2 dependent manner.

GSK-3 β was found to modulate BDNF expression (Hur and Zhou, 2010; Li et al., 2011). Our results demonstrated the ability of candesartan to increase the expression of BDNF in both in vivo and in vitro settings. Accordingly we were interested in assessing the activity of GSK-3 β under the different treatment conditions we were using. Candesartan treatment significantly increased the phosphorylation of GSK-3 β at the inhibitory serine 9 residue in cells treated with high dose AngII (Figure 8A). The higher phosphorylation was reversed by the concomitant treatment with AT2 blocker PD-123319 (Figure 8A).

Discussion:

Our results demonstrate, for the first time, the ability of candesartan to promote a proangiogenic state in hCMECs in an AngII concentration independent manner, where candesartan was able to maintain the angiogenic effect of AngII at the low dose, and reverse the antiangiogenic effect of the high dose. The proangiogenic effect of candesartan was found to be mainly dependent on BDNF and is mediated through AngII stimulation of the AT2 receptor. In addition, we have demonstrated the ability of

JPET #197483

candesartan to induce an angiogenic response in endothelial cells; even in the absence of exogenously added AngII.

Following an ischemic insult in the brain, induction of angiogenesis has been shown to ameliorate the damage and is coupled to neurogenesis leading to enhanced recovery and better outcome (Navaratna et al., 2009; Xiong et al., 2010). ARBs administration was found to increase vascular density in heart and brain following MI (Sladek et al., 1996) and stroke (Kozak et al., 2009; Guan et al., 2011) which was attributed to an increase in VEGF expression in stroke (Guan et al., 2011). In contrast to in vivo data, in vitro angiogenesis studies demonstrated an antiangiogenic effect of ARBs in endothelial cells treated with AngII (Hu et al., 2007; Herr et al., 2008). This antiangiogenic effect was mediated mainly through inhibiting AngII induced increase in VEGF signaling (Fujiyama et al., 2001; Herr et al., 2008). The majority of these studies were conducted in either HUVECs (Kou et al., 2007; Buharalioglu et al., 2011) or coronary artery endothelial cells (Fujiyama et al., 2001; Hu et al., 2007) and none of them evaluated the effect of AngII or AngII and ARBs in hCMECs which are phenotypically different from other endothelial cells (Feletou, 2011). In addition, the majority of the studies focused on the involvement of VEGF (Buharalioglu et al., 2011), angiopoietins (Herr et al., 2008) and EGFR transactivation (Fujiyama et al., 2001) in the angiogenic response to AngII. This ignores the possible role of other angiogenic mediators, like BDNF, which has been shown to induce VEGF expression (Caporali and Emanuelli, 2009) and produce an angiogenic response comparable to that of VEGF in endothelial cells (Li et al., 2006). Results from this study demonstrate a dose dependent effect of AngII on brain angiogenesis, where low concentrations of AngII induce angiogenesis and higher doses

JPET #197483

inhibit it. This is consistent with that shown in other vascular beds (Kou et al., 2007; Buharalioglu et al., 2011).

The design of this study is unique in many aspects. In a typical *in vitro* study, cells are initially treated with candesartan followed by AngII treatment. This precludes detection of a possible AngII independent interaction between candesartan and endothelial cells. This design has limited clinical relevance since candesartan will be introduced to the system in response to the effects of AngII or other circulating vasoactive mediators. In our design, we attempted to model the temporal relationship of treatment introduction. Data about the effect of AngII on BDNF expression in endothelial cells were lacking and the only reports addressing the effect of AngII on BDNF expression were in the adrenal cortex (Szekeres et al., 2010) and the brain (Chan et al., 2010). Accordingly, we did a time and dose response study in hCMECs to determine the incubation time and AngII dose to be used in the *in vitro* studies. This dose and time response study was followed by assessing the effect of those AngII on the viability of hCMECs (supplementary data S1). Also, the concentration of candesartan used in the *in vitro* studies was calculated to produce the midpoint therapeutic steady state concentration in humans (Schulz and Schmoldt, 2003). The calculated concentration was tested for its angiogenic and migratory effect and compared to other candesartan doses in a dose response curve (Figures 2C and D). The *in vivo* dose was determined based on previously reported data demonstrating neurovascular protection in SHR (Kozak et al., 2008).

Our findings support and expand our previously reported data on the ability of cerebral spinal fluid (CSF) from candesartan treated non-stroked animals to induce angiogenesis in endothelial cells (Kozak et al., 2009). This proangiogenic effect of candesartan was

JPET #197483

only partially attributed to VEGF (Fagan et al., 2006), suggesting the involvement of other angiogenic factors. In this study we have identified BDNF as another important mediator of the angiogenic effect of candesartan. BDNF neutralization using receptor inhibitor K252a, antiBDNF antibody or TrkB-Fc almost ablated the effects of candesartan in all reported assays.

The proliferative and angiogenic effects of AngII have been largely attributed to AT1 mediated signaling (Otani et al., 1998; Fujiyama et al., 2001; Hu et al., 2007; Herr et al., 2008; Buharalioglu et al., 2011) while AT2 mediated signaling was thought to either counteract AT1 induced angiogenesis (Fujiyama et al., 2001; Javier Carbajo-Lozoya and Hans-Peter Hammes, 2012) or not have an effect on AngII induced angiogenesis (Otani et al., 1998; Hu et al., 2007). In contrast to this notion, AT2 mediated signaling has been demonstrated to promote angiogenesis in ischemic myocardium (Munk et al., 2007) and retinal endothelial cells (Sarlos et al., 2003). In this study, AT2 mediated signaling was found to be largely responsible for the angiogenic response in hCMECs. AT2 involvement in angiogenesis was initially suggested by the lack of effect on AngII – mediated angiogenesis when AT1 was blocked using candesartan and was further confirmed when the angiogenic response was totally prevented upon AT2 blockade using PD-123319. This finding suggests the importance of unopposed AT2 stimulation following stroke (McCarthy et al., 2009; Oprisiu-Fournier R, 2009) and may explain the lack of protective effect of ARBs in the absence of AT2 signaling following cerebral ischemia (Iwai et al., 2004; Li et al., 2005; Lu et al., 2005; Faure et al., 2008) . Our data demonstrates the indispensable role of AT2 for angiogenesis in hCMECs which has

JPET #197483

been linked to neurogenesis and improving recovery after CNS ischemic insults (Madri, 2009; Navaratna et al., 2009; Xiong et al., 2010).

The activity of glycogen synthase kinase 3 beta (GSK-3 β) has been shown to be involved in the expression of neurotrophins in the brain (Wada, 2009) and in the cross talk between endothelial and neural stem cells through regulating BDNF and VEGF expression (Li et al., 2011). In addition GSK-3 β has been recently demonstrated to regulate the angiogenic response in endothelial cells (Flugel et al., 2012). Data from this study highlights a possible involvement of GSK-3 β in mediating the effects of AT1 antagonism in hCMECs in an Ang II dose-dependent manner. Candesartan increased phosphorylation of GSK-3 β at the inhibitory serine 9 residue which will inhibit the activity of GSK-3 β when used alone or in combination with 1 μ M AngII the concentration that induced antiangiogenic effects in hCMECs. Concomitantly this increased inhibition of GSK-3 β was associated with increased BDNF expression and angiogenic response in hCMECs. These findings are consistent with previously published data about the interaction between GSK-3 β and BDNF expression (Wada, 2009; Li et al., 2011).

In this study, candesartan demonstrated the ability to modulate the angiogenic response of hCMECs in the absence of exogenously added AngII. This effect was found to be dose dependent. The therapeutically relevant candesartan concentration induces a proangiogenic effect; whereas the other two concentrations didn't affect the angiogenic potential of the cells. This finding may help explain the controversial angiogenic effect of ARBs between the in vivo and the in vitro data (Willis et al., 2011). Previously published reports on the in vitro antiangiogenic effect of ARBs used supra-therapeutic concentrations (Hu et al., 2007; Herr et al., 2008). As demonstrated by the findings of

JPET #197483

this study, the supra-therapeutic doses of ARBs might have direct antiangiogenic effects on endothelial cells.

Another unique aspect of this work is the demonstration of candesartan's ability to increase the expression of BDNF in both hCMECs and the brain tissue of hypertensive animals. Hashikawa-Hobara et al. recently demonstrated the ability of candesartan to stimulate neurite growth in an AT2 dependent manner through Akt signaling (Hashikawa-Hobara et al., 2012). This is consistent with our findings, highlighting the involvement of the AT2 receptor in BDNF expression which is known to stimulate neurite growth (Parrish et al., 2007) and Akt signaling (Caporali and Emanuelli, 2009). Interestingly, the effect of candesartan on BDNF expression appears to be independent of blood pressure lowering. Candesartan increased BDNF expression in wistar rats (supplemental data S3); whereas, hydralazine had no effect on BDNF expression in SHRs (supplemental data S4).

An interesting finding in this study is the involvement of AT2 in AT1 expression. Cross-talk between AngII receptors have been previously reported at both expression and functional levels (De Paolis et al., 1999; Saavedra, 1999; Miura et al., 2010). These reports consistently demonstrated the involvement of AT1 in AT2 expression (De Paolis et al., 1999). In addition, it has been shown that AT2 mediated signaling antagonizes some aspects of AT1 mediated effects (Miura et al., 2010). Our results demonstrate for the first time that AT1 expression in hCMECs is positively regulated by AT2 mediated signaling.

JPET #197483

In this study all efforts were made to confirm each result using different methods but the following limitations can be identified; the in vitro studies were conducted in an endothelial cell line rather than primary endothelial cells. In addition, BDNF neutralization studies were performed using mainly pharmacologic methods although a genetic approach using RNA interference would provide more power to the study. In addition, in our study we have used human derived cell line in our in vitro study while using a murine model as an in vivo model. In our research the main goal and focus is to model and understand the changes that accompany ischemic stroke in humans. Because of the inability to directly probe human brain samples, we are using rats to study the in vivo changes in response to candesartan or any pharmacologic agent of interest. In addition whenever we had the chance to use human derived tissues or cells we do use them to give a better understanding of what changes are taking place. In this investigation we employed human cerebrovascular endothelial cells as an in vitro system and we used SHRs as an in vivo system.

In conclusion, our findings demonstrate the ability of candesartan to modulate the behavior of endothelial cells to promote a proangiogenic state. In hCMECs, this modulatory effect of candesartan can be largely attributed to BDNF and is mediated through the AT2 receptor. In addition, the dose dependent proangiogenic effect of candesartan and even the mechanism involved may help explain the disparate findings on the angiogenic potential of ARBs that prevails in the biomedical literature.

JPET #197483

JPET #197483

Authorship contributions:

Participated in research design: Alhusban, Ergul, and Fagan.

Conducted experiments: Alhusban, and Kozak.

Performed data analysis: Alhusban, Ergul, and Fagan.

**Wrote or contributed to the writing of the manuscript: Alhusban, Kozak, Ergul,
and Fagan.**

References

- Björn Dahlöf RBD, Sverre E Kjeldsen, Stevo Julius, Gareth Beevers, Ulf de Faire, Frej Fyhrquist, Hans Ibsen, Krister Kristiansson OL-P, Lars H Lindholm, Markku S Nieminen, Per Omvik, Suzanne Oparil, and Hans Wedel ftLsg (2002) Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* **359**:8.
- Buharalioglu CK, Song CY, Yaghini FA, Ghafoor HU, Motiwala M, Adris T, Estes AM and Malik KU (2011) Angiotensin II-induced process of angiogenesis is mediated by spleen tyrosine kinase via VEGF receptor-1 phosphorylation. *Am J Physiol Heart Circ Physiol* **301**:H1043-1055.
- Caporali A and Emanuelli C (2009) Cardiovascular actions of neurotrophins. *Physiol Rev* **89**:279-308.
- Chan SH, Wu CW, Chang AY, Hsu KS and Chan JY (2010) Transcriptional upregulation of brain-derived neurotrophic factor in rostral ventrolateral medulla by angiotensin II: significance in superoxide homeostasis and neural regulation of arterial pressure. *Circ Res* **107**:1127-1139.
- De Paolis P, Porcellini A, Gigante B, Giliberti R, Lombardi A, Savoia C, Rubattu S and Volpe M (1999) Modulation of the AT₂ subtype receptor gene activation and expression by the AT₁ receptor in endothelial cells. *J Hypertens* **17**:1873-1877.
- Engelhorn T, Goerike S, Doerfler A, Okorn C, Forsting M, Heusch G and Schulz R (2004) The angiotensin II type 1-receptor blocker candesartan increases cerebral blood flow, reduces infarct size, and improves neurologic outcome after transient cerebral ischemia in rats. *J Cereb Blood Flow Metab* **24**:467-474.
- Fagan SC, Kozak A, Hill WD, Pollock DM, Xu L, Johnson MH, Ergul A and Hess DC (2006) Hypertension after experimental cerebral ischemia: candesartan provides neurovascular protection. *J Hypertens* **24**:535-539.
- Faure S, Bureau A, Oudart N, Javellaud J, Fournier A and Achard JM (2008) Protective effect of candesartan in experimental ischemic stroke in the rat mediated by AT₂ and AT₄ receptors. *J Hypertens* **26**:2008-2015.
- Feletou M (2011) Series on Integrated Systems Physiology: From Molecule to Function to Disease., in *The Endothelium: Part 1: Multiple Functions of the Endothelial Cells—Focus on Endothelium-Derived Vasoactive Mediators* (Granger DN GJ ed), Morgan & Claypool Life Sciences, San Rafael (CA).
- Flugel D, Gorchach A and Kietzmann T (2012) GSK-3 β regulates cell growth, migration, and angiogenesis via Fbw7 and USP28-dependent degradation of HIF-1 α . *Blood* **119**:1292-1301.
- Fujiyama S, Matsubara H, Nozawa Y, Maruyama K, Mori Y, Tsutsumi Y, Masaki H, Uchiyama Y, Koyama Y, Nose A, Iba O, Tateishi E, Ogata N, Jyo N, Higashiyama S and Iwasaka T (2001) Angiotensin AT₁ and AT₂ receptors differentially regulate angiopoietin-2 and vascular endothelial growth factor expression and angiogenesis by modulating heparin binding-epidermal growth factor (EGF)-mediated EGF receptor transactivation. *Circ Res* **88**:22-29.

- Greenberg ME, Xu B, Lu B and Hempstead BL (2009) New insights in the biology of BDNF synthesis and release: implications in CNS function. *J Neurosci* **29**:12764-12767.
- Guan W, Somanath PR, Kozak A, Goc A, El-Remessy AB, Ergul A, Johnson MH, Alhusban A, Soliman S and Fagan SC (2011) Vascular protection by angiotensin receptor antagonism involves differential VEGF expression in both hemispheres after experimental stroke. *PLoS One* **6**:e24551.
- Hashikawa-Hobara N, Hashikawa N, Inoue Y, Sanda H, Zamami Y, Takatori S and Kawasaki H (2012) Candesartan Cilexetil Improves Angiotensin II Type 2 Receptor-Mediated Neurite Outgrowth via the PI3K-Akt Pathway in Fructose-Induced Insulin-Resistant Rats. *Diabetes* **61**:925-932.
- Herr D, Rodewald M, Fraser HM, Hack G, Konrad R, Kreienberg R and Wulff C (2008) Regulation of endothelial proliferation by the renin-angiotensin system in human umbilical vein endothelial cells. *Reproduction* **136**:125-130.
- Hu C, Dandapat A and Mehta JL (2007) Angiotensin II induces capillary formation from endothelial cells via the LOX-1 dependent redox-sensitive pathway. *Hypertension* **50**:952-957.
- Hur EM and Zhou FQ (2010) GSK3 signalling in neural development. *Nat Rev Neurosci* **11**:539-551.
- Iwai M, Liu HW, Chen R, Ide A, Okamoto S, Hata R, Sakanaka M, Shiuchi T and Horiuchi M (2004) Possible inhibition of focal cerebral ischemia by angiotensin II type 2 receptor stimulation. *Circulation* **110**:843-848.
- Javier Carbajo-Lozoya SL, Yuxi Feng, Jens Kroll, and Hans-Peter Hammes TW (2012) Angiotensin II modulates VEGF-driven angiogenesis by opposing effects of type 1 and type 2 receptor stimulation in the microvascular endothelium. *Cellular Signalling* **24**:1261-1269.
- Kou B, Vatish M and Singer DR (2007) Effects of angiotensin II on human endothelial cells survival signalling pathways and its angiogenic response. *Vascul Pharmacol* **47**:199-208.
- Kozak A, Ergul A, El-Remessy AB, Johnson MH, Machado LS, Elewa HF, Abdelsaid M, Wiley DC and Fagan SC (2009) Candesartan augments ischemia-induced proangiogenic state and results in sustained improvement after stroke. *Stroke* **40**:1870-1876.
- Kozak W, Kozak A, Johnson MH, Elewa HF and Fagan SC (2008) Vascular protection with candesartan after experimental acute stroke in hypertensive rats: a dose-response study. *J Pharmacol Exp Ther* **326**:773-782.
- Li J, Culman J, Hortnagl H, Zhao Y, Gerova N, Timm M, Blume A, Zimmermann M, Seidel K, Dirnagl U and Unger T (2005) Angiotensin AT2 receptor protects against cerebral ischemia-induced neuronal injury. *FASEB J* **19**:617-619.
- Li Q, Ford MC, Lavik EB and Madri JA (2006) Modeling the neurovascular niche: VEGF- and BDNF-mediated cross-talk between neural stem cells and endothelial cells: an in vitro study. *J Neurosci Res* **84**:1656-1668.
- Li Q, Michaud M, Canosa S, Kuo A and Madri JA (2011) GSK-3beta: a signaling pathway node modulating neural stem cell and endothelial cell interactions. *Angiogenesis* **14**:173-185.

JPET #197483

- Lu Q, Zhu YZ and Wong PT (2005) Neuroprotective effects of candesartan against cerebral ischemia in spontaneously hypertensive rats. *Neuroreport* **16**:1963-1967.
- Madri JA (2009) Modeling the neurovascular niche: implications for recovery from CNS injury. *J Physiol Pharmacol* **60 Suppl 4**:95-104.
- McCarthy CA, Vinh A, Callaway JK and Widdop RE (2009) Angiotensin AT2 receptor stimulation causes neuroprotection in a conscious rat model of stroke. *Stroke* **40**:1482-1489.
- Miura S, Matsuo Y, Kiya Y, Karnik SS and Saku K (2010) Molecular mechanisms of the antagonistic action between AT1 and AT2 receptors. *Biochem Biophys Res Commun* **391**:85-90.
- Munk VC, Sanchez de Miguel L, Petrimpol M, Butz N, Banfi A, Eriksson U, Hein L, Humar R and Battegay EJ (2007) Angiotensin II induces angiogenesis in the hypoxic adult mouse heart in vitro through an AT2-B2 receptor pathway. *Hypertension* **49**:1178-1185.
- Navaratna D, Guo S, Arai K and Lo EH (2009) Mechanisms and targets for angiogenic therapy after stroke. *Cell Adh Migr* **3**:216-223.
- Oprisiu-Fournier R FS, Mazouz H, Boutitie F, Serot JM, Achard JM, Godefroy O, Hanon O, Temmar M, Albu A, Strandgaard S, Wang J, Black SE, Fournier A. (2009) Angiotensin AT1-receptor blockers and cerebrovascular protection: do they actually have a cutting edge over angiotensin-converting enzyme inhibitors? *Expert Rev Neurother* **9**.
- Otani A, Takagi H, Suzuma K and Honda Y (1998) Angiotensin II potentiates vascular endothelial growth factor-induced angiogenic activity in retinal microcapillary endothelial cells. *Circ Res* **82**:619-628.
- Parrish JZ, Emoto K, Kim MD and Jan YN (2007) Mechanisms that regulate establishment, maintenance, and remodeling of dendritic fields. *Annu Rev Neurosci* **30**:399-423.
- Saavedra JM (1999) Emerging features of brain angiotensin receptors. *Regul Pept* **85**:31-45.
- Sarlos S, Rizkalla B, Moravski CJ, Cao Z, Cooper ME and Wilkinson-Berka JL (2003) Retinal angiogenesis is mediated by an interaction between the angiotensin type 2 receptor, VEGF, and angiopoietin. *Am J Pathol* **163**:879-887.
- Schieffer B, Wirger A, Meybrunn M, Seitz S, Holtz J, Riede UN and Drexler H (1994) Comparative effects of chronic angiotensin-converting enzyme inhibition and angiotensin II type 1 receptor blockade on cardiac remodeling after myocardial infarction in the rat. *Circulation* **89**:2273-2282.
- Schrader J, Luders S, Kulschewski A, Berger J, Zidek W, Treib J, Einhaupl K, Diener HC and Dominiak P (2003) The ACCESS Study: evaluation of Acute Candesartan Cilexetil Therapy in Stroke Survivors. *Stroke* **34**:1699-1703.
- Schrader J, Luders S, Kulschewski A, Hammersen F, Plate K, Berger J, Zidek W, Dominiak P and Diener HC (2005) Morbidity and Mortality After Stroke, Eprosartan Compared with Nitrendipine for Secondary Prevention: principal results of a prospective randomized controlled study (MOSES). *Stroke* **36**:1218-1226.

JPET #197483

- Schulz M and Schmoltdt A (2003) Therapeutic and toxic blood concentrations of more than 800 drugs and other xenobiotics. *Pharmazie* **58**:447-474.
- Sladek T, Sladkova J, Kolar F, Papousek F, Cicutti N, Korecky B and Rakusan K (1996) The effect of AT1 receptor antagonist on chronic cardiac response to coronary artery ligation in rats. *Cardiovasc Res* **31**:568-576.
- Szekeres M, Nadasy GL, Turu G, Supeki K, Szidonya L, Buday L, Chaplin T, Clark AJ and Hunyady L (2010) Angiotensin II-induced expression of brain-derived neurotrophic factor in human and rat adrenocortical cells. *Endocrinology* **151**:1695-1703.
- Wada A (2009) Lithium and neuropsychiatric therapeutics: neuroplasticity via glycogen synthase kinase-3beta, beta-catenin, and neurotrophin cascades. *J Pharmacol Sci* **110**:14-28.
- Willis LM, El-Remessy AB, Somanath PR, Deremer DL and Fagan SC (2011) Angiotensin receptor blockers and angiogenesis: clinical and experimental evidence. *Clin Sci (Lond)* **120**:307-319.
- Xiong Y, Mahmood A and Chopp M (2010) Angiogenesis, neurogenesis and brain recovery of function following injury. *Curr Opin Investig Drugs* **11**:298-308.

JPET #197483

Footnotes:

This work was supported by the Veterans affairs Merit Review [Grants BX000347, BX000891], – SCF); National Institutes of Health [Grants NS070239, NS063965], and Jordan University of Science and Technology predoctoral fellowship.

JPET #197483

Figure legends:

Figure 1: Hypertension and AT1 blockade affects the expression of BDNF. Blood pressure transmitters were implanted intraperitoneally in SHR. Animals had sham surgery and received a single dose of candesartan (0.3mg/kg) and the mean arterial blood pressure was monitored (A). Arrow indicates time of candesartan administration; n=6. SHR underwent sham surgery and randomized to receive either candesartan (0.3mg/kg) or saline intravenously (n=6 per group). 24 hours later the animals were sacrificed and the brains were extracted and. Right and left hemispheres were separated and processed for immunoblotting (B).

Figure 2: Angiotensin II and AT1 blockade affects the expression of BDNF and the angiogenic potential in hCMECs. hCMECs were cultured to confluence followed by serum starvation for 16 hours. Cells were treated with a concentration range of Angiotensin II (AngII) for different time periods. The expression of BDNF was assessed using immunoblotting (A) n=3-5. hCMECs were treated with AngII (1fM or 1 μ M) for 6 hours followed by treatment with candesartan (0.16 μ M) for 10 hours and the expression of BDNF was assessed (B) n=3. The ability of candesartan to modulate the angiogenic response of hCMECs was evaluated using in vitro matrigel tube formation assay (C) and wound recovery assay (D) in response to treatment with a concentration range of candesartan (0-1.6 μ M) n=3. For panels A and B, data presented as mean \pm SEM, * p<0.05. For panels C and D * significantly different from control; \$ significantly different from cand 0.16 μ M. Overall p=0.0014, F=8.19 (C) and p<0.0001; F=22.44 (D).

JPET #197483

Figure 3: Angiotensin II modulates the proliferation of hCMECs. Proliferative response of hCMECs was evaluated using BrdU incorporation assay. hCMECs were treated with AngII (1fM or 1 μ M) for 6 hours followed by candesartan (0.16 μ M) alone or in combination with other treatments. The plates were then processed according to the manufacturer recommendations. Data are presented as mean \pm SEM of 3 different experiments each in triplicate. * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall p=0.005, F=8.85 for AngII 10⁻⁹ and p<0.0001, F=34.97 for AngII 1 μ M

Figure 4: Angiotensin II modulates the migration of hCMECs. Insert; representative image of control, AngII 1 μ M and AngII+candesartan showing candesartan induced migration of hCMECs. hCMECs were cultured to confluence followed by 10 hours serum starvation. Cells were treated with either AngII 1fM or 1 μ M for 6 hours and then a scratch was introduced in the monolayer. Cells were then incubated with AngII 1fM or 1 μ M alone or with candesartan (A and D). The involvement of BDNF was assessed using a number of inhibitors for BDNF functions (B and E) which were added to the media 30 minutes before AngII treatment. Data are presented as mean \pm SEM of 3 different experiments each in triplicate. * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall p<0.0001, F=16.08 for AngII 1fM and p<0.0001, F=22.08 for AngII 1 μ M.

Figure 5: Angiotensin II modulates the angiogenic potential of hCMECs. Insert; representative images of in vitro tube formation (arrows) showing reduced tube

JPET #197483

formation rate in AngII 1 μ M treated hCMECs and the reversal of AngII 1 μ M antiangiogenic effect by candesartan.

hCMECS (2×10^4 cells/well) were suspended in a 30:60 solution of matrigel and serum free media. Angiogenic response of hCMECs to AngII 1 fM (A) and 1 μ M (D) in the presence and absence of candesartan was evaluated 24 hours after treatment. The involvement of BDNF was assessed through using K252a (Trk inhibitor) or TrkB-Fc (soluble chimeric receptor) (B and E). Data are presented as mean \pm SEM of 3 different experiments each in triplicate. * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall $p < 0.0001$, $F = 12.74$ for AngII 1 fM and $p < 0.0001$, $F = 9.54$ for AngII 1 μ M.

Figure 6: AT2 receptor mediates the angiogenic response in hCMECs. The involvement of AT2 receptor in the angiogenic response to AngII 1 fM alone or in combination with candesartan (A) and to the combination of AngII 1 μ M and candesartan (B) was assessed using the AT2 antagonist PD-123319 (0.1 μ M). The angiogenic response was evaluated using matrigel tube formation assay as described in the methods section. Data are presented as mean \pm SEM of 3 different experiments each in triplicate. * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall $p < 0.0001$, $F = 8.779$ for both AngII 1 fM and AngII 1 μ M.

Figure 7: Candesartan induced BDNF expression is mediated through AT2 receptor. To evaluate the involvement of AT2 receptor in candesartan induced BDNF

JPET #197483

expression hCMECs were pretreated with PD-123319 or vehicle followed by AngII (1fM or 1 μ M) in the presence or absence of candesartan (0.16 μ M) (Figure 7A). To further confirm the role of AT2 hCMECs were treated with CGP-42112A (0.1 μ M) or vehicle for 16 hours (Figure 7B). * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall $p=0.02$, $F=3.086$. $n=4-6$.

Figure 8: AT1 antagonism affects the expression of AT1 receptor in an AT2 receptor mediated manner. To assess the expression of AT1 (A and B) and AT2 (Cand D) receptors in response to the different treatments used. Cells were incubated with PD-123319 (0.1 μ M) or vehicle for 30 minutes followed by 6 hours of AngII 1fM (A and C) or 1 μ M (B and D). After 6 hours of AngII treatment cells were co-incubated with candesartan or vehicle for 10 hours. Receptor expression was assessed using immunoblotting. Data presented as mean \pm SEM; $n=3-5$. * Significantly different from control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall $p<0.0042$, $F=8.742$ for AngII 1 μ M.

Figure 9: AT1 antagonism modulates the phosphorylation of GSK-3 β in an AT2 receptor mediated manner. To assess the phosphorylation of GSK-3 β at the inhibitory serine 9 residue, hCMECs were treated as described for AngII receptors expression evaluation. Panel (A) represents the response to AngII 1fM while (B) represents AngII 1 μ M response. Data presented as mean \pm SEM; $n=3-5$. * Significantly different from

JPET #197483

control, # significantly different from AngII in the same group, \$ significantly different from AngII+cand in the same group; overall $p=0.0038$, $F=6.94$ for AngII 1 μ M.

Figure 10: A schematic representation of the results. By blocking AT1 receptors ARBs induce an unopposed stimulation of AT2 receptors. AT2 stimulation induces the expression of BDNF which will bind to its TrkB receptor to promote a proangiogenic state in hCMECs.

Figure 1





Figure 2










Figure 3





Downloaded from jpet.aspetjournals.org at ASPET Journals on December 27, 2024

Figure 4




downloaded from jpet.aspetjournals.org at ASPET Journals on December 27, 2004

Figure 5





Downloaded from jpet.aspetjournals.org at ASPET Journals on December 7, 2024

Figure 6



Downloaded from jpet.aspetjournals.org at ASPET Journals on December 27, 2024

Figure 7



Downloaded from jpet.aspetjournals.org at ASPET Journals on December 21, 2024

Figure 8







Figure 9






Figure 10




Supplementary Data:




S1: Angiotensin II increases the viability of human cerebrovascular endothelial cells (hCMECs). hCMECs were cultured and treated with different doses of AngII and their viability was assessed using MTT metabolism. AngII significantly increased the viability of hCMECs in all doses. * significantly different from ctrl; overall p=0.0001, F=18.36.

Ahmed Alhusban, Anna Kozak, Adviye Ergul, Susan C Fagan.
AT1 receptor antagonism is proangiogenic in the brain: BDNF a novel mediator
Journal of pharmacology and experimental therapeutics




S2: AT1 expression is induced through AT2 stimulation. hCMECs were serum starved for 16 hours followed by treatment with either the AT2 agonist CGP-42112A (0.1 μ M) or vehicle and the expression of AT1 was quantified. Data presented as mean \pm SEM, n=3.

Ahmed Alhusban, Anna Kozak, Adviye Ergul, Susan C Fagan.
AT1 receptor antagonism is proangiogenic in the brain: BDNF a novel mediator
Journal of pharmacology and experimental therapeutics



S3: AT1 blockade affects the expression of BDNF in wistar rats. Wistar rats were subjected to sham surgery and randomized to receive either saline or candesartan (1mg/kg) 3 hours after surgery. Candesartan significantly increased BDNF expression in both hemispheres as compared to the corresponding hemispheres in saline treated animals. p=0.04; n=4 per group.

Ahmed Alhusban, Anna Kozak, Adviye Ergul, Susan C Fagan.
AT1 receptor antagonism is proangiogenic in the brain: BDNF a novel mediator
Journal of pharmacology and experimental therapeutics



S4: Blood pressure reduction with hydralazine did not affect BDNF expression. Spontaneously hypertensive rats were exposed to sham surgery and randomized to receive either saline or hydralazine (1mg/kg) 3 hours after surgery. Hydralazine treatment resulted in a trend toward increased BDNF expression in the brain; but this trend did not reach statistical significance level. n=4 per group.