

JPET #200733

**A novel macrolide solithromycin exerts superior anti-inflammatory effect via
NF- κ B inhibition**

Yoshiki Kobayashi, Hiroo Wada, Christos Rossios, Dai Takagi, Manabu Higaki,
Shin'ichiro Mikura, Hajime Goto, Peter J. Barnes and Kazuhiro Ito

Airway Disease Section, National Heart and Lung Institute, Imperial College London,
London, United Kingdom

1: Airway Disease Section, National Heart and Lung Institute, Imperial College London,
London, United Kingdom: YK, CR, DT, PJB and KI

2: Department of Respiratory Medicine, Kyorin University, School of Medicine, Tokyo,
Japan: HW, MH, SM and HG

JPET #200733

A running title: Anti-inflammatory effect of solithromycin

Corresponding author: Kazuhiro Ito, Airway Disease Section, National Heart and Lung Institute, Imperial College London, Guy Scadding Building, Royal Brompton Campus, Dovehouse Street, London SW3 6LY, UK

Tel: +44-207-352-8121, Fax: +44-207-351-8126

E-mail: k.ito@imperial.ac.uk

The number of text pages:29

The number of tables: 1

The number of figures: 6

The number of references: 59

The number of words in the Abstract: 221

The number of words in the Introduction: 568

The number of words in the Discussion: 1122

Abbreviation

CAP; community-acquired pneumonia

COPD; chronic obstructive pulmonary disease

DPB; diffuse panbronchiolitis

LPS; lipopolysaccharide

MMP9; matrix metalloproteinase 9

NF- κ B; nuclear factor- κ B

PBMCs; peripheral blood mononuclear cells

JPET #200733

PMA; phorbol 12-myristate 13-acetate

TNF α ; tumor necrosis factor α

JPET #200733

Abstract

Macrolides are reported to reduce exacerbation of chronic inflammatory respiratory disease such as chronic obstructive pulmonary disease (COPD), and also show anti-inflammatory effects *in vitro* and *in vivo*. However the anti-inflammatory efficacies of current macrolides are relatively weak. Here we found that a novel macrolide/fluoroketolide solithromycin (CEM-101) showed superior anti-inflammatory effects to macrolides in current clinical use. The effects of solithromycin (SOL) on LPS-induced TNF α and/or CXCL8 release, PMA-induced MMP9 activity and NF- κ B activity under conditions of oxidative stress have been evaluated and compared with the effects of erythromycin, clarithromycin, azithromycin and telithromycin in human monocytic U937 cells and peripheral blood mononuclear cells (PBMC) obtained from COPD patients. We also examined effect of SOL on cigarette smoke-induced airway inflammation in mice. SOL exerted superior inhibitory effects on TNF α /CXCL8 production and MMP9 activity in U937 cells. In addition, SOL suppressed TNF α release and MMP9 activity in PBMC from COPD patients at 10 μ M, which is 100 times more potent than the other macrolides tested. Activated NF- κ B due to oxidative stress was completely reversed by SOL. SOL also inhibited cigarette smoke-induced neutrophilia and proMMP9 production *in vivo* although erythromycin did not inhibit them. Thus, SOL showed better anti-inflammatory profiles compared with macrolides currently used in the clinic, and may be a promising anti-inflammatory and anti-microbial macrolide for the treatment of COPD in the future.

Introduction

It has long been recognized that macrolides exert not only antibacterial activity, but also anti-inflammatory and immunoregulatory effects (Kano and Rubin, 2010). Erythromycin was found to induce a dramatic and unexpected improvement in diffuse panbronchiolitis (DPB) (Kudoh et al., 1998), leading to an investigation of its anti-inflammatory actions. Several studies indicate that macrolides inhibit pro-inflammatory cytokine production, such as CXCL8 and TNF α , as the mechanism of the effects of macrolides on DPB (Sakito et al., 1996). Macrolides also inhibit release of leukotriene B₄ (LTB₄), a neutrophil chemotactic mediator (Oda et al., 1995), and soluble adhesion molecules (Mukae et al., 1997), resulting in reduction of neutrophil chemotaxis and infiltration into the airway epithelium. It has also been reported that macrolides increase antioxidant activity in alveolar macrophages (Morikawa et al., 2000), enhance apoptosis of neutrophils (Inamura et al., 2000), decrease superoxide anion release by neutrophils (Tamaoki et al., 2004) and reduce mucus secretion in the airways (Kaneko et al., 2003).

Macrolides were found to be effective for DPB at doses below the concentrations required for their anti-bacterial activity, by inhibiting not only biofilm-formation and virulence factors produced from *Pseudomonas aeruginosa*, but also quorum-sensing systems in which bacteria coordinately regulate specific virulence genes (Schultz, 2004; Tateda et al., 2004). Thus, macrolides suppress airway inflammation observed in DPB, resulting in an improvement of lung function and prolonged survival (Oda et al., 1995; Kudoh et al., 1998; Kadota et al., 2003).

In addition to DPB, it has been reported that macrolides provide beneficial effects, although to a lesser degree, in other inflammatory airway diseases, such as

JPET #200733

bronchiectasis, cystic fibrosis, sarcoidosis, asthma and chronic obstructive pulmonary disease (COPD) (Trisolini et al., 2008; Kanoh and Rubin, 2010; Albert et al., 2011). Although DPB is geographically limited to East Asia, COPD is a serious worldwide health problem without any safe and effective anti-inflammatory therapy (Ito and Barnes, 2009). There are few reports investigating the clinical benefit of macrolides for COPD, with some studies showing reduced exacerbations after long-term therapy COPD (Simpson et al., 2008; He et al., 2010; Albert et al., 2011). However, the reduction in exacerbations could be due to the antibiotic action of macrolides rather than any anti-inflammatory effect and there is no evidence for any significant effects on inflammatory biomarkers in these studies.

The anti-inflammatory effects of macrolides have been previously investigated. For example, azithromycin inhibits pro-inflammatory cytokines from M1 macrophages in cystic fibrosis (Meyer et al., 2009). Clarithromycin and erythromycin inhibit airway inflammation in a cigarette smoke model (Nakanishi et al., 2009; Mikura et al., 2011). One of the molecular mechanisms of anti-inflammatory effects of macrolides is inhibition of pro-inflammatory transcription factors, such as nuclear factor- κ B (NF- κ B) (Li et al., 2012) and activated protein-1 (AP-1) (Bosnar et al., 2011).

Solithromycin (CEM-101)) has been developed as a novel oral and intravenous fluoroketolide antibacterial agent related to 14-member-ring macrolides that is currently in clinical development for severe community-acquired bacterial pneumonia (CABP). Solithromycin exhibits a greater potency and a wider spectrum of *in vitro* activity against respiratory tract pathogens associated with CABP, including macrolide-resistant strains, compared with other available macrolides (Farrell et al., 2010; McGhee et al., 2010). However, the anti-inflammatory efficacies of solithromycin

JPET #200733

have not yet been explored.

Here we compared the anti-inflammatory effects of solithromycin on release of pro-inflammatory mediators, such as TNF α , CXCL8 and MMP9, and activation of NF- κ B compared to other macrolides currently used clinically, such as erythromycin, clarythromycin, azithromycin and telithromycin

Methods

Reagents.

Solithromycin((3*aS*,4*R*,7*S*,9*R*,10*R*,11*R*,13*R*,15*R*,15*aR*)-1-[4-[4-(3-aminophenyl)-1*H*-1,2,3-triazol-1-yl]butyl]-4-ethyl-7-fluorooctahydro-11-methoxy-3*a*,7,9,11,13,15-hexamethyl-10-[[3,4,6-trideoxy-3-(dimethylamino)- β -D-xylo-hexopyranosyl]oxy]-2*H*-Oxacyclotetradecino[4,3-*d*]oxazole-2,6,8,14(1*H*,7*H*,9*H*)-tetrone) and telithromycin ((1*S*,2*R*,5*R*,7*R*,8*R*,9*S*,11*R*,13*R*,14*R*)-8-[(2*S*,3*R*,4*S*,6*R*)-4-dimethylamino-3-hydroxy-6-methyl-oxan-2-yl]oxy-2-ethyl-9-methoxy-1,5,7,9,11,13-hexamethyl-15-[4-(4-pyridin-3-ylimidazol-1-yl)butyl]-3,17-dioxa-15-azabicyclo[12.3.0]heptadecane-4,6,12,16-tetrone) were provided by Cempra Pharmaceuticals, Inc (Chapel Hill, NC). Erythromycin ((3*R*,4*S*,5*S*,6*R*,7*R*,9*R*,11*R*,12*R*,13*S*,14*R*)-6-[[[(2*S*,3*R*,4*S*,6*R*)-4-(dimethylamino)-3-hydroxy-6-methyloxan-2-yl]oxy]-14-ethyl-7,12,13-trihydroxy-4-[[[(2*R*,4*R*,5*S*,6*S*)-5-hydroxy-4-methoxy-4,6-dimethyloxan-2-yl]oxy]-3,5,7,9,11,13-hexamethyl-1-oxacyclotetradecane-2,10-dione), clarithromycin ((3*R*,4*S*,5*S*,6*R*,7*R*,9*R*,11*S*,12*R*,13*S*,14*S*)-6-[[[(2*S*,3*R*,4*S*,6*R*)-4-(dimethylamino)-3-hydroxy-6-methyloxan-2-yl]oxy]-14-ethyl-12,13-dihydroxy-4-[[[(2*R*,4*S*,5*S*,6*S*)-5-hydroxy-4-methoxy-4,6-dimethyloxan-2-yl]oxy]-7-methoxy-3,5,7,9,11,13-hexamethyl-1-oxacyclotetradecane-2,10-dione), azithromycin ((2*R*,3*S*,4*R*,5*R*,8*R*,10*R*,11*R*,12*S*,13*S*,14*R*)-2-ethyl-3,4,10-trihydroxy-3,5,6,8,10,12,14-heptomethyl-15-oxo-11-[[3,4,6-trideoxy-3-(dimethylamino)- β -D-xylo-]oxy]-1-oxa-6-azacyclopentadec-13-yl-2,6-dideoxy-3-*C*-methyl-3-*O*-methyl- α -L-*ribo*-hexopyranoside)), hydrogen peroxide, phorbol 12-myristate 13-acetate (PMA), and lipopolysaccharide (LPS) were purchased from Sigma Aldrich (Poole, UK).

Cells. The human monocytic cell line U937 was obtained from the American Type

JPET #200733

Culture Collection (ATCC, Rockville, MD). PBMCs from COPD patients were obtained from Brompton hospital and separated by AccuSPIN (Sigma–Aldrich). The characteristics of subjects are shown in Supplemental Table 1. Cells were cultured in complete growth medium (RPMI 1640) (Sigma–Aldrich) supplemented with 10% fetal bovine serum (FBS) and 1% L-glutamine at 37°C in a humidified atmosphere with 5% CO₂. U937 cells were differentiated into adherent macrophage-like morphology by exposure to PMA (50 ng/ml) for 48 hrs in complete growth medium. Cell viability was assessed microscopically by trypan blue staining. Cell toxicity was determined by MTT assay as needed. This study was approved by the ethics committee of the Royal Brompton and Harefield Hospitals, and all subjects gave written informed consent.

Cell Lysis. Whole cell extracts were prepared as previously described (Kobayashi et al., 2011). Briefly, cell protein extracts were prepared using modified RIPA buffer (50 mM Tris HCL pH 7.4, 0.5% NP-40, 0.25% Na-deoxycholate, 150 mM NaCl with freshly added complete protease inhibitor cocktail (Roche, Mannheim, Germany)). Protein concentration was determined using the BCA Protein Assay (Thermo Fisher Scientific, Waltham, MA).

Cytokine ELISA. TNF α and IL-8 concentrations in the supernatant of cell cultures were determined by sandwich ELISA according to the manufacturer's instructions (R&D Systems Europe, Abingdon, UK).

Zymography. MMP9 enzyme activity was measured by gelatin zymography. Cell culture supernatants were diluted with equal amount of Laemli sample buffer (Bio-Rad,

JPET #200733

Hertfordshire, UK) and loaded on a Novex® 10 % Zymogram (Gelatin) gel (Invitrogen Ltd, Paisley, UK). After electrophoresis, gels were incubated and rinsed with Novex® zymogram renaturing buffer (Invitrogen) for 30 min at room temperature. The gels were then rinsed in Novex® zymogram developing buffer (Invitrogen) for 30 min at room temperature prior to overnight incubation in the developing buffer at 37 °C. After incubation, the gels were stained using a Colloidal Blue Staining Kit (Invitrogen) to visualize the zymogen bands.

NF-κB activity. The activation of NF-κB (p65 binding activity to NF-κB binding sequence) was determined using a TransAM NF-κB p65 Assay kit (Active Motif, Inc., Carlsbad, CA) according to the manufacturer's instruction. As shown above, whole cell extracts were prepared from PMA-differentiated U937 cells, and 20 μl of each extract was used for this study. Results were determined by measuring the spectrophotometric absorbance at 450 nm with a reference wavelength of 655 nm.

Cigarette smoke exposure to mice. C57BL/6J mice (male, 4 weeks) were purchased from CLEA Japan, Inc, (Tokyo, Japan) and adapted for 1 week. Mice were exposed to cigarette smoke (4% cigarette smoke diluted with compressed air) for 30 min/day for 12 days or 4 days using the commercially marketed un-filtered Peace cigarettes (28 mg of tar and 2.3 mg of nicotine per cigarette; Japan Tobacco Inc., Tokyo, Japan) using a Tobacco Smoke Inhalation Experiment System for small animals (Model INH06-CIGR02A; MIPS, Osaka, Japan) as described previously (Nakamaru et al., 2009).

In cohort 1, solithromycin (100 mg/kg, orally) was suspended into 0.5%

JPET #200733

carboxymethylcysteine and administered orally for the last 8 days of 12 days tobacco smoke exposure (1hr before each tobacco smoke exposure). In cohort 2, solithromycin (100 mg/kg, orally) or erythromycin (100 mg/kg, orally) was suspended into 0.5% carboxymethylcysteine and administered orally at 1hr before each tobacco smoke exposure (1hr before each tobacco smoke exposure) for 4 times. Bronchoalveolar lavage was performed at 24hrs after the last cigarette smoke exposure as previously described (Nakamaru et al., 2009), and the number of alveolar macrophages and neutrophils were determined from Diff-Quick stained specimens. The proMMP-9 protein concentration in BAL fluid supernatant was assessed using a commercially available mouse proMMP-9 ELISA kit (R&D, Abingdon, UK) according to the manufacturer's instructions. This animal study was approved by The Experimental Animal Ethics Committee in Kyorin University.

Statistical analysis. The results were expressed as the mean \pm SEM. Comparisons of data in two groups were performed using the Student's *t* test or the Wilcoxon signed rank test. Multiple comparisons were made by one-way ANOVA with *post hoc* test (Dunnett's) as appropriate. The difference was considered significant at $p < 0.05$. IC₅₀ values (50% inhibitory concentration) for macrolides for production of cytokines or MMP9 were calculated using Prism 4.0 (GraphPad Software Inc., San Diego, CA).

Results

Anti-inflammatory effects of macrolides in U937 cells. LPS significantly increased TNF α and CXCL8 production in PMA-differentiated U937 cells (TNF α , 63.1 ± 2.6 fold in LPS vs. non-stimulated; and CXCL8, 2.0 ± 0.1 fold in LPS vs. non-stimulated cells, n=3). Solithromycin significantly inhibited both TNF α and CXCL8 at 100 μ M (Fig. 1A and Fig. 2A). Although clarithromycin showed modest effects on both TNF α and CXCL8 production at a higher concentration (333 μ M) (Fig. 1 C and Fig. 2C), erythromycin and azithromycin had no inhibitory effect (Fig. 1 B, D, Fig.2 B, D). Telithromycin (100 μ M) was similarly without effect on TNF α and CXCL8 (Fig. 1E, Fig. 2E). The IC₅₀ values for solithromycin on TNF α and CXCL8 release were 41.6 ± 1.9 μ M and 78.2 ± 9.5 μ M, respectively, and were superior to those for clarithromycin (IC₅₀, 426.3 ± 63.9 μ M for TNF α and 506.5 ± 44.0 μ M for CXCL8) (Table 1). None of compounds affected cell viability evaluated by MTT assay at maximal concentrations used in this test.

We also investigated the effects of macrolides on MMP9 activity, which was clearly elevated by PMA stimulation in U937 cells (9.9 ± 2.0 fold in PMA vs. non-stimulation, n=3). Solithromycin markedly reduced MMP9 activity, with an IC₅₀ of 14.9 ± 3.1 μ M (Fig. 3A and Table 1). In contrast, clarithromycin and azithromycin showed 10-fold lower inhibitory effects than solithromycin, whereas erythromycin showed no effect (Fig. 3B, C, D and Table 1). Telithromycin also inhibited MMP9 activity, although to lesser extent than solithromycin, with an IC₅₀ of 83.4 μ M (Fig. 3E).

Anti-inflammatory effects of macrolides in PBMC from COPD patients.

JPET #200733

Anti-inflammatory effects of solithromycin were evaluated in PBMC from COPD patients. Solithromycin (10 μ M) slightly but significantly inhibited LPS-induced TNF α release from PBMC of COPD patients, while erythromycin, clarithromycin and azithromycin failed to suppress TNF α release, even at a higher concentration (100 μ M) (Fig. 4 A-D). Furthermore, solithromycin clearly reduced MMP9 activity in PBMCs from COPD patients at 10 μ M. Although clarithromycin slightly but significantly inhibited MMP9 activity at a higher concentration (100 μ M), erythromycin and azithromycin showed no effect (Fig. 4 E-H). Telithromycin showed cytotoxicity at less than 100 μ M, and could not get meaningful data.

Solithromycin inhibited NF- κ B activity under conditions of oxidative stress. As the secretion of TNF α , CXCL8 and MMP9 are reported to be NF- κ B dependent, we investigated the effects of solithromycin on NF- κ B activity as defined by p65 binding activity to NF- κ B DNA binding sequence under conditions of oxidative stress. NF- κ B activity was already high in PMA-differentiated U937 cells and 4hrs exposure of H₂O₂ (200 μ M) significantly enhanced NF- κ B activity (114.1 \pm 1.1 % of non-treatment control, n=4). Clarithromycin and azithromycin restored H₂O₂-induced enhancement of NF- κ B activity at a high concentration (100 μ M) (Fig. 5C,D), whereas solithromycin provided remarkable inhibitory effects even at a lower concentration (10 μ M) (Fig. 5A). Erythromycin did not show any effects (Fig. 5B).

Inhibition of inflammatory cell accumulation in cigarette smoke-exposed mice *in vivo* by solithromycin. C57BL/6J mice were exposed to cigarette smoke (4%) for 12

JPET #200733

days (30 min/day) and treated with solithromycin at 100 mg/kg orally once daily 1hr before each smoke exposure during the last 8 days of the 12 day smoking course. At 24hrs after the last cigarette smoke exposure, bronchoalveolar lavage fluid was collected. Cigarette smoke exposure significantly increased the number of alveolar macrophages (air: 48828 ± 2847 cells/ml, smoke + vehicle: 240712 ± 22917) (Fig. 6A), neutrophils (air: 30.9 ± 20.3 cells/ml, smoke + vehicle: 5702.0 ± 725.4) (Fig. 6B) and lymphocytes (Air: 32.9 ± 21.7 cells/ml, smoke + vehicle: 1024.0 ± 138.1) in bronchoalveolar lavage fluid.

Solithromycin did not inhibit alveolar macrophage accumulation (smoke + SOL: 280660 ± 40121) (Fig. 6A), but significantly inhibited neutrophil accumulation by 49% (smoke + SOL: 2635.0 ± 829.3 , $p < 0.05$) (Fig. 6B) and lymphocyte accumulation by 75% (smoke + SOL: 276.5 ± 202.2 , $p < 0.05$). In addition, although cigarette smoke exposure decreased body weight, solithromycin significantly restored the body weight (Fig. 6C).

The effects of erythromycin were also compared to that of solithromycin. Head-to-head comparison was conducted in 4 days smoking model. C57BL/6J mice were exposed to cigarette smoke (4%) for 4 days (30 min/day) and treated with vehicle, solithromycin at 100 mg/kg or erythromycin at 100 mg/kg orally once daily 1hr before each smoke exposure. At 24hrs after the last cigarette smoke exposure, bronchoalveolar lavage fluid was collected. As shown in Fig. 6D, although solithromycin significantly inhibited airway neutrophilia, erythromycin did not inhibit it (air: 46.3 ± 25.7 cells/ml, smoke + vehicle: 13500 ± 1360 , smoke + SOL: 8320 ± 1160 , $p < 0.05$, smoke + EM: 15100 ± 1140 , *not significant*). In addition, proMMP9 (not active form) was also elevated in BAL by smoke exposure, and significantly inhibited by solithromycin, but not by

JPET #200733

erythromycin (air: 1.6 ± 0.21 ng/ml, smoke + vehicle: 3.9 ± 0.29 , smoke + SOL: 2.5 ± 0.17 , $p < 0.05$, smoke + EM: 3.9 ± 0.70 , *not significant*)

Discussion

In this study, we demonstrated that a novel macrolide/fluoroketolide, solithromycin has beneficial anti-inflammatory effects in addition to the anti-microbial ability previously shown. The inhibitory activities on production of TNF α , CXCL8 and MMP9, as well as NF- κ B activation, were superior to those of other macrolide antibiotics currently used in the clinic. Even more importantly, solithromycin also inhibited pro-inflammatory mediators released by PBMC obtained from COPD patients and inhibited neutrophilia and proMMP9 production in cigarette smoke-exposed mice *in vivo*, which are known to be corticosteroid resistant.

TNF α and CXCL8 were increased in bronchoalveolar lavage, sputum and serum in COPD (Stankiewicz et al., 2002; Vermooy et al., 2002) and are believed to play an important role in the pathogenesis of COPD (Barnes, 2009). There are many *in vitro* studies indicating that macrolides inhibit pro-inflammatory mediators, such as CXCL8 and TNF α released from monocytes (Iino et al., 1992; Kikuchi et al., 2002), neutrophils (Tsuchihashi et al., 2002), bronchial epithelial cells (Takizawa et al., 1997) and induced sputum cells (Marjanovic et al., 2011). Macrolides also exerted inhibitory effects on neutrophil-dominant inflammation in *in vivo* models (Leiva et al., 2008; Bosnar et al., 2009). Furthermore, macrolides inhibited release of inflammatory mediators including TNF α and CXCL8 from sputum cells obtained from patients with COPD (Marjanovic et al., 2011). In addition to these *in vitro* results, clinical benefits by long-term macrolide therapy have been reported in COPD patients. Although short-term treatment did not provide any advantages for patients with stable COPD (Banerjee et al., 2005), long-term (more than 6 months) macrolide therapy reduces exacerbation of COPD (Seemungal et al., 2008; Blasi et al., 2010; He et al., 2010).

JPET #200733

Macrolides have also been reported to exert beneficial effects in asthma. Low-dose erythromycin therapy in children with asthma for 6 months improved their symptoms and serum CXCL8 levels (Korematsu et al., 2010). Telithromycin showed beneficial effects on acute exacerbations of adult asthma (Johnston et al., 2006). Even short-term (8 week) clarithromycin treatment reduced sputum CXCL8, neutrophils, neutrophil elastase and MMP9 and also improved QOL scores in patients with refractory neutrophilic asthma (Simpson et al., 2008). Some *in vitro* studies have supported these clinical findings. Erythromycin and clarithromycin reduced CXCL8 release from human eosinophils in patients with allergic diseases, including asthma (Kohyama et al., 1999). Moreover, azithromycin inhibited IL-5 release from Th0 and Th2 cells in atopic asthmatic children (Lin et al., 2011). Thus, macrolides have shown great promise in their ability to reduce airway inflammation, including improvements in quality of life and exacerbation rates in airways diseases such as COPD and asthma (Simpson et al., 2009), but the efficacies are relatively moderate.

Excessive or inappropriate expression of MMP is closely associated with pathogenesis of tissue destructive processes in a wide variety of diseases including lung diseases such as COPD (Ohbayashi, 2002). In peripheral lung from COPD patients, MMP9 expression and activity are up-regulated and related to disease severity (Nakamaru et al., 2009). Furthermore, overexpression of human MMP9 in macrophages induced adult onset emphysema in mice and was associated with the loss of alveolar elastin (Foronjy et al., 2008). MMP9 produced from macrophages might be important in the development of emphysema (Russell et al., 2002). Ribeiro et al. demonstrated that azithromycin prevented the up-regulation of MMP9 genes, which were triggered by inflammatory stimuli (Ribeiro et al., 2009). In addition, there were a few *in vivo* studies

JPET #200733

evaluating effect of macrolide on MMP9 activity (Simpson et al., 2008; Ogawa et al., 2009). Simpson et al. showed that MMP9 concentrations were reduced in sputum from refractory asthmatics receiving clarithromycin. As shown in Figure 3 and 4, the effect of solithromycin on MMP9 production was more potent than that of clarithromycin, promising higher efficacy of solithromycin. The inhibition of MMP9 by solithromycin was also confirmed in smoke exposed mice (Fig. 6E)

The inflammation in COPD is amplified by increased oxidative stress (Rahman and Adcock, 2006). Oxidative stress activates NF- κ B, a transcription factor, that orchestrates the expression of multiple inflammatory genes, including TNF α , CXCL8 and MMP9 (Barnes, 2009). In fact, NF- κ B is reported to be regulated by macrolides. For example, azithromycin inhibited pro-inflammatory mediators, such as CXCL8, via inhibition of NF- κ B *in vitro* models (Aghai et al., 2007; Matsumura et al., 2011), although it inhibited LPS-induced pulmonary neutrophilia and IL-1 β with concomitant suppression of AP-1, but not NF- κ B *in vivo* (Bosnar et al., 2011). In addition, erythromycin and clarithromycin also inhibited CXCL8 with inactivation of NF- κ B and/or AP-1 in human bronchial epithelial cells *in vitro* (Abe et al., 2000; Desaki et al., 2000) as the effects were moderate. In this study, we also confirmed that solithromycin inhibited activated NF- κ B under oxidative stress and the efficacy was superior to those of other macrolides currently used clinically. As the mechanism of NF- κ B inactivation by macrolides, it is reported that macrolides are able to suppress the degradation of I κ B α , an inhibitor of NF- κ B, and/or to affect downstream of the dissociation from I κ B α in the NF- κ B signalling pathway (Desaki et al., 2004; Aghai et al., 2007).

JPET #200733

There have been some negative findings on the anti-inflammatory effects of macrolides. For example, neither erythromycin nor azithromycin directly inhibited inflammatory mediators such as CXCL8 and MMPs in human gingival cells (Kamemoto et al., 2009). Neither clarithromycin nor azithromycin had beneficial effects on CXCL8 release in human whole blood cells and alveolar macrophages (Kurdowska et al., 2001). These inconsistent results on anti-inflammatory effects could be due to time-dependent responses to macrolides via cross talk of signaling pathway (Shinkai et al., 2006) , due to weak/moderate efficacies, or variable cellular intake, the results became variable.

Telithromycin, a commercially available ketolide, has also been reported to exert anti-inflammatory effects through NF- κ B inhibition (Leiva et al., 2008). We therefore compared the effect of telithromycin with that of solithromycin. We found that telithromycin inhibited MMP9 activity (IC_{50} ; 83.4 μ M), although to a lesser extent than solithromycin (IC_{50} ; 14.9 μ M). However, it did not inhibit TNF α or CXCL8 release in macrophage system. In PBMC, as telithromycin showed cytotoxicity (significant at 100 μ M), we could not evaluate the efficacies properly. In addition, the clinical use of telithromycin is limited due to its side effects that include liver failure, exacerbation of myasthenia gravis, and visual disturbances. The pyridine-imidazole-containing moiety, which is a part of telithromycin, could induce inhibition of the nicotinic acetyl-choline receptors at the vagus nerve innervating the liver, the neuromuscular junction receptors, and the ciliary ganglion, resulting in these side effects, whereas solithromycin does not contain this moiety (Bertrand et al., 2010).

In conclusion, our findings revealed that a novel macrolide/ketolide, solithromycin exerts potent anti-inflammatory effects compared with other macrolides

JPET #200733

used clinically, possibly through NF- κ B inhibition. This suggests that with its superior anti-microbial and anti-inflammatory activities, solithromycin might be a potential therapy for refractory chronic inflammatory respiratory diseases such as COPD.

JPET #200733

Acknowledgement

We would like to thank Dr. Prabhavathi Fernandes (Cempra Phamaceuticals) for supply of CEM-101 and relevant information for the compound. We would like to thank Mr. Arnell Colongon and Ms. Sally Meah for their support in the recruitment of patients.

Authorship Contributions

Participated in research design: Kobayashi, Goto, Wada, Barnes, and Ito.

Conducted experiments: Kobayashi, Wada, Rossios, Takagi, Higaki, and Mikura.

Performed data analysis: Kobayashi, Wada and Ito.

Wrote or contributed to the writing of the manuscript: Kobayashi, Barnes and Ito.

References

- Abe S, Nakamura H, Inoue S, Takeda H, Saito H, Kato S, Mukaida N, Matsushima K and Tomoiike H (2000) Interleukin-8 gene repression by clarithromycin is mediated by the activator protein-1 binding site in human bronchial epithelial cells. *Am J Respir Cell Mol Biol* **22**:51-60.
- Aghai ZH, Kode A, Saslow JG, Nakhla T, Farhath S, Stahl GE, Eydelman R, Strande L, Leone P and Rahman I (2007) Azithromycin suppresses activation of nuclear factor-kappa B and synthesis of pro-inflammatory cytokines in tracheal aspirate cells from premature infants. *Pediatr Res* **62**:483-488.
- Albert RK, Connett J, Bailey WC, Casaburi R, Cooper JA, Jr., Criner GJ, Curtis JL, Dransfield MT, Han MK, Lazarus SC, Make B, Marchetti N, Martinez FJ, Madinger NE, McEvoy C, Niewoehner DE, Porsasz J, Price CS, Reilly J, Scanlon PD, Sciurba FC, Scharf SM, Washko GR, Woodruff PG and Anthonisen NR (2011) Azithromycin for prevention of exacerbations of COPD. *N Engl J Med* **365**:689-698.
- Banerjee D, Khair OA and Honeybourne D (2005) The effect of oral clarithromycin on health status and sputum bacteriology in stable COPD. *Respir Med* **99**:208-215.
- Barnes PJ (2009) The cytokine network in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* **41**:631-638.
- Bertrand D, Bertrand S, Neveu E and Fernandes P (2010) Molecular characterization of off-target activities of telithromycin: a potential

JPET #200733

role for nicotinic acetylcholine receptors. *Antimicrob Agents Chemother* **54**:5399-5402.

Blasi F, Bonardi D, Aliberti S, Tarsia P, Confalonieri M, Amir O, Carone M, Di Marco F, Centanni S and Guffanti E (2010) Long-term azithromycin use in patients with chronic obstructive pulmonary disease and tracheostomy. *Pulm Pharmacol Ther* **23**:200-207.

Bosnar M, Bosnjak B, Cuzic S, Hrvacic B, Marjanovic N, Glojnaric I, Culic O, Parnham MJ and Erakovic Haber V (2009) Azithromycin and clarithromycin inhibit lipopolysaccharide-induced murine pulmonary neutrophilia mainly through effects on macrophage-derived granulocyte-macrophage colony-stimulating factor and interleukin-1beta. *J Pharmacol Exp Ther* **331**:104-113.

Bosnar M, Cuzic S, Bosnjak B, Nujic K, Ergovic G, Marjanovic N, Pasalic I, Hrvacic B, Polancec D, Glojnaric I and Erakovic Haber V (2011) Azithromycin inhibits macrophage interleukin-1beta production through inhibition of activator protein-1 in lipopolysaccharide-induced murine pulmonary neutrophilia. *Int Immunopharmacol* **11**:424-434.

Desaki M, Okazaki H, Sunazuka T, Omura S, Yamamoto K and Takizawa H (2004) Molecular mechanisms of anti-inflammatory action of erythromycin in human bronchial epithelial cells: possible role in the signaling pathway that regulates nuclear factor-kappaB activation. *Antimicrob Agents Chemother* **48**:1581-1585.

Desaki M, Takizawa H, Ohtoshi T, Kasama T, Kobayashi K, Sunazuka T,

JPET #200733

- Omura S, Yamamoto K and Ito K (2000) Erythromycin suppresses nuclear factor-kappaB and activator protein-1 activation in human bronchial epithelial cells. *Biochem Biophys Res Commun* **267**:124-128.
- Farrell DJ, Sader HS, Castanheira M, Biedenbach DJ, Rhomberg PR and Jones RN (2010) Antimicrobial characterisation of CEM-101 activity against respiratory tract pathogens, including multidrug-resistant pneumococcal serogroup 19A isolates. *Int J Antimicrob Agents* **35**:537-543.
- Foronjy R, Nkyimbeng T, Wallace A, Thankachen J, Okada Y, Lemaitre V and D'Armiento J (2008) Transgenic expression of matrix metalloproteinase-9 causes adult-onset emphysema in mice associated with the loss of alveolar elastin. *Am J Physiol Lung Cell Mol Physiol* **294**:L1149-1157.
- He ZY, Ou LM, Zhang JQ, Bai J, Liu GN, Li MH, Deng JM, MacNee W and Zhong XN (2010) Effect of 6 months of erythromycin treatment on inflammatory cells in induced sputum and exacerbations in chronic obstructive pulmonary disease. *Respiration* **80**:445-452.
- Iino Y, Toriyama M, Kudo K, Natori Y and Yuo A (1992) Erythromycin inhibition of lipopolysaccharide-stimulated tumor necrosis factor alpha production by human monocytes in vitro. *Ann Otol Rhinol Laryngol Suppl* **157**:16-20.
- Inamura K, Ohta N, Fukase S, Kasajima N and Aoyagi M (2000) The effects of erythromycin on human peripheral neutrophil apoptosis. *Rhinology*

38:124-129.

Ito K and Barnes PJ (2009) COPD as a disease of accelerated lung aging.

Chest 135:173-180.

Johnston SL, Blasi F, Black PN, Martin RJ, Farrell DJ and Nieman RB

(2006) The effect of telithromycin in acute exacerbations of asthma. *N*

Engl J Med 354:1589-1600.

Kadota J, Mukae H, Ishii H, Nagata T, Kaida H, Tomono K and Kohno S

(2003) Long-term efficacy and safety of clarithromycin treatment in patients with diffuse panbronchiolitis. *Respir Med* 97:844-850.

Kamemoto A, Ara T, Hattori T, Fujinami Y, Imamura Y and Wang PL (2009)

Macrolide antibiotics like azithromycin increase lipopolysaccharide-induced IL-8 production by human gingival fibroblasts. *Eur J Med Res* 14:309-314.

Kaneko Y, Yanagihara K, Seki M, Kuroki M, Miyazaki Y, Hirakata Y, Mukae

H, Tomono K, Kadota J and Kohno S (2003) Clarithromycin inhibits overproduction of muc5ac core protein in murine model of diffuse panbronchiolitis. *Am J Physiol Lung Cell Mol Physiol* 285:L847-853.

Kanoh S and Rubin BK (2010) Mechanisms of action and clinical application

of macrolides as immunomodulatory medications. *Clin Microbiol Rev* 23:590-615.

Kikuchi T, Hagiwara K, Honda Y, Gomi K, Kobayashi T, Takahashi H, Tokue

Y, Watanabe A and Nukiwa T (2002) Clarithromycin suppresses lipopolysaccharide-induced interleukin-8 production by human

monocytes through AP-1 and NF-kappa B transcription factors. *J Antimicrob Chemother* **49**:745-755.

Kobayashi Y, Mercado N, Barnes PJ and Ito K (2011) Defects of protein phosphatase 2A causes corticosteroid insensitivity in severe asthma. *PLoS One* **6**:e27627.

Kohyama T, Takizawa H, Kawasaki S, Akiyama N, Sato M and Ito K (1999) Fourteen-member macrolides inhibit interleukin-8 release by human eosinophils from atopic donors. *Antimicrob Agents Chemother* **43**:907-911.

Korematsu S, Yamamoto K, Nagakura T, Miyahara H, Okazaki N, Akiyoshi K, Maeda T, Suenobu S and Izumi T (2010) The indication and effectiveness of low-dose erythromycin therapy in pediatric patients with bronchial asthma. *Pediatr Allergy Immunol* **21**:489-492.

Kudoh S, Azuma A, Yamamoto M, Izumi T and Ando M (1998) Improvement of survival in patients with diffuse panbronchiolitis treated with low-dose erythromycin. *Am J Respir Crit Care Med* **157**:1829-1832.

Kurdowska A, Noble JM and Griffith DE (2001) The effect of azithromycin and clarithromycin on ex vivo interleukin-8 (IL-8) release from whole blood and IL-8 production by human alveolar macrophages. *J Antimicrob Chemother* **47**:867-870.

Leiva M, Ruiz-Bravo A and Jimenez-Valera M (2008) Effects of telithromycin in in vitro and in vivo models of lipopolysaccharide-induced airway inflammation. *Chest* **134**:20-29.

JPET #200733

- Li M, Zhong X, He Z, Wen M, Li J, Peng X, Liu G, Deng J, Zhang J and Bai J (2012) Effect of erythromycin on cigarette-induced histone deacetylase protein expression and nuclear factor-kappaB activity in human macrophages in vitro. *Int Immunopharmacol*.
- Lin SJ, Lee WJ, Liang YW, Yan DC, Cheng PJ and Kuo ML (2011) Azithromycin Inhibits IL-5 Production of T Helper Type 2 Cells from Asthmatic Children. *Int Arch Allergy Immunol* 156:179-186.
- Marjanovic N, Bosnar M, Michielin F, Wille DR, Anic-Milic T, Culic O, Popovic-Grle S, Bogdan M, Parnham MJ and Erakovic Haber V (2011) Macrolide antibiotics broadly and distinctively inhibit cytokine and chemokine production by COPD sputum cells in vitro. *Pharmacol Res* 63:389-397.
- Matsumura Y, Mitani A, Suga T, Kamiya Y, Kikuchi T, Tanaka S, Aino M and Noguchi T (2011) Azithromycin May Inhibit Interleukin-8 Through the Suppression of Rac1 and a Nuclear Factor KappaB Pathway in KB Cells Stimulated With Lipopolysaccharide. *J Periodontol*.
- McGhee P, Clark C, Kosowska-Shick KM, Nagai K, Dewasse B, Beachel L and Appelbaum PC (2010) In vitro activity of CEM-101 against *Streptococcus pneumoniae* and *Streptococcus pyogenes* with defined macrolide resistance mechanisms. *Antimicrob Agents Chemother* 54:230-238.
- Meyer M, Huaux F, Gavilanes X, van den Brule S, Lebecque P, Lo Re S, Lison D, Scholte B, Wallemacq P and Leal T (2009) Azithromycin reduces

JPET #200733

exaggerated cytokine production by M1 alveolar macrophages in cystic fibrosis. *Am J Respir Cell Mol Biol* **41**:590-602.

Mikura S, Wada H, Higaki M, Yasutake T, Ishii H, Kamiya S and Goto H (2011) Erythromycin prevents the pulmonary inflammation induced by exposure to cigarette smoke. *Transl Res* **158**:30-37.

Morikawa T, Kadota JI, Kohno S and Kondo T (2000) Superoxide dismutase in alveolar macrophages from patients with diffuse panbronchiolitis. *Respiration* **67**:546-551.

Mukae H, Kadota J, Ashitani J, Taniguchi H, Mashimoto H, Kohno S and Matsukura S (1997) Elevated levels of soluble adhesion molecules in serum of patients with diffuse panbronchiolitis. *Chest* **112**:1615-1621.

Nakamaru Y, Vuppusetty C, Wada H, Milne JC, Ito M, Rossios C, Elliot M, Hogg J, Kharitonov S, Goto H, Bemis JE, Elliott P, Barnes PJ and Ito K (2009) A protein deacetylase SIRT1 is a negative regulator of metalloproteinase-9. *FASEB J* **23**:2810-2819.

Nakanishi Y, Kobayashi D, Asano Y, Sakurai T, Kashimura M, Okuyama S, Yoneda Y, Shapiro SD and Takayama K (2009) Clarithromycin prevents smoke-induced emphysema in mice. *Am J Respir Crit Care Med* **179**:271-278.

Oda H, Kadota J, Kohno S and Hara K (1995) Leukotriene B4 in bronchoalveolar lavage fluid of patients with diffuse panbronchiolitis. *Chest* **108**:116-122.

Ogawa M, Suzuki J, Takayama K and Isobe M (2009) Matrix

JPET #200733

metalloproteinase suppression induced by clarithromycin in murine cardiac allografts. *Transplant Proc* 41:395-397.

Ohbayashi H (2002) Matrix metalloproteinases in lung diseases. *Curr Protein Pept Sci* 3:409-421.

Rahman I and Adcock IM (2006) Oxidative stress and redox regulation of lung inflammation in COPD. *Eur Respir J* 28:219-242.

Ribeiro CM, Hurd H, Wu Y, Martino ME, Jones L, Brighton B, Boucher RC and O'Neal WK (2009) Azithromycin treatment alters gene expression in inflammatory, lipid metabolism, and cell cycle pathways in well-differentiated human airway epithelia. *PLoS One* 4:e5806.

Russell RE, Culpitt SV, DeMatos C, Donnelly L, Smith M, Wiggins J and Barnes PJ (2002) Release and activity of matrix metalloproteinase-9 and tissue inhibitor of metalloproteinase-1 by alveolar macrophages from patients with chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 26:602-609.

Sakito O, Kadota J, Kohno S, Abe K, Shirai R and Hara K (1996) Interleukin 1 beta, tumor necrosis factor alpha, and interleukin 8 in bronchoalveolar lavage fluid of patients with diffuse panbronchiolitis: a potential mechanism of macrolide therapy. *Respiration* 63:42-48.

Schultz MJ (2004) Macrolide activities beyond their antimicrobial effects: macrolides in diffuse panbronchiolitis and cystic fibrosis. *J Antimicrob Chemother* 54:21-28.

Seemungal TA, Wilkinson TM, Hurst JR, Perera WR, Sapsford RJ and

JPET #200733

Wedzicha JA (2008) Long-term erythromycin therapy is associated with decreased chronic obstructive pulmonary disease exacerbations. *Am J Respir Crit Care Med* **178**:1139-1147.

Shinkai M, Foster GH and Rubin BK (2006) Macrolide antibiotics modulate ERK phosphorylation and IL-8 and GM-CSF production by human bronchial epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **290**:L75-85.

Simpson JL, Phipps S and Gibson PG (2009) Inflammatory mechanisms and treatment of obstructive airway diseases with neutrophilic bronchitis. *Pharmacol Ther* **124**:86-95.

Simpson JL, Powell H, Boyle MJ, Scott RJ and Gibson PG (2008) Clarithromycin targets neutrophilic airway inflammation in refractory asthma. *Am J Respir Crit Care Med* **177**:148-155.

Stankiewicz W, Dabrowski MP, Chcialowski A and Plusa T (2002) Cellular and cytokine immunoregulation in patients with chronic obstructive pulmonary disease and bronchial asthma. *Mediators Inflamm* **11**:307-312.

Takizawa H, Desaki M, Ohtoshi T, Kawasaki S, Kohyama T, Sato M, Tanaka M, Kasama T, Kobayashi K, Nakajima J and Ito K (1997) Erythromycin modulates IL-8 expression in normal and inflamed human bronchial epithelial cells. *Am J Respir Crit Care Med* **156**:266-271.

Tamaoki J, Kadota J and Takizawa H (2004) Clinical implications of the

JPET #200733

immunomodulatory effects of macrolides. *Am J Med* **117 Suppl 9A**:5S-11S.

Tateda K, Standiford TJ, Pechere JC and Yamaguchi K (2004) Regulatory effects of macrolides on bacterial virulence: potential role as quorum-sensing inhibitors. *Curr Pharm Des* **10**:3055-3065.

Trisolini R, Cancellieri A, Paioli D, Burzi M, Orlandi P and Patelli M (2008) Sarcoidosis in the setting of idiopathic chronic bronchiolitis with airway colonization from *P. aeruginosa*: treatment with low-dose macrolides. *Intern Med* **47**:537-542.

Tsuchihashi Y, Oishi K, Yoshimine H, Suzuki S, Kumatori A, Sunazuka T, Omura S, Matsushima K and Nagatake T (2002) Fourteen-member macrolides suppress interleukin-8 production but do not promote apoptosis of activated neutrophils. *Antimicrob Agents Chemother* **46**:1101-1104.

Vernooy JH, Kucukaycan M, Jacobs JA, Chavannes NH, Buurman WA, Dentener MA and Wouters EF (2002) Local and systemic inflammation in patients with chronic obstructive pulmonary disease: soluble tumor necrosis factor receptors are increased in sputum. *Am J Respir Crit Care Med* **166**:1218-1224.

JPET #200733

Footnotes

This work was supported by a project grant from Cembra Pharmaceuticals, Inc. KI is currently employed by RespiVert Ltd and has honorary contract with Imperial College. PJB has served on Scientific Advisory Boards of AstraZeneca, Boehringer-Ingelheim, Chiesi, Daiichi-Sankyo, GlaxoSmithKline, Novartis, Nycomed, Pfizer, Teva and UCB and has received research funding from Aquinox Pharmaceutiocal, AstraZeneca, Boehringer-Ingelheim, Chiesi, Daiichi-Sankyo, GlaxoSmithKline, Novartis, Nycomed, Pfizer, RespiVert and Prosonix. The other authors have declared that no competing interests exist for this research.

Legend for figures

Fig. 1. Effects of macrolides on LPS-induced TNF α release in PMA-differentiated U937 cells. Cells were pretreated with solithromycin (A) (10 to 100 μ M) or erythromycin (B), clarithromycin (C), azithromycin (D) (33 to 333 μ M), telithromycin (E) (10 to 100 μ M) for 1 hr, followed by LPS (100 ng/ml) stimulation for 4 hrs. LPS-induced TNF α release was evaluated by ELISA. Values represent means of three experiments \pm SEM. ^{##} $p < 0.01$ (vs. non-treatment control), * $p < 0.05$, ** $p < 0.01$ (vs. treatment with LPS only).

Fig. 2. Effects of macrolides on LPS-induced CXCL8 release in PMA-differentiated U937 cells. Cells were pretreated with solithromycin (A) (10 to 100 μ M) or erythromycin (B), clarithromycin (C), azithromycin (D) (33 to 333 μ M), telithromycin (E) (10 to 100 μ M) for 1 hr, followed by LPS (100 ng/ml) stimulation for 4 hrs. LPS-induced CXCL8 release was evaluated by ELISA. Values represent means of three experiments \pm SEM. ^{##} $p < 0.01$ (vs. non-treatment control), * $p < 0.05$, ** $p < 0.01$ (vs. treatment with LPS only).

Fig. 3. Effects of macrolides on PMA-induced MMP9 activation in U937 cells. Cells were pretreated with solithromycin (A) (10 to 100 μ M) or erythromycin (B), clarithromycin (C), or azithromycin (D) (33 to 333 μ M), or telithromycin (E) (10 to 100 μ M) for 1 hr, followed by PMA (50 ng/ml) treatment for 48 hrs. After 48 hrs supernatants were collected for zymography. MMP9 enzyme activity was measured by gelatin zymography. Data are expressed relative to standard. Values represent means of

JPET #200733

four (A) or three (B, C and D) experiments \pm SEM. ^{###} $p < 0.01$ (vs. non-treatment control), * $p < 0.05$, ** $p < 0.01$ (vs. treatment with PMA only).

Fig. 4. Effects of macrolides on TNF α release and MMP9 production by PBMC from COPD patients. PBMC were incubated with macrolide compounds (solithromycin (A, E), 10 μ M; erythromycin (B, F), clarithromycin (C, G) and azithromycin (D, H), 100 μ M) overnight. TNF α release was evaluated by ELISA. MMP9 enzyme activity in supernatants was measured by gelatin zymography. Seven or eight individual values are shown. ^{###} $p < 0.01$, [#] $p < 0.05$ (vs. non-treatment control).

Fig. 5. Effects of macrolides on NF- κ B activity under conditions of oxidative stress. PMA-differentiated U937 cells were pretreated with solithromycin (A) (10 and 33 μ M) or erythromycin (B), clarithromycin (C), or azithromycin (D) (33 and 100 μ M) for 1 hr, followed by H₂O₂ (200 μ M) stimulation for 4 hrs. Binding activity of p65 to the NF- κ B binding sequence was measured by spectrophotometer. Data were expressed relative to standard (1 mg of recombinant p65). Values represent means of three experiments \pm SEM. ^{###} $p < 0.01$ (vs. non-treatment control), * $p < 0.05$, ** $p < 0.01$ (vs. treatment with H₂O₂ only).

Fig. 6. Anti-inflammatory effects of solithromycin in cigarette smoke-exposed mice. (A-C) C57BL/6J mice were exposed to cigarette smoke (4%) for 30min/day for 12 days. Solithromycin (100mg/kg orally: SOL) or vehicle was administered once daily 1hr before cigarette smoke exposure for the last 8 days of smoke exposure. The number of

JPET #200733

alveolar macrophages (A) and neutrophils (B) in BAL fluid were calculated. (C) The effects of solithromycin on an increase in body weight over 12 days. . (D,E) C57BL/6J mice were exposed to cigarette smoke (4%) for 30min/day for 4 days. Solithromycin or erythromycin (100mg/kg orally: SOL/EM) or vehicle was administered once daily 1hr before cigarette smoke exposure. Neutrophils (D), proMMP9 in BAL (E). $p < 0.05$ vs. smoke control (SM).NS: not significant

JPET #200733


Table 1. Effect of macrolides on inhibition of LPS-induced CXCL8 and TNF α release, and PMA-induced MMP9 activation in U937 cells

	IC ₅₀ (μ M)				
	Solithro- mycin	Erythro- mycin	Clarithro- mycin	Azithro- mycin	Telithro- mycin
LPS- induce CXCL8 release	78.2	NE at 333 μ M	506.5	NE at 333 μ M	NE at 100 μ M
LPS- induced TNF α release	41.6	NE at 333 μ M	426.3	NE at 333 μ M	NE at 100 μ M
PMA- induced MMP9 activation	14.9	NE at 333 μ M	118.0	212.1	83.4


NE: no effect

Figure 1


A




B



C



D



E




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



**A novel macrolide solithromycin exerts superior anti-inflammatory effect via
NF- κ B inhibition**

Yoshiki Kobayashi, Hiroo Wada, Christos Rossios, Dai Takagi, Manabu Higaki,
Shin'ichiro Mikura, Hajime Goto, Peter J. Barnes and Kazuhiro Ito

Airway Disease Section, National Heart and Lung Institute, Imperial College London,
London, United Kingdom

1: Airway Disease Section, National Heart and Lung Institute, Imperial College London,
London, United Kingdom: YK, CR, DT, PJB and KI

2: Department of Respiratory Medicine, Kyorin University, School of Medicine, Tokyo,
Japan: HW, MH, SM and HG

Supplemental Table 1.

Study subjects for PBMCs

	COPD (n=8)
Age (yr)	67.3 ± 1.9
Gender (M / F)	4 / 4
FEV ₁	1.36 ± 0.19
FEV ₁ (%pred)	56.6 ± 7.5
FEV ₁ /FVC	50.9 ± 5.5