

JPET #241901

The cannabinoid CB₂ agonist GW405833 suppresses inflammatory and neuropathic pain through a CB₁ mechanism that is independent of CB₂ receptors in mice

Ai-Ling Li, Lawrence M. Carey, Ken Mackie and Andrea G. Hohmann

Department of Psychological and Brain Sciences, Indiana University, Bloomington, IN, United States (A-L L, L.M.C., K.M., A.G.H.)

Program in Neuroscience, Indiana University, Bloomington, IN, United States (L.M.C., K.M., A.G.H.)

Gill Center for Biomolecular Science, Indiana University, Bloomington, IN, United States (K.M., A.G.H.)

JPET #241901

Running title: Anti-allyodynamic effect of GW405833 is CB₁ dependent

Corresponding author: Andrea G. Hohmann, Department of Psychological and Brain Sciences, Indiana University, 1101 E 10th Street, Bloomington, IN 47405-7007, USA.

Phone: +1 8128560672, email: hohmanna@indiana.edu

Number of text pages:	23
Number of tables:	1
Number of figures:	8
Number of references:	35
Number of words in Abstract:	246
Number of words in Introduction:	750
Number of words in Discussion:	1468

Abbreviations:

CB₁ or CB₂, cannabinoid receptor 1 or 2; CFA, complete Freund's adjuvant; CNS, central nervous system; HEK, human embryonic kidney; KO, knock out; PSNL, partial sciatic nerve ligation; Δ^9 -THC, Δ^9 -tetrahydrocannabinol; WT, wild type.

Recommended section: Behavioral Pharmacology

ABSTRACT

GW405833, widely accepted as a cannabinoid CB₂ agonist, suppresses pathological pain in preclinical models without unwanted central side effects of CB₁ agonists. However, recent *in vitro* studies suggest that GW405833 may also behave as a non-competitive CB₁ antagonist, suggesting that its pharmacology is more complex than initially appreciated. Here, we further investigated the pharmacological specificity of *in vivo* antinociceptive actions of GW405833 in models of neuropathic (i.e. partial sciatic nerve ligation (PSNL) model) and inflammatory (i.e. complete Freund's adjuvant (CFA) model) pain using CB₂ and CB₁ knockout (KO) mice, their respective wild-type (WT) mice and both CB₂ and CB₁ antagonists. GW405833 (3, 10, and 30 mg/kg i.p.) dose-dependently reversed established mechanical allodynia in both pain models in WT mice. However, the anti-allodynic effects of GW405833 were fully preserved in CB₂KO mice and absent in CB₁KO mice. Furthermore, anti-allodynic efficacy of GW405833 (30 mg/kg i.p.) was completely blocked by the CB₁ antagonist rimonabant (10 mg/kg i.p.) but not by the CB₂ antagonist SR144528 (10 mg/kg i.p.). Thus, the antinociceptive properties of GW405833 are dependent upon CB₁ receptors. GW405833 (30 mg/kg i.p.) was also inactive in a tetrad of tests measuring cardinal signs of CB₁ activation. Additionally, unlike rimonabant (10 mg/kg i.p.), GW405833 (10 mg/kg, i.p.) did not act as a CB₁ antagonist *in vivo* to precipitate withdrawal in mice treated chronically with Δ^9 -tetrahydrocannabinol. The present results suggest that anti-allodynic efficacy of GW405833 is CB₁-dependent but does not seem to involve engagement of the CB₁ receptor's orthosteric site.

INTRODUCTION

Cannabinoids exhibit antinociceptive effects in behavioral and electrophysiological studies in rodents (Iversen and Chapman, 2002; Guindon and Hohmann, 2009). There are two well characterized cannabinoid receptors, cannabinoid receptor 1 (CB₁) (Matsuda *et al.*, 1990) and cannabinoid receptor 2 (CB₂) (Munro *et al.*, 1993). Agonists that bind to CB₁ receptors have many desirable therapeutic properties but also induce undesirable psychoactive side effects due to the abundant expression of CB₁ receptors in the central nervous system (CNS) (Herkenham *et al.*, 1991; Galiegue *et al.*, 1995). By contrast, CB₂ receptors are mainly restricted to the periphery and immune tissues (Galiegue *et al.*, 1995), but may nonetheless be induced in the CNS by inflammation or injury (Zhang *et al.*, 2003; Guindon and Hohmann, 2008; Viscomi *et al.*, 2009; Atwood and Mackie, 2010). Therefore, much research interest has emerged in developing CB₂ agonists as suitable analgesics.

GW405833 (also known as L768242; see Fig. 1) binds with preferential affinity to CB₂ over CB₁ receptors (Gallant, 1996; Valenzano *et al.*, 2005; Yao *et al.*, 2008, Table 1). Anti-hyperalgesic effects of GW405833 were first described in a carrageenan model of inflammatory nociception (Clayton *et al.*, 2002). GW405833 was subsequently shown to reduce allodynia in other inflammatory pain (Valenzano *et al.*, 2005; Whiteside *et al.*, 2005; Beltramo *et al.*, 2006), traumatic nerve injury (Valenzano *et al.*, 2005; Whiteside *et al.*, 2005; Beltramo *et al.*, 2006; Brownjohn and Ashton, 2012; Hu *et al.*, 2009; Leichsenring *et al.*, 2009) and incisional injury (LaBuda *et al.*, 2005; Valenzano *et al.*, 2005) models. Despite substantial penetration to the CNS (Bouchard *et al.*, 2012), GW405833 did not produce cannabimimetic deficits below doses of 100 mg/kg i.p. (Valenzano *et al.*, 2005; Whiteside *et al.*, 2005). Anti-allodynic efficacy of GW405833 was opioid independent as systemic administration of naltrexone did not block the anti-hyperalgesic or antinociceptive effects of GW405833 (Whiteside *et al.*, 2005). Thus, GW405833 was advanced as a more specific CB₂ agonist compared to AM1241, which exhibited naloxone-sensitivity under some (Ibrahim *et al.*, 2005) but not all conditions (Rahn *et al.*, 2010) and substantial protean agonism (Yao *et al.*, 2009).

In *in vitro* studies, GW405833 was reported to behave as a partial agonist at human CB₂ receptors (Valenzano *et al.*, 2005) and, alternately, a potent inverse agonist at both human and rat CB₂ receptors and weak agonist at rat CB₁ receptors (Yao *et al.*, 2008). GW405833 was further

JPET #241901

proposed to be a protean agonist whose pharmacological properties were dependent on the constitutive activity of the CB₂ receptor (Mancini *et al.*, 2009). More recently, GW405833 was suggested to act as a non-competitive CB₁ antagonist as GW405833 non-competitively antagonized CP55,940-induced adenylyl cyclase activity, ERK1/2 phosphorylation, PIP₂ signaling and CB₁ internalization *in vitro* in HEK cells transfected with CB₁ and showed a complex, time dependent effect on arrestin recruitment in CHO cells (Dhopeshwarkar *et al.*, 2017).

Anti-allodynic effects of GW405833 have been reported in several preclinical pain models, but pharmacological specificity has been poorly characterized and discrepancies exist among studies. In a few studies, the anti-hyperalgesic effect of GW405833 was blocked by the CB₂ antagonist SR144528 (Clayton *et al.*, 2002; Beltramo *et al.*, 2006), whereas effects of GW405833 did not differ from vehicle in CB₂ KO mice (Valenzano *et al.*, 2005). The possible involvement of CB₁ was never investigated in these studies. By contrast, the antinociceptive effect of GW405833 in reducing lactic acid-stimulated stretching was partially blocked by the CB₁ antagonist rimonabant but not by the CB₂ antagonist SR144528, whereas GW405833-induced attenuation of acid-induced depression of intracranial self-stimulation was not blocked by either SR144528 or rimonabant (Kwilasz and Negus, 2013). Because of these discrepancies, we elected to further characterize the pharmacological specificity of GW405833 in two mechanistically distinct animal pain models, neuropathic pain induced by partial sciatic nerve ligation (PSNL) and inflammatory pain induced by intraplantar injection of complete Freund's adjuvant (CFA). To characterize pharmacological specificity of the *in vivo* profile of GW405833 we used CB₂ knockout (CB₂KO), CB₁ knockout (CB₁KO), and respective wild-type (WT) mice as well as CB₁ and CB₂ antagonists. The highest effective dose of GW405833 (30 mg/kg i.p.) was also evaluated for cardinal signs of CNS CB₁ activation in the cannabinoid tetrad of tests. GW405833 also behaved as non-competitive CB₁ antagonist *in vitro* (Dhopeshwarkar *et al.*, 2017), but it remains unclear whether the reported *in vitro* effects could account for the pharmacological effects of this compound observed *in vivo*. We, therefore also evaluated whether GW405833 would behave similarly to the CB₁ antagonist rimonabant and precipitate a withdrawal syndrome in mice treated chronically with Δ^9 -tetrahydrocannabinol (Δ^9 -THC).

JPET #241901

MATERIALS AND METHODS

Subjects

Mice (5 ± 2 months) of both sexes from different strains were used as specified in each study. CB₂KO mice and WT littermates on a C57BL/6J background (The Jackson Laboratory, Bar Harbor, ME) and CB₁KO mice and WT mice on a CD1 background (Charles River Laboratories, Wilmington, MA) were included. Animals were single-housed at relatively constant temperature ($73 \pm 2^\circ\text{F}$) and humidity (45%) under light-dark cycles of 12/12 h. All the experimental procedures were approved by Bloomington Institutional Animal Care and Use Committee of Indiana University and followed the guidelines for the treatment of animals of the International Association for the Study of Pain (Zimmermann, 1983).

Drugs and chemicals

GW405833 was purchased from two different commercial sources (Tocris Bioscience, Bristol, United Kingdom and Cayman Chemical Company, Ann Arbor, MI, USA). CFA and WIN55, 212-2 were purchased from Sigma-Aldrich (St. Louis, MO). Rimonabant (SR141716A), SR144528 and Δ^9 -THC were obtained from the National Institutes of Health Drug Supply Program (Research Triangle Institute, RTP, NC). All drugs except Δ^9 -THC were dissolved in a vehicle of DMSO (Sigma-Aldrich, St. Louis, MO), emulphor (Alkamuls EL 620L, Solvay, Princeton, NJ), ethanol (Sigma-Aldrich, St. Louis, MO), and sterile saline (Aquilite System, Hospira Inc, Lake Forest, IL) at a ratio of 5:2:2:16 respectively. Δ^9 -THC was dissolved in a vehicle of 95% ethanol, Cremophor® EL (Sigma Aldrich, St. Louis, MO) and sterile saline (Aquilite System; Hospira, Inc, Lake Forest, IL) in a ratio of 1:1:18 respectively. All compounds were delivered via intraperitoneal injection (i.p.) in a volume of 5 ml/kg.

Partial sciatic nerve ligation (PSNL)

PSNL was performed to induce neuropathic pain. Mice were anesthetized with isoflurane (5% for induction followed by 2-2.5 % for maintenance) in 1.5 L/min of oxygen. The right high thigh area was then shaved and disinfected followed by a 1 to 1.5 cm longitudinal incision. The muscle layers were bluntly dissected to expose the underlying sciatic nerve. One third to half of the sciatic nerve at the location just above its trifurcation was tightly ligated using 8-0 silk suture (Sharpoint DA-2526N). The muscle layers were then closed with 5-0 silk thread (Ethicon 682 G)

JPET #241901

and the skin was closed by staples (Autoclips 9mm No 7631). Animals were allowed at least two weeks to recover and fully develop neuropathic pain.

Complete Freund's adjuvant (CFA)-induced inflammatory pain

A single intraplantar (i.pl.) injection was used to deliver CFA (20 μ l; diluted 1:1 in sterile saline) unilaterally into the right hind paw. Animals were given 48 hours to fully develop inflammatory pain prior to pharmacological manipulations.

Assessment of mechanical allodynia

Animals were habituated in individual transparent Plexiglass test chambers (10.5 x 9 x 7 cm) atop a metal mesh platform for at least 30 min prior to the testing. An electronic von Frey anesthesiometer with a 90-g probe (IITC Life Science, Woodland Hills, California) was employed to determine the paw withdrawal threshold to mechanical stimulation of the hind paw. A rigid tip with a diameter of 0.8 mm attached to the probe was applied vertically against the plantar surface of the paw with gradually increasing force. The force in grams necessary to produce paw withdrawal was recorded in duplicate for each paw.

Baseline mechanical paw withdrawal thresholds were measured in each paw for each animal prior to performing either PSNL or injecting CFA. Another pre-drug baseline was then taken after painful peripheral neuropathy or inflammatory pain was fully established. GW405833 was administered (i.p.) 30 min prior to evaluation of the impact of drug manipulations on mechanical paw withdrawal thresholds. Different doses of GW405833 were injected (i.p.) within subjects in the order of vehicle (0), 3, 10, and 30 mg/kg. Sufficient time was allowed to lapse between each dose to verify that mechanical paw withdrawal thresholds returned to the pre-drug levels prior to dose escalation. For detailed timeline information, please see Fig 1B and C. In the groups where CB₁ or CB₂ antagonists were tested, rimonabant or SR144528 (10 mg/kg i.p.) were administered 20 min prior to GW405833 injection.

Tetrad test

Rotarod

Motor performance was measured using an accelerating Rotarod (IITC Life Science) (4-40 rpm, 300 s cutoff time). Male CD1 mice were trained over two consecutive days and on day

JPET #241901

three the baseline latencies to descend from the rotating drum were measured. Mice that did not meet exclusion criteria prior to baseline measurements (i.e. ability to stay on rotating drum for at least 30 s on baseline day) were removed from the study and did not receive pharmacological treatments.

Rectal Temperature

Rectal temperature was measured to assess hypothermia by insertion of a mouse rectal probe (Braintree Laboratories Inc., Braintree, MA.) 1.2 cm into the rectum using a thermometer (Physitemp Instruments Inc., Clifton, NJ).

Tail-flick test

To assess the effects of drug treatments on acute antinociception, mice were gently restrained in a towel and the tip of the tail (~1 cm) was submerged in a hot water bath maintained at 54-55 °C. The latency to withdraw the tail was measured.

Ring test

The ring test (Pertwee, 1972) was performed to assess possible cataleptic effects produced by drug administration. Briefly, each mouse was positioned on top of an elevated ring attached to a vertical stainless steel rod suspended at a height of 16 – 17 cm above a flat platform for a total duration of 5 min. The amount of time spent immobile was recorded as an index for catalepsy.

Behavioral assessment of the development of tolerance to Δ^9 -THC

Male mice of the C57 background strain received once daily injections of Δ^9 -THC (50 mg/kg i.p.) for 9 consecutive days. Mice were assessed in three of the behavioral assays from the tetrad test to quantify the development of tolerance to the behavioral effects of Δ^9 -THC. Motor coordination was measured using the rotarod test, body temperature was measured using a rectal temperature probe and antinociception was measured using the tail-flick test. Responses were measured at baseline and 30 minutes following administration of Δ^9 -THC on days 1 and 8 of Δ^9 -THC administration. Mice received Δ^9 -THC in the same environment each day and remained for 4 hours post- Δ^9 -THC injection prior to being returned to the colony room.

Assessment of antagonist-precipitated withdrawal symptoms in Δ^9 -THC-treated mice

JPET #241901

On day 9 of Δ^9 -THC administration, all mice were challenged with vehicle (i.p.) followed 30 min later by a second challenge (i.p.) with either rimonabant (10 mg/kg) or GW405833 (10 mg/kg). Mice were placed in clear Plexiglass chambers on an observation platform and withdrawal behaviors were videotaped and scored by an investigator blinded to treatment conditions. On the day of withdrawal testing, mice were injected with Δ^9 -THC (50 mg/kg, i.p.) and allowed to acclimate to the chamber for 30 minutes. Next, all mice received a single injection of vehicle (i.p.) and behavior was videotaped for 30 minutes. Immediately following vehicle challenge, mice received a single i.p. injection of either rimonabant or GW405833 and behavior was videotaped for 30 minutes. The number of front paw tremors, headshakes, scratching, grooming and rearing behaviors were counted. All mice received vehicle followed by drug challenge so within subjects comparisons could be made.

Statistical analysis

Differences in mechanical paw withdrawal thresholds were evaluated using mixed ANOVA (with different pharmacological treatments as the within-subject variable and genotype as between-subject variable) followed by Bonferroni post hoc tests. For tetrad tests, one way ANOVA was performed separately on pre-drug (baseline) and post-drug responses, followed by Bonferroi post hoc tests. Paired or independent t-tests were performed with the specific comparisons of interest as indicated. Analysis of variance for repeated measures was used to determine the time course of tolerance development to the effects of Δ^9 -THC, followed by paired-samples t-tests to confirm the development of tolerance. Paired samples t-tests were used to assess differences in withdrawal symptoms elicited by vehicle vs. drug challenge. Statistical analyses were performed using IBM SPSS Statistics version 24 (IBM Corporation, Armonk, New York) or GraphPad Prism Version 5.02 statistical software for windows (GraphPad Software, San Diego, CA USA; www.graphpad.com). Figures were generated using GraphPad Prism Version 5.02 statistical software. All data are presented as mean \pm SEM, $p < 0.05$ was considered significant.

RESULTS

GW405833-induced suppressions of neuropathic pain in the PSNL model are CB₁ receptor-dependent and CB₂ receptor-independent.

JPET #241901

GW405833-induced anti-allodynic effects in PSNL model are fully preserved in CB₂ KO mice and are not blocked by a CB₂ antagonist.

Partial sciatic nerve ligation lowered mechanical paw withdrawal thresholds in CB₂KO and respective WT mice relative to baseline, consistent with the development of neuropathic pain ($F_{1,10} = 27.434, p < 0.001$) but these effects did not differ between WT and CB₂ KO mice ($F_{1,10} = 0.003, p = 0.988$) (Fig. 2A). Vehicle (0 mg/kg, i.p.) alone did not alter mechanical paw withdrawal thresholds in either group ($p > 0.05$) (Fig. 2A). ANOVA revealed a main effect of GW405833 treatment on mechanical allodynia ($F_{3,30} = 11.381, p < 0.001$) and Bonferroni post hoc tests indicated that GW405833 elevated paw withdrawal thresholds in a dose-dependent manner (Fig. 2A). No main effect of genotype ($F_{1,10} = 3.620, p = 0.086$) or significant interactions were observed ($p = 0.598$) (Fig. 2A). These results indicate that the anti-allodynic effect of GW405833 was fully preserved in CB₂KO mice. GW405833 (10 and 30 mg/kg i.p.) fully restored mechanical paw withdrawal thresholds relative to baseline at doses as low as 10 mg/kg i.p. in both WT ($t_5 = 0.270, p = 0.798$) and CB₂KO ($t_5 = 1.959, p = 0.107$) mice (Fig. 2A). Paired t tests indicated that the CB₂ antagonist SR144528 (10 mg/kg i.p.) did not block the anti-allodynic efficacy of GW405833 (30 mg/kg i.p.) in CB₂KO mice ($t_5 = 1.790, p = 0.133$), but attenuated the efficacy of GW405833 in WT mice ($t_5 = 2.833, p = 0.037$). Although anti-allodynic efficacy of GW405833 (30 mg/kg i.p.) was attenuated in WT mice by SR144528, residual efficacy was, nonetheless, comparable to that observed in WT mice receiving GW405833 alone at a highly efficacious dose of 10 mg/kg i.p. ($t_5 = 0.800, p = 0.460$). Moreover, the anti-allodynic efficacy of GW405833 in WT mice did not differ between doses of 10 mg/kg i.p. and 30 mg/kg i.p. ($p = 1$). These observations suggest that, in WT mice, SR144528-induced blockade of the anti-allodynic efficacy of GW405833, if present at all, was minimal.

GW405833-induced anti-allodynic effects in PSNL model are absent in CB₁ KO mice and are blocked by a CB₁ antagonist.

Partial sciatic nerve ligation lowered mechanical paw withdrawal thresholds in CB₁KO and respective WT mice relative to baseline, consistent with the development of robust mechanical allodynia ($F_{1,14} = 42.594, p < 0.001$), but these effects did not differ between WT and CB₁ KO mice ($F_{1,14} = 1.079, p = 0.317$) (Fig. 3A). Vehicle (0 mg/kg, i.p.) did not alter mechanical paw withdrawal thresholds after the neuropathic pain was fully established ($p >$

JPET #241901

0.05). There was a main effect of GW405833 on mechanical allodynia ($F_{3,42} = 5.025, p = 0.005$), a main effect of genotype ($F_{1,14} = 14.820, p = 0.002$), and a trend was observed for the interaction that approached statistical significance ($F_{3,42} = 2.487, p = 0.074$) (Fig. 3A). Bonferroni post hoc tests indicated that GW405833 dose-dependently elevated mechanical paw withdrawal thresholds in WT mice, but these anti-allodynic effects were absent in CB₁KO mice (Fig. 3A). The highest dose of GW405833 (30 mg/kg i.p.) did not fully restore mechanical paw withdrawal thresholds to baseline (pre-injury) levels ($t_7 = 3.33, p = 0.013$). In WT mice, pretreatment with the CB₁ antagonist rimonabant (10 mg/kg i.p.) before GW405833 (30 mg/kg i.p.) completely blocked the anti-allodynic effect of GW405833 ($t_7 = 3.097, p = 0.017$), and the resulting mechanical thresholds did not differ from those of CB₁KO mice receiving GW405833 alone (30 mg/kg i.p.) ($t_{14} = -0.582, p = 0.597$) (Fig. 3A).

Neither PSNL nor pharmacological manipulations altered the mechanical thresholds in the contralateral (uninjured) paw in any genotype ($p > 0.27$) (Fig. 2B & 3B).

GW405833-induced anti-allodynic effects in the CFA model of inflammatory pain are CB₁ receptor-dependent and CB₂ receptor-independent.

GW405833-induced anti-allodynic effects in CFA model are fully preserved in CB₂ KO mice and are not blocked by a CB₂ antagonist.

Intraplantar CFA injection reduced mechanical paw withdrawal thresholds in both WT and CB₂KO mice ($F_{1,14} = 222.594, p < 0.001$) (Fig. 4A). There was a main effect of GW405833 treatment on mechanical allodynia ($F_{3,42} = 37.257, p < 0.001$), and Bonferroni post hoc tests indicated that GW405833 suppressed CFA-induced allodynia in a dose-dependent manner (Fig. 4A). There was no main effect of genotype ($F_{1,14} = 0.146, p = 0.708$) or interaction ($F_{3,42} = 1.156, p = 0.338$). Efficacy of the 10 mg/kg and 30 mg/kg i.p. doses of GW405833 did not differ from each other ($p = 0.236$). Together, these data indicate that the anti-allodynic efficacy of GW405833 was fully preserved in CB₂KO mice relative to WT mice. The highest dose of GW405833 (30 mg/kg i.p.) did not produce a full reversal of mechanical allodynia in WT relative to pre-CFA baseline ($t_7 = 5.024, p = 0.002$), but there was no statistically significant difference between pre-CFA baseline and GW405833 (30 mg/kg i.p.)-reversed mechanical withdrawal thresholds in CB₂KO ($t_7 = 2.097, p = 0.074$) mice. Moreover, there was no difference in responding between WT and CB₂KO mice ($t_{14} = 0.437, p = 0.165$) (Fig. 4A).

JPET #241901

Paired-samples t-tests showed that the CB₂ antagonist SR144528 did not block the anti-allodynic efficacy of GW405833 in either WT ($t_7 = 0.858, p = 0.419$) or CB₂KO ($t_7 = 0.760, p = 0.472$) mice (Fig. 4A).

GW405833-induced anti-allodynic effects in CFA model are absent in CB₁ KO mice and blocked by a CB₁ antagonist in WT mice.

Intraplantar CFA injection lowered mechanical paw withdrawal thresholds in both WT and CB₁KO mice, suggesting that both genotypes developed inflammatory pain ($F_{1,12} = 145.506, p < 0.001$) (Fig. 5A). Vehicle (i.p.) did not alter mechanical paw withdrawal thresholds after CFA-induced inflammation was established ($p > 0.05$). There was a main effect of GW405833 treatment on mechanical allodynia ($F_{3,36} = 26.618, p < 0.001$), a main effect of genotype ($F_{1,12} = 8.348, p < 0.001$), and the interaction was significant ($F_{3,36} = 8.212, p = 0.014$) (Fig. 5A). Bonferroni post hoc tests revealed that GW405833 produced a dose-dependent elevation of mechanical thresholds in WT mice, but such anti-allodynic effects were absent in CB₁KO mice (Fig. 5A). The highest dose of GW405833 (30 mg/kg i.p.) fully restored mechanical paw withdrawal thresholds relative to baseline (pre-CFA) levels ($t_7 = 1.374, p = 0.212$) in WT mice. Moreover, the CB₁ antagonist rimonabant completely blocked the anti-allodynic efficacy of GW405833 in WT mice ($t_7 = 5.58, p = 0.001$) (Fig. 5A). Mechanical paw withdrawal thresholds observed in CB₁KO mice treated with GW405833 (30 mg/kg i.p.) did not differ from those observed in WT mice pretreated with rimonabant prior to GW405833 (30 mg/kg i.p.) ($t_{12} = 1.150, p = 0.189$) (Fig. 5A). Moreover, local administration of GW405833 (30 µg or 300 µg in 10 µl) in the CFA-injected paw of CB₁ KOs did not produce anti-allodynic efficacy (data not shown).

Neither CFA injection nor pharmacological manipulations altered mechanical thresholds in the contralateral (non-inflamed) paw in any genotype ($p > 0.33$) (Fig. 4B & 5B).

GW405833 does not induce typical cannabinimimetic effects in the tetrad tests.

Prior to pharmacological manipulations, responding did not differ between the three groups in either the ring test ($F_{2,15} = 0.344, p = 0.715$), rota-rod test ($F_{2,15} = 0.027, p = 0.973$), rectal temperature assessment ($F_{2,15} = 0.443, p = 0.650$) or tail-flick test ($F_{2,15} = 0.411, p = 0.670$) (Fig. 6A-D). Pharmacological manipulations altered post-drug responding in the ring test

JPET #241901

($F_{2,15} = 31.010, p < 0.001$) (Fig. 6A). Bonferroni post hoc tests revealed that the positive control WIN55, 212-2 (5 mg/kg i.p.) increased immobility in animals compared to the vehicle or GW405833 (30 mg/kg i.p.) injection ($p < 0.001$) (Fig. 6A). Immobility time in the ring test in groups receiving GW405833 did not differ from vehicle ($p = 1$) or the pre-injection responding ($p = 0.156$) (Fig. 6A). Therefore, GW405833 (30 mg/kg i.p.) failed to induce catalepsy.

Drug manipulations tended to alter rota-rod descend latencies ($F_{2,15} = 3.264, p = 0.067$). Nonetheless, planned comparison independent t-tests revealed that rotarod latencies were lower in groups receiving WIN55, 212-2 (5 mg/kg i.p.) compared to GW405833 ($p = 0.046$). By contrast, effects of GW405833 (30 mg/kg i.p.) did not differ from vehicle ($p = 0.302$). Thus, GW405833 at a dose of 30 mg/kg i.p. did not produce motor ataxia in WT mice.

Pharmacological manipulations altered body temperature ($F_{2,15} = 52.875, p < 0.001$). Bonferroni post hoc tests revealed that the positive control WIN55, 212-2 (5 mg/kg i.p.) reduced rectal temperature compared to body temperatures measured in mice receiving either vehicle or GW405833 ($p < 0.001$ for each comparison) (Fig. 6C). By contrast, post-injection body temperatures in mice receiving GW405833 did not differ from those observed in mice receiving vehicle ($p = 0.555$) and did not differ from pre-injection (baseline) body temperatures ($p = 0.119$) (Fig. 6C). Therefore, GW405833 failed to induce hypothermia.

Pharmacological manipulations altered post-injection tail-flick latencies ($F_{2,15} = 7.993, p = 0.004$). Bonferroni post hoc tests indicated that the positive control WIN55, 212-2 (5 mg/kg i.p.) increased tail-flick latencies compared to either vehicle or GW405833 treatments ($p < 0.05$) (Fig. 6D). By contrast, post-injection tail-flick latencies in groups receiving GW405833 did not differ from vehicle ($p = 0.772$) or the pre-injection latency ($p = 0.691$) (Fig. 6D). Therefore, GW405833 failed to induce acute tail-flick antinociception.

Mice develop tolerance to motor ataxic, hypothermic and antinociceptive properties of Δ^9 -THC.

Prior to withdrawal manipulations, there were no differences between any of the groups in any of the dependent measures used to assess the development of tolerance ($p > 0.1$). Therefore, pre-withdrawal responses of all mice were pooled for statistical analysis of tolerance development (Fig. 7).

JPET #241901

Mice treated chronically with Δ^9 -THC (50 mg/kg/day i.p. x 8 days) exhibited motor impairment ($F_{2,20} = 108.8$, $p < 0.0001$; Fig. 7A), and there was no main effect of group ($p > 0.8$) or interaction ($p > 0.3$). Δ^9 -THC produced motor impairment on day 1 of administration relative to baseline ($t_{11} = 8.899$, $p < 0.0001$; Fig. 7A) but full tolerance developed by day 8 of drug administration ($t_{11} = 12.39$, $p < 0.0001$; Fig. 7A), indicating that animals developed tolerance to the motor ataxic effects of Δ^9 -THC. Latencies to descend from the rotarod were also higher on day 8 relative to baseline ($t_{11} = 6.154$, $p < 0.0001$; Fig. 7A), indicating that rotarod performance may have improved with additional training when acute motor impairment from Δ^9 -THC was no longer present.

Δ^9 -THC decreased rectal temperature ($F_{2,20} = 140.2$, $p < 0.0001$; Fig. 7B), and there was no main effect of group ($p > 0.3$) or interaction ($p > 0.3$). Δ^9 -THC decreased rectal temperature on day 1 of administration relative to baseline ($t_{11} = 14.66$, $p < 0.0001$; Fig. 7B) but reached ~95% tolerance by day 8 of administration ($t_{11} = 10.62$, $p < 0.0001$; Fig. 7B), indicating mice had become tolerant to the hypothermic effects of Δ^9 -THC. Δ^9 -THC also decreased rectal temperature on day 8 relative to baseline ($t_{11} = 2.367$, $p < 0.05$; Fig. 7B).

In the hot water tail-flick test, Δ^9 -THC treatment (i.p.) produced antinociception in all mice ($F_{2,20} = 19.18$, $p < 0.0001$; Fig. 7C) and there was no effect of group ($p > 0.1$) or interaction ($p > 0.4$). Δ^9 -THC increased the latency to withdraw the tail on day 1 relative to baseline ($t_{11} = 4.644$, $p < 0.001$; Fig. 7C) and responding on day 8 ($t_{11} = 4.134$, $p < 0.01$; Fig. 7C) of administration. Tail withdrawal latencies did not differ on day 8 of administration relative to baseline ($p > 0.1$), indicative of the development of tolerance to the antinociceptive effects of Δ^9 -THC.

Rimonabant elicits classical signs of cannabinoid withdrawal while GW405833 does not.

In mice chronically treated with Δ^9 -THC (50 mg/kg/day x 9 days, i.p.), rimonabant challenge (10 mg, i.p.) increased paw tremors ($t_5 = 3.95$, $p < 0.05$; Fig. 8A), head shakes ($t_5 = 2.716$, $p < 0.05$; Fig. 8A), grooming behavior ($t_5 = 2.963$, $p < 0.05$; Fig. 8A) and rearing behaviors ($t_5 = 3.943$, $p < 0.05$, Fig. 8A) relative to vehicle (i.p.) challenge.

JPET #241901

In mice chronically treated with Δ^9 -THC (50 mg/kg/day x 9 days, i.p.), GW405833 (10 mg/kg, i.p.) decreased paw tremors ($t_5 = 3.955$, $p < 0.05$; Fig 8B), but failed to alter head shakes, scratching, grooming or rearing behaviors ($p > 0.05$) relative to vehicle challenge (i.p.).

DISCUSSION

GW405833 has been widely described in the literature as a cannabinoid CB₂ agonist, although the pharmacological specificity of its *in vivo* profile of actions has not been thoroughly evaluated. We characterized the pharmacological specificity of the anti-allodynic effects of GW405833 using both neuropathic (i.e. induced by PSNL) and inflammatory (i.e. induced by CFA) pain models in mice. Consistent with other studies (Valenzano *et al.*, 2005; Whiteside *et al.*, 2005), GW405833 dose-dependently reversed mechanical hypersensitivity in both neuropathic and inflammatory pain models following systemic administration. GW405833 (3 mg/kg i.p.) did not produce anti-allodynic efficacy but doses of 10 and 30 mg/kg i.p. robustly reversed established allodynia. Neither PSNL, CFA nor GW405833 treatment altered mechanical thresholds in the contralateral paw, suggesting that GW405833 suppressed established allodynia induced by neuropathic or inflammatory insults without altering basal nociceptive thresholds observed in the absence of injury.

The most striking observation of our study was that anti-allodynic efficacy of GW405833 was not mediated by CB₂ receptors. GW405833-induced antinociception was, in fact, fully preserved in CB₂ KO mice in models of established neuropathic and inflammatory pain. GW405833 reversed established allodynia with similar efficacy in WT and CB₂KO mice in both pain models. By contrast, anti-allodynic efficacy of GW405833 was absent in CB₁ KO mice and blocked in WT mice by the CB₁ antagonist rimonabant. In neuropathic WT mice, the CB₂ antagonist SR144528 (10 mg/kg i.p.) produced only a modest attenuation of the anti-allodynic efficacy of GW405833 (30 mg/kg i.p.); anti-allodynic efficacy was suppressed to a level comparable to that of a fully efficacious lower dose (10 mg/kg i.p.). Moreover, anti-allodynic efficacy of GW405833 was not blocked at all by SR144528 in the CFA model. The antinociceptive effects of GW405833 are widely believed to be CB₂ receptor-mediated because of the high selectivity of GW405833 to CB₂ receptor over CB₁ receptor described *in vitro* (Valenzano *et al.*, 2005), and reports of blockade of GW405833 effects by a CB₂ antagonist (Clayton *et al.*, 2002; Beltramo *et al.*, 2006), and absence of GW405833 antinociceptive efficacy

JPET #241901

in CB₂KO mice (Valenzano *et al.*, 2005). However, our results suggest that GW405833 does not behave as a CB₂ agonist in mice following systemic administration.

Strikingly, the anti-allodynic effects of GW405833 observed in the current study were CB₁ receptor-mediated. These observations are consistent with a previous report suggesting that the antinociceptive effects of GW405833 on lactic acid-induced nociceptive stretch responses were partially blocked by the CB₁ antagonist rimonabant but not by the CB₂ antagonist SR144528 (Kwilasz and Negus, 2013). In the few studies suggesting the antinociceptive effects of GW405833 are CB₂-mediated (Clayton *et al.*, 2002; Valenzano *et al.*, 2005; Beltramo *et al.*, 2006), neither CB₁ antagonists nor CB₁KOs were employed to investigate the possible contribution of CB₁ receptors to the *in vivo* antinociceptive profile of GW405833. Only one prior study included CB₂KO mice (Valenzano *et al.*, 2005). To our knowledge, our study is the first to include CB₁KO mice to study possible involvement of CB₁ receptors in the anti-allodynic effects of GW405833. GW405833 has been reported to behave as a partial agonist of the orphan G protein-coupled receptor 55 (GPR55) in evaluation of L-alpha-lysophosphatidylinositol (LPI) – induced mitogen-activated protein kinase (MAPK) activation *in vitro* (Anavi-Goffer *et al.*, 2012). However, it is unclear how GPR55 agonism would produce antinociception. The involvement of GPR55 in pain is controversial (Staton *et al.*, 2008; Carey *et al.*, 2017) and difficult to address due to lack of sufficiently selective agonists and antagonists for this receptor.

In the two studies where the CB₂ antagonist SR144528 was used to block the antinociceptive effect of GW405833 (Clayton *et al.*, 2002; Beltramo *et al.*, 2006), different pain models (i.e. carrageenan and formalin inflammatory pain models), different methodologies (i.e. weight bearing, paw edema, paw licking and flinching) and different species (i.e. rats were used by Clayton and colleagues) were used compared with our study, which may contribute to the different pattern of results obtained. One possibly noteworthy methodological difference is that GW405833 was administered before the noxious insult in both prior studies, while in our study GW405833 was administered after the establishment of both neuropathic pain and inflammatory pain. Thus, it is possible that GW405833 blocks pain through a CB₂ mediated mechanism during the development of pain, but through a CB₁-mediated mechanism during the maintenance of pathological pain. In the study by Valenzano *et al.* (2005), the anti-allodynic effect of GW405833 (30 mg/kg i.p.) in the CFA model in CB₂KO mice is reported to not differ from

JPET #241901

vehicle. However, this latter study did not evaluate whether efficacy of GW405833 differed statistically in WT and CB₂KO mice. Inadequate statistical power could account for failure to observe differences in responding between CB₂KO mice receiving vehicle and GW405833 at the single time point evaluated previously (Valenzano *et al.*, 2005). By contrast, in our study, GW405833 exhibited similar anti-allodynic effects in CB₂KO and WT mice at doses of 10 mg/kg i.p. and 30 mg/kg i.p. Both our study and that of Valenzano *et al.* (2005) employed CB₂KO mice on the C57BL/6 background, and the same concentration of CFA in the paw (20 µl, 50% CFA diluted in 0.9% saline). The effect of GW405833 (30 mg/kg i.p.) was studied 24 h after CFA injection by Valenzano *et al.*, while in our study GW405833 was tested ≥ 48 h after CFA injection. If GW405833 engages different antinociceptive mechanisms during different stages of the maintenance of inflammatory pain, then inclusion of different testing time points between studies could potentially contribute to the observed discrepancies.

Although GW405833 binds with higher affinity to the orthosteric site on CB₂ receptors relative to CB₁ receptors *in vitro*, the degree of selectivity for CB₂ over CB₁ is quite variable among studies, ranging from 37 fold up to ~1200 fold selective to human CB₂ receptors over human CB₁ receptors (Gallant, 1996; Valenzano *et al.*, 2005; Yao *et al.*, 2008). Although it is possible that species differences exist between rat and mouse CB₂ receptors, they are much more highly conserved compared to human CB₂ receptors (Liu *et al.*, 2008). Binding of GW405833 to native mouse CB₂ receptors, to our knowledge, has never been determined. Considering that GW405833 penetrates into the CNS at high levels (Valenzano *et al.*, 2005; Evens *et al.*, 2011), cannabimimetic CB₁-like effects could be expected in the tetrad tests if GW405833 binds directly to the orthosteric binding site of CB₁ receptors. However, the highest effective dose of GW405833 (30 mg/kg i.p.) identified in the present study was inactive in the tetrad tests. Specifically, GW405833 (30 mg/kg i.p.) did not produce catalepsy, motor ataxia, hypothermia or acute antinociception in the tail-flick test in mice. These observations are in agreement with prior studies (Valenzano *et al.*, 2005; Whiteside *et al.*, 2005) suggesting that GW405833 did not produce cannabimimetic effects at doses lower than 100 mg/kg i.p. A recent paper has shown that GW405833 can act as a non-competitive CB₁ antagonist *in vitro*, because it non-competitively antagonized CP55,940-induced adenylyl cyclase inhibition, ERK1/2 phosphorylation, PIP₂ signaling and CB₁ internalization *in vitro* (Dhopeswarkar *et al.*, 2017). In addition, GW405833 induced a complex, time-dependent activation of arrestin through CB₁

JPET #241901

(Dhopeshwarkar *et al.*, 2017). Therefore, to assess whether GW405833 behaves as an orthosteric CB₁ antagonist *in vivo* we evaluated whether challenge with GW405833 would induce CB₁-dependent cannabinoid withdrawal in mice treated chronically with Δ^9 -THC (50 mg/kg, i.p. x 9 days). In contrast to challenge with the classical CB₁ antagonist/inverse agonist rimonabant (10 mg/kg, i.p.), challenge with GW405833 (10mg/kg, i.p.) did not induce classic withdrawal responses in THC-tolerant mice. The lack of precipitated withdrawal by GW405833 observed in our withdrawal assay indicates that GW405833 does not behave as a CNS-penetrant competitive CB₁ antagonist *in vivo*. Thus, caution must be exerted before attributing the mechanism of action underlying *in vivo* effects of novel ligands to mechanisms identified *in vitro*.

In summary, the anti-allodynic effect of GW405833 in the established neuropathic and inflammatory pain models examined here is CB₁-mediated and not CB₂-mediated. The anti-allodynic effects of GW405833 were absent in CB₁KO mice and blocked by a CB₁ antagonist, but were fully preserved in CB₂KO mice and were largely unaffected by a CB₂ antagonist. Although GW405833 is brain penetrant (Valenzano *et al.*, 2005) and reduced neuropathic and inflammatory pain through a CB₁-mediated mechanism in our study, it did not induce cannabimimetic CB₁-mediated side effects at the highest therapeutic dose evaluated (30 mg/kg i.p.). Moreover, GW405833 (10 mg/kg i.p.) did not function as a competitive CB₁ antagonist in our attempts to precipitate withdrawal with this agent in Δ^9 -THC tolerant mice. Together, our results indicate that GW405833 reversed established neuropathic and inflammatory pain through a CB₁-mediated mechanism without producing characteristic cannabimimetic effects associated with direct orthosteric binding to the CB₁ receptor. GW405833 could possibly act as a CB₁ allosteric modulator or interact with the endocannabinoid system. More research is needed to elucidate the mechanism of antinociceptive actions of GW405833.

JPET #241901

Author Contributions:

Participated in research design and oversaw project: Hohmann

Conducted experiments: Li, Carey

Performed data analysis: Li, Carey

Wrote or contributed to the writing of the manuscript: Li, Carey, Mackie and Hohmann

JPET #241901

References

- Anavi-Goffer S, Baillie G, Irving AJ, Gertsch J, Greig IR, Pertwee RG, and Ross RA (2012) Modulation of L-alpha-lysophosphatidylinositol/GPR55 mitogen-activated protein kinase (MAPK) signaling by cannabinoids. *J Biol Chem* **287**:91–104.
- Atwood BK, and Mackie K (2010) CB 2: A cannabinoid receptor with an identity crisis. *Br J Pharmacol* **160**:467–479.
- Beltramo M, Bernardini N, Bertorelli R, Campanella M, Nicolussi E, Fredduzzi S, and Reggiani A (2006) CB2 receptor-mediated antihyperalgesia: Possible direct involvement of neural mechanisms. *Eur J Neurosci* **23**:1530–1538.
- Bouchard J, Truong J, Bouchard K, Dunkelberger D, Desrayaud S, Moussaoui S, Tabrizi SJ, Stella N, and Muchowski PJ (2012) Cannabinoid receptor 2 signaling in peripheral immune cells modulates disease onset and severity in mouse models of Huntington's disease. *J Neurosci* **32**:18259–68.
- Brownjohn PW, and Ashton JC (2009) Novel targets in pain research: The case for CB2 receptors as a biorational pain target. *Curr Anaesth Crit Care* **20**:198–203, Elsevier Ltd.
- Brownjohn PW, and Ashton JC (2012) Spinal cannabinoid CB2 receptors as a target for neuropathic pain: An investigation using chronic constriction injury. *Neuroscience* **203**:180–193, Elsevier Inc.
- Carey LM, Gutierrez T, Deng L, Lee W-H, Mackie K, and Hohmann AG (2017) Inflammatory and Neuropathic Nociception is Preserved in GPR55 Knockout Mice. *Sci Rep* **7**:944, Springer US.
- Clayton N, Marshall FH, Bountra C, and O'Shaughnessy CT (2002) CB1 and CB2 cannabinoid receptors are implicated in inflammatory pain. *Pain* **96**:253–260.
- Dhopeswarkar A, Murataeva N, Makriyannis A, Straiker A, and Mackie KP (2017) Two Janus cannabinoids that are both CB2 agonists and CB1 antagonists. *J Pharmacol Exp Ther* **360**:300–311.
- Evens N, Vandeputte C, Muccioli GG, Lambert DM, Baekelandt V, Verbruggen AM, Debyser

JPET #241901

- Z, Van Laere K, and Bormans GM (2011) Synthesis, in vitro and in vivo evaluation of fluorine-18 labelled FE-GW405833 as a PET tracer for type 2 cannabinoid receptor imaging. *Bioorganic Med Chem* **19**:4499–4505.
- Galiegue S, Mary S, Marchand J, Dussossoy D, Carriere D, Carayon P, Bouaboula M, Shire D, Fur G, and Casellas P (1995) Expression of Central and Peripheral Cannabinoid Receptors in Human Immune Tissues and Leukocyte Subpopulations. *Eur J Biochem* **232**:54–61.
- Gallant M, Dufresne C, Gareau Y, Guay D, Leblanc Y, Prasit P, Rochette C, Sawyer N, Slipetz DM, Tremblay N, Metters KM, and Labelle M (1996) New class of potent ligands for the human peripheral cannabinoid receptor. *Bioorg Med Chem Lett* **6**:2263–2268.
- Guindon J, and Hohmann AG (2008) Cannabinoid CB2 receptors: a therapeutic target for the treatment of inflammatory and neuropathic pain. *Br J Pharmacol* **153**:319–334.
- Guindon J, and Hohmann AG (2009) The endocannabinoid system and pain. *CNS Neurol Disord Drug Targets* **8**:403–21.
- Herkenham M, Lynn AB, Johnson MR, Melvin LS, de Costa BR, and Rice KC (1991) Characterization and localization of cannabinoid receptors in rat brain: a quantitative in vitro autoradiographic study. *J Neurosci* **11**:563–83.
- Hu B, Doods H, Treede RD, and Ceci A (2009) Depression-like behaviour in rats with mononeuropathy is reduced by the CB2-selective agonist GW405833. *Pain* **143**:206–212, International Association for the Study of Pain.
- Ibrahim MM, Porreca F, Lai J, Albrecht PJ, Rice FL, Khodorova A, Davar G, Makriyannis A, Vanderah TW, Mata HP, and Malan TP (2005) CB2 cannabinoid receptor activation produces antinociception by stimulating peripheral release of endogenous opioids. *Proc Natl Acad Sci U S A* **102**:3093–8.
- Iversen L, and Chapman V (2002) Cannabinoids: A real prospect for pain relief? *Curr Opin Pharmacol* **2**:50–55.
- Kwilasz A, and Negus S (2013) Effects of the cannabinoid 2 receptor-selective agonist GW405833 in assays of acute pain-stimulated and paindepressed behavior in rats. *FASEB J* **27**:886.9-886.9, Federation of American Societies for Experimental Biology.

JPET #241901

LaBuda CJ, Koblisch M, and Little PJ (2005) Cannabinoid CB2 receptor agonist activity in the hindpaw incision: Model of postoperative pain. *Eur J Pharmacol* **527**:172–174.

Leichsenring A, Andriske M, Bäcker I, Stichel CC, and Lübbert H (2009) Analgesic and antiinflammatory effects of cannabinoid receptor agonists in a rat model of neuropathic pain. *Naunyn Schmiedebergs Arch Pharmacol* **379**:627–636.

Liu Q-R, Pan C-H, Hishimoto A, Li C-Y, Xi Z-X, Llorente-Berzal A, Viveros M-P, Ishiguro H, Arinami T, Onaivi ES, and Uhl GR (2008) Species differences in cannabinoid receptor 2 (CNR2 gene): identification of novel human and rodent CB2 isoforms, differential tissue expression, and regulation by cannabinoid receptor ligands. *Genes Brain Behav* **144**:724–732.

Mancini I, Brusa R, Quadrato G, Foglia C, Scandroglio P, Silverman L, Tulshian D, Reggiani A, and Beltramo M (2009) Constitutive activity of cannabinoid-2 (CB 2) receptors plays an essential role in the protean agonism of (+)AM1241 and L768242. *Br J Pharmacol* **158**:382–391.

Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, and Bonner TI (1990) Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* **346**:561–564.

Munro S, Thomas KL, and Abu-Shaar M (1993) Molecular characterization of a peripheral receptor for cannabinoids. *Nature* **365**:61–65.

Pertwee RG (1972) The ring test: a quantitative method for assessing the “cataleptic” effect of cannabis in mice. *Br J Pharmacol* **46**:753–763.

Rahn EJ, Zvonok AM, Makriyannis A, and Hohmann AG (2010) Antinociceptive effects of racemic AM1241 and its chirally synthesized enantiomers: lack of dependence upon opioid receptor activation. *AAPS J* **12**:147–57.

Staton PC, Hatcher JP, Walker DJ, Morrison AD, Shapland EM, Hughes JP, Chong E, Mander PK, Green PJ, Billinton A, Fulleylove M, Lancaster HC, Smith JC, Bailey LT, Wise A, Brown AJ, Richardson JC, and Chessell IP (2008) The putative cannabinoid receptor GPR55 plays a role in mechanical hyperalgesia associated with inflammatory and neuropathic pain. *Pain* **139**:225–236.

JPET #241901

- Valenzano KJ, Tafesse L, Lee G, Harrison JE, Boulet JM, Gottshall SL, Mark L, Pearson MS, Miller W, Shan S, Rabadi L, Rotshteyn Y, Chaffer SM, Turchin PI, Elsemore DA, Toth M, Koetzner L, and Whiteside GT (2005) Pharmacological and pharmacokinetic characterization of the cannabinoid receptor 2 agonist, GW405833, utilizing rodent models of acute and chronic pain, anxiety, ataxia and catalepsy. *Neuropharmacology* **48**:658–672.
- Viscomi MT, Oddi S, Latini L, Pasquariello N, Florenzano F, Bernardi G, Molinari M, and Maccarrone M (2009) Selective CB2 receptor agonism protects central neurons from remote axotomy-induced apoptosis through the PI3K/Akt pathway. *J Neurosci* **29**:4564–4570.
- Whiteside GT, Gottshall SL, Boulet JM, Chaffer SM, Harrison JE, Pearson MS, Turchin PI, Mark L, Garrison AE, and Valenzano KJ (2005) A role for cannabinoid receptors, but not endogenous opioids, in the antinociceptive activity of the CB2-selective agonist, GW405833. *Eur J Pharmacol* **528**:65–72.
- Yao BB., Mukherjee S., Fan Y., Garrison TR., Daza AV., Grayson GK., Hooker BA., Dart MJ., Sullivan JP, Meyer MD, P.; SJ, Meyer MD, Sullivan JP, and Meyer MD (2009) In vitro pharmacological characterization of AM1241: a protean agonist at the cannabinoid CB2 receptor? *Br J Pharmacol* **149**:145–154.
- Yao BB, Hsieh GC, Frost JM, Fan Y, Garrison TR, Daza a V, Grayson GK, Zhu CZ, Pai M, Chandran P, Salyers a K, Wensink EJ, Honore P, Sullivan JP, Dart MJ, and Meyer MD (2008) In vitro and in vivo characterization of A-796260: a selective cannabinoid CB2 receptor agonist exhibiting analgesic activity in rodent pain models. *Br J Pharmacol* **153**:390–401.
- Zhang J, Hoffert C, Vu HK, Groblewski T, Ahmad S, and O'Donnell D (2003) Induction of CB2 receptor expression in the rat spinal cord of neuropathic but not inflammatory chronic pain models. *Eur J Neurosci* **17**:2750–2754.
- Zimmermann M (1983) Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* **16**:109–10.

JPET #241901

Footnotes:

This work was supported by DA041229 (to AGH and KM), DA009158 (to KM and AGH), and DA021696 (to KM). LMC is supported by T32 NIDA training grant DA024628 and the Harlan Scholars Research Program.

JPET #241901

Figure legends

Figure 1. A. Chemical structure of GW405833, also referred as L768242 (Gallant *et al.*, 1996; Valenzano *et al.*, 2005). Cayman Chemical (Cat. No. 20219) and Tocris Bioscience (Cat. No. 2374) identify this ligand as a selective, high affinity CB₂ receptor partial agonist. **B & C:** Timeline for the dose response of GW405833 and blockade with antagonist in PSNL and CFA model. **B.** In the PSNL model, the baseline was tested a day before the surgery. Animals were allowed (≥ 14 days) to fully develop neuropathic pain prior to pharmacological manipulations. After the full establishment of neuropathic pain, different doses of GW405833 were tested in an ascending order with variable intervals to ensure no carryover effect. Specifically, 0 mg/kg was tested followed by 3 mg/kg on the same day with 3-4 hours interval; 10 mg/kg was then tested the next day followed by 30 mg/kg tested 2 days later. The CB₁ or CB₂ antagonist was tested 3-4 days after the last dose of 30 mg/kg. Groups: C57BL/6J: n= 6 (male); CB₂KO: n = 6 (male); CD1: n = 8 (mixed sex); CB₁KO: n = 8 (mixed sex). **C.** In the CFA model, the baseline was tested on the same day before the CFA injection, the test of dose response started 48 hours after the injection. The timeline for the dose response of GW405833 was the same as the timeline in PSNL model. Groups: C57BL/6J: n = 8 (mixed sex), CB₂KO: n = 8 (mixed sex), CD1: n = 8 (male), CB₁KO: n = 6 (male).

Figure 2. GW405833 dose-dependently reversed PSNL-induced mechanical allodynia in both WT and CB₂KO mice (A). The effect of GW405833 (30 mg/kg i.p.) was only slightly attenuated by pretreatment with the CB₂ antagonist SR144528 (10 mg/kg i.p.). Mechanical thresholds were not changed in the contralateral paw after PSNL or by drug treatments (B). WT (C57BL/6J): n = 6 male; CB₂KO: n = 6 male. # $p < 0.05$ vs. baseline; * $p < 0.05$ vs. vehicle (0 mg/kg i.p.); ^ $p < 0.05$ vs. GW405833 (3 mg/kg i.p.); & $p < 0.05$ vs. GW405833 (30 mg/kg i.p.) in WT. BL: pre-injury baseline. PSNL: partial sciatic nerve ligation. SR2: CB₂ antagonist SR144528. Ipsilateral: injured side; contralateral: uninjured side.

Figure 3. GW405833 dose-dependently reversed PSNL-induced mechanical allodynia in WT, but not in CB₁KO mice (A). The anti-allodynic effect of GW405833 (30 mg/kg i.p.) was completely blocked by the CB₁ antagonist rimonabant (10 mg/kg i.p.) (A). Mechanical threshold was not changed in the contralateral paw after PSNL or by drug treatments (B). Mixed sex groups were used; WT (CD1): n = 8, CB₁KO: n = 8. # $p < 0.05$ vs. baseline; * $p < 0.05$ vs.

JPET #241901

vehicle (0 mg/kg i.p.); $^{\wedge} p < 0.05$ vs. GW405833 (3 mg/kg i.p.); $+ p < 0.05$ vs. CB₁KO, & $p < 0.05$ vs. GW405833 (30 mg/kg i.p.) in WT. BL: pre-injury baseline. PSNL: partial sciatic nerve ligation. SR1: CB₁ antagonist rimonabant. Ipsilateral: injured side; contralateral: uninjured side.

Figure 4. GW405833 dose-dependently reversed CFA-induced mechanical allodynia in both WT and CB₂KO mice (A). The anti-allodynic effect of GW405833 (30 mg/kg i.p.) was unaffected by the CB₂ antagonist SR144528 (10 mg/kg i.p.). Mechanical paw withdrawal thresholds in the contralateral paw were not changed by CFA injection or drug treatments (B). Mixed sex groups were used; WT (C57BL/6J): n = 8, CB₂KO: n = 8. # $p < 0.05$ vs. baseline; * $p < 0.05$ vs. vehicle (0 mg/kg i.p.); $^{\wedge} p < 0.05$ vs. GW405833 (3 mg/kg i.p.) BL: pre-injection baseline. CFA: complete Freund's adjuvant injection. SR2: CB₂ antagonist SR144528. Ipsilateral: injected paw; contralateral: uninjected paw.

Figure 5. GW405833 dose-dependently reversed CFA-induced mechanical allodynia in WT, but not in CB₁KO mice. The anti-allodynic effect of GW405833 (30 mg/kg i.p.) in WT mice was completely blocked by the CB₁ antagonist rimonabant (10 mg/kg i.p.) (B). Mechanical withdrawal thresholds in the contralateral paw were not changed by CFA injection or treatments (B). WT (CD1): n = 8 male, CB₁KO: n = 6 male. # $p < 0.05$ vs. baseline; * $p < 0.05$ vs. vehicle (0 mg/kg i.p.); $^{\wedge} p < 0.05$ vs. GW405833 (3 mg/kg i.p.); \$ $p < 0.05$ vs. GW405833 (10 mg/kg i.p.); $+ p < 0.05$ vs. CB₁KO, & $p < 0.05$ vs. GW405833 (30 mg/kg i.p.) in WT. BL: pre-injection baseline. CFA: complete Freund's adjuvant injection. SR1: CB₁ antagonist rimonabant. Ipsilateral: injected paw; contralateral: uninjected paw.

Figure 6. GW405833 (30 mg/kg i.p.) did not induce catalepsy (A), motor ataxia (B), hypothermia (C) or acute tail-flick antinociception (D) in mice. n = 6 CD1 male mice per group. * $p < 0.05$, *** $p < 0.001$ vs. all the other post injection groups (vehicle and WIN55, 212-2, 5 mg/kg i.p.).

Figure 7. Mice develop behavioral tolerance to Δ^9 -THC administration. Δ^9 -THC (50 mg/kg i.p.) produces motor impairment (A), hypothermia (B) and antinociception (C) on day 1 of repeated dosing relative to baseline and day 8 of Δ^9 -THC administration (A-C), indicating that mice had become tolerant to the pharmacological effects of Δ^9 -THC. Data are expressed as mean \pm SEM (n = 12 C57BL/6J male mice). *** $p < 0.0001$, ** $p < 0.01$ vs. baseline or day 8, #### $p < 0.0001$, # $p < 0.05$ vs. baseline, paired-samples t-test.

JPET #241901

Figure 8. Rimonabant challenge elicits cannabinoid CB₁-dependent withdrawal behaviors whereas GW405833 does not. Rimonabant (10 mg/kg i.p.) challenge increases the number of paw tremors, headshakes, grooming and rearing behaviors in mice chronically-treated with Δ^9 -THC (see tolerance development for these mice in Figure 7). (A). GW405833 (10 mg/kg i.p.) decreases the levels of paw tremors in mice chronically-treated with Δ^9 -THC. (B). Data are expressed as mean \pm SEM (n = 6 C57BL/6J male mice per group). * p<0.05 vs. vehicle, paired-samples t-test.

JPET #241901


Table 1 *In vitro* binding profile of GW405833

Ki (nM)						Assay	Reference
hCB ₁	hCB ₂	hCB ₂ /hCB ₁ selectivity	rCB ₁	rCB ₂	rCB ₂ /rCB ₁ selectivity		
4772	3.92	1217	273	3.6	76	[³ H] CP55,940 binding	Valenzano et al., 2005
282	7.62	37	NA	11.1	NA	[³ H] CP55,940 binding	Yao et al., 2008
1917	12	160	NA	NA	NA	[³ H] WIN55,212-2 binding	Gallant, 1996

NA: not available; hCB: human cannabinoid receptor; rCB: rat cannabinoid receptor.


JPET #241901

Figure 1




JPET #241901

Figure 2




JPET #241901

Figure 3




JPET #241901

Figure 4




JPET #241901

Figure 5




JPET #241901

Figure 6



JPET #241901

Figure 7



JPET #241901

Figure 8

