

Characterization of Isoprenaline- and Salmeterol-Stimulated Interactions between β_2 -Adrenoceptors and β -Arrestin 2 Using β -Galactosidase Complementation in C2C12 Cells

Alison A. Carter and Stephen J. Hill

Institute of Cell Signalling, Medical School, Queens Medical Centre, Nottingham, United Kingdom

Received May 2, 2005; accepted July 26, 2005

ABSTRACT

β -Arrestin is an adaptor protein that has been shown to couple G protein-coupled receptors (GPCRs) to clathrin-coated pits and target them for subsequent internalization. More recently, β -arrestin 2 has also been shown to be involved in the activation of mitogen-activated protein kinase cascades by G protein-coupled receptors independently of G protein activation. Direct interactions between proteins can be monitored using enzyme complementation between two inactive deletion mutants of β -galactosidase (β -gal; $\Delta\alpha$ and $\Delta\omega$). In the present study, we have used fusion proteins of the human β_2 -adrenoceptor (C-terminal β -gal $\Delta\alpha$) and β -arrestin 2 (β -gal $\Delta\omega$) to study directly the pharmacology of this interaction in C2C12 cells expressing the β_2 -adrenoceptor- β -gal $\Delta\alpha$ fusion protein at low physiological levels (38.2 ± 2.7 fmol \cdot mg protein $^{-1}$). Isoprenaline, noradrenaline, and adrenaline ($-\log EC_{50} = 5.9, 5.5,$ and 5.7 , respectively) stimulated an association between the

β_2 -adrenoceptor and β -arrestin 2 at much higher concentrations than required for activation of cAMP accumulation ($-\log EC_{50} = 7.6, 6.3,$ and 7.7 , respectively). This was sensitive to inhibition by the β_2 -adrenoceptor antagonists propranolol, timolol, and ICI 118551. Both salbutamol and terbutaline behaved as partial agonists of β -gal complementation. Furthermore, the long-acting β_2 -agonist salmeterol ($-\log K_D$ for inhibition of [3 H]CGP12177 binding = 8.7) behaved as an antagonist of isoprenaline-stimulated β_2 -adrenoceptor-arrestin 2 interactions ($-\log K_D = 8.0$), whereas acting as a full agonist of cAMP accumulation in the same cells ($-\log EC_{50} = 9.2$). These data suggest that salmeterol can discriminate between receptor- G_s protein and receptor-arrestin 2 complexes (in terms of efficacy and affinity) in a way that is favorable for its long duration of action.

The β_2 -adrenoceptor is the most thoroughly investigated member of the G protein-coupled receptor (GPCR) family, which classically signals through activation of G_s heterotrimeric G proteins (Kobilka, 1992; Benovic, 2002). Agonist stimulation of the β_2 -adrenoceptor leads to activation of adenylyl cyclase, resulting in an increase in the intracellular levels of cAMP (Kobilka, 1992; Benovic, 2002). cAMP activates protein kinase A (PKA) and causes phosphorylation of multiple targets within the cell (Alto et al., 2002). Persistent stimulation of this receptor does not cause a continuous response from the cell, but instead, the cell adapts to the

presence of the stimuli and becomes desensitized (Carman and Benovic, 1998).

Desensitization of the β_2 -adrenoceptor can be mediated either by a mechanism that is dependent upon agonist occupancy of the β_2 -adrenoceptor (homologous desensitization) or by a mechanism that involves the activation of PKA (heterologous desensitization; Carman and Benovic, 1998; Ferguson et al., 1998). However, phosphorylation of the β_2 -adrenoceptor by PKA only causes a partial uncoupling of the receptor from its G protein (Yuan et al., 1994). This phosphorylation can be triggered by low agonist occupancy of the receptor, since it only requires a small increase in cAMP to fully activate PKA (Clark et al., 1999). In contrast, homologous desensitization requires high agonist occupancy by a strong agonist and involves the recruitment of G protein-coupled receptor kinases (GRKs) to the receptor (Kahout and

A.A.C. holds a Biotechnology and Biological Sciences Research Council research studentship.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.
doi:10.1124/jpet.105.088914.

ABBREVIATIONS: GPCR, G protein-coupled receptor; PKA, protein kinase A; GRK, G protein-coupled receptor kinase; MAP, mitogen-activated protein; β -gal, β -galactosidase; CGP 12177, (-)-4-(3-*tert*-utylamino-2-hydroxypropoxy)-benzimidazol-2-one; DMEM, Dulbecco's modified Eagle's medium; DMEM F12, Dulbecco's modified Eagle's medium Ham's F-12; β -AR, β_2 -adrenoceptor; FCS, fetal calf serum; CHO, Chinese hamster ovary; HBH, Hanks' balanced salt solution; PBS, phosphate-buffered saline; ICI 118551, (-)-1-(2,3-dihydro-7-methyl-1*H*-inden-4-yl)oxy)-3-[(1-methylethyl)-amino]-2-butanol; CGP 20712A, 2-hydroxy-5-(2-[[hydroxyl-3-(4-[1-methyl-4-trifluoromethyl-2-imidazolyl]phenoxy)-propyl]amino]ethoxy)benamide.

Lefkowitz, 2003; Marchese et al., 2003). This leads to phosphorylation of serine residues within the C-terminal tail of the β_2 -adrenoceptor (Seibold et al., 2000) and the subsequent binding of β -arrestins (Barak et al., 1997; Ferguson et al., 1998). Binding of arrestin causes a disruption of the receptor- G_s -protein interaction and prevents further signaling from the receptor via the G_s -protein (Barak et al., 1997).

Arrestin has also been found to act as an adaptor molecule coupling GPCRs to clathrin-coated pits and targeting the receptor for subsequent internalization (Luttrell and Lefkowitz, 2002; Marchese et al., 2003). More recently, β -arrestin 2 has also been shown to act as a scaffold for activation of MAP kinase cascades by GPCRs and can function to retain activated MAP kinase enzymes within the cytosol rather than allowing them to translocate to the nucleus (Luttrell et al., 2001; Seta et al., 2002; Tohgo et al., 2002; Wei et al., 2003). Furthermore, it is now clear that this interaction between GPCRs and β -arrestins can activate intracellular signaling via MAP kinase independently of the involvement of heterotrimeric G proteins (Seta et al., 2002; Azzi et al., 2003; Wei et al., 2003; Terrillon and Bouvier, 2004).

The translocation of arrestins to the plasma membrane and their physical interaction with GPCRs therefore represents an important early step in receptor internalization and G protein-independent receptor signaling. However, at the present time, only a few investigations have used technologies capable of detecting direct protein-protein interactions to probe specifically and quantitatively the characteristics of the interaction between GPCRs and β -arrestins. These have included the use of bioluminescence resonance energy transfer (Berglund et al., 2003) and the study of the translocation of fluorescently labeled arrestins to the plasma membrane (Oakley et al., 2002). In both of these examples, complex detection techniques (e.g., confocal microscopy or other ratio-metric optical techniques) and/or algorithms are required to obtain quantitative information on the extent of these protein-protein interactions. In this study, we have used β -galactosidase (β -gal) complementation and a simple luminescence detection methodology to investigate the pharmacology of these interactions.

Direct interactions between different proteins can be monitored by use of enzyme complementation between two inactive deletion mutants of β -galactosidase (β -gal; $\Delta\alpha$ and $\Delta\omega$; Mohler and Blau, 1996). Both mutants of the bacterial enzyme contain inactivating mutations in different essential domains and have been expressed as a fusion protein to target proteins, which are known to associate via protein-protein interactions (Rossi et al., 1997, 2000). When the two target proteins interact the deletion mutants of the galactosidase enzyme are brought into proximity, allowing complementation and the recreation of an active enzyme (Rossi et al., 1997, 2000). The aim of the present study was to investigate the pharmacology of the β_2 -adrenoceptor- β -arrestin 2 interaction at low physiological levels of receptor expression using a cell line containing β -galactosidase mutants fused to either the C terminus of the β_2 -adrenoceptor ($\Delta\alpha$) or the C terminus of β -arrestin 2 ($\Delta\omega$).

Materials and Methods

Materials. Cell culture reagents were from Sigma Chemical (Poole, Dorset, UK), except fetal calf serum (PAA Laboratories,

Teddington, Middlesex, UK). [3 H]Adenine, [3 H]CGP 12177 and [14 C]cAMP were obtained from GE Healthcare (Buckinghamshire, Little Chalfont, UK). Salmeterol, sotalol, CGP 12177, and propranolol were from Tocris Cookson Inc. (Avonmouth, Bristol, UK). The Gal-Screen reagent was from Applied Biosystems (Bedford, MA). All assay plates were obtained from Corning Glassworks (Corning, NY). Lipofectamine was obtained from Invitrogen (Paisley, UK), whereas Opti-MEM reduced serum medium was obtained from Invitrogen cell culture systems. The Hit hunter DiscoverX cAMP assay kit was purchased from DiscoverX (Cardiff, UK). All other reagents were supplied by Sigma Chemical.

Cell Culture. C2C12 (mouse myoblast) cells stably expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins (Yan et al., 2002; Applied Biosystems) were grown at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 4 mM L-glutamine and 20% FCS in a humidified 5% CO₂; 95% air atmosphere. Expression of the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins was maintained using resistance to geneticin (0.5 mg/ml) and hygromycin (0.2 mg/ml), respectively. In addition a cell line expressing only the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein (Yan et al., 2002; Applied Biosystems) was used to investigate the characteristics of the endogenous murine β_2 -adrenoceptor in C2C12 cells. Chinese hamster ovary (CHO) cells were grown at 37°C in Dulbecco's modified Eagle's medium/Nutrient mix F-12 (DMEM F12) supplemented with 2 mM L-glutamine and 10% FCS in a humidified 5% CO₂, 95% air atmosphere.

Transient Transfection of β_2 AR- β -Gal- $\Delta\alpha$ in CHO Cells. CHO cells were grown to 60 to 70% confluence in T75 flasks and then transfected with 10 μ g of β_2 AR- β -gal- $\Delta\alpha$ (Applied Biosystems) in pcDNA3.1 using Lipofectamine and Opti-MEM according to the manufacturer's instructions. Cells were incubated overnight at 37°C, 5% CO₂; after this time the transfection mix was removed and replaced with 20 ml of DMEM F12 media (containing 2 mM glutamine and 10% FCS). Cells were incubated for a further 24 h at 37°C, 5% CO₂, after which time they were plated into 96-half-well white view plates.

[3 H]cAMP Accumulation. Cells were grown to 80% confluence in 24-well plates then prelabeled with [3 H]adenine (2 μ Ci ml⁻¹) for 2 h at 37°C in 500 μ l well⁻¹ Hanks' balanced salt solution containing 20 nM HEPES, pH 7.4 (HBH). The [3 H]adenine was removed, each well washed with 1 ml of HBH, and then incubated for 30 min with 1 ml of medium containing 3-isobutyl-1-methyl xanthine (100 μ M). Agonists were added at 10 μ l per well, and cells were incubated for a further 10 min before the reaction was terminated by the addition of 50 μ l of concentrated HCl. [3 H]Cyclic AMP was separated from other [3 H]adenine nucleotides by acid alumina chromatography. Each column was corrected for efficiency by comparison with [14 C]cAMP recovery as described previously (Alvarez and Daniels, 1992).

cAMP Accumulation Using the DiscoverX Assay. Cells were grown to 80% confluence in 96-half-well white view plates. The media was then aspirated and replaced with 20 μ l well⁻¹ of assay media (C2C12 cells, DMEM media containing 4 mM glutamine; CHO cells, DMEM F12 media containing 2 mM glutamine) containing agonist where appropriate. Cells were incubated for 30 min at 37°C. Cells were then lysed and monitored for cAMP content according to the manufacturer's instructions (DiscoverX).

β_2 -Adrenoceptor- β -Gal- $\Delta\alpha$ and β -Arrestin 2- β -Gal- $\Delta\omega$ Fusion Protein Complementation. Cells were grown to 80% confluence in 96-well white view plates; media were aspirated and replaced with 180 μ l well⁻¹ assay media (DMEM media containing 4 mM glutamine), plus or minus antagonist. Before agonist addition, cells were incubated for 30 min at 37°C with antagonist. Agonists were added at 20 μ l well⁻¹, and cells were incubated for a further 1 h. Reactions were terminated via aspiration of all media. Cells were washed with phosphate-buffered saline (PBS) at 200 μ l well⁻¹, and lysed with the addition of Gal-Screen buffer/substrate (with the substrate diluted 1:25) at 100 μ l well⁻¹. Cells were then incubated at

room temperature in Gal-Screen buffer/substrate (between 21 and 26°C) for 1 h. β -Galactosidase activity was determined using a Top-Count microplate scintillation and luminescence counter.

[³H]CGP 12177 Binding. Cells were grown to confluence in 96-well white view plates. The media were removed, and 200 μ l of serum-free media DMEM containing [³H]CGP 12177 (0.05–8 nM) and β_2 -ligands was added to each well. The plates were incubated for 1.5 h at 37°C in a 5% CO₂ atmosphere. Nonspecific binding was determined using 1 μ M ICI 118551. The media and drugs were then removed, and the cells were washed twice with 200 μ l of PBS. Then, 200 μ l of Microscint 20 was then added to each well, and the plates were counted on a TopCount.

Data Analysis. A maximal isoprenaline concentration was included in each separate experiment for both β -galactosidase complementation, cAMP, and [³H]cAMP accumulation to allow agonist responses to be expressed as a percentage of the isoprenaline maximum. cAMP data were corrected for recovery of tracer [¹⁴C]cyclic AMP, from acid alumina chromatography.

Agonist concentration response curves were fitted to a four-parameter logistic equation through the computer assisted nonlinear regression using the program Prism 2 as described previously (Baker et al., 2003a). Antagonist dissociation curves were assessed at fixed antagonist concentrations (assuming competitive antagonism) by observing the shift in the agonist concentration-response curve using the equation $DR = 1 + [A]/K_D$, where DR (dose ratio) is the ratio of the concentrations of agonist required to produce an identical response in the presence and absence of antagonist, [A] is the concentration of antagonist, and K_D is the antagonist dissociation constant.

Where antagonists produced a decrease in the maximum isoprenaline response, K_D values were obtained as described by Christopoulos et al. (1999). Data were normalized against the isoprenaline standard included in each experiment, and the EC₂₅ was then determined and used to calculate K_D values using the above-mentioned equation. Where appropriate, partial agonist dissociation constants were estimated according to the method of Stephenson (1956). In the case of salmeterol inhibiting the β -galactosidase response to a fixed concentration of isoprenaline (10 μ M), the apparent K_D value was determined from the relationship $C/C' = (IC_{50}/K_D) + 1$, where C is the concentration of isoprenaline (10 μ M) and C' is the concentration of isoprenaline (alone) that produces response equivalent to 50% of that produced by 10 μ M isoprenaline in the presence of an IC₅₀ concentration of salmeterol.

K_D values were also obtained from inhibition of the specific binding of [³H]CGP 12177 using the relationship $K_D = IC_{50}/(1 + [L]/K_L)$, where [L] is the concentration of [³H]CGP 12177 and K_L is its dissociation constant. Saturation binding curves were fitted to the following expression: specific binding (SB) = $B_{MAX} \times [L]/(K_L + [L])$, where B_{MAX} is the maximal specific binding capacity. The curve in Fig. 2 was also fitted to the expression $SB = B_{MAX} \times [L]/(K_L + [L]) + m \times [L]$, where m is the slope of the line for specific binding in native cells only expressing the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein.

All data are represented as mean \pm S.E.M. The "n" in the text refers to the number of separate experiments.

Results

Agonist-Stimulated Interactions between the β_2 -Adrenoceptor and β -Arrestin 2. Isoprenaline produced a marked stimulation of β -galactosidase activity (β_2 -adrenoceptor/ β -arrestin 2 complementation) in C2C12 cells stably expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins but not in native C2C12 cells (Fig. 1a). Similarly, no stimulation of β -galactosidase activity was detected in C2C12 cells expressing only the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein ($n = 4$; data not shown). However, it is interesting to note that in the native C2C12 cells, a considerable degree of endogenous β -galactosidase activity

was detectable (Fig. 1a). The presence of endogenous β -galactosidase activity in mammalian cells has been detected previously (Fisher et al., 1967; Hendriks et al., 1994; Weiss et al., 1997). Similar levels of endogenous β -galactosidase activity were detected in CHO-K1 cells expressing the human β_2 -adrenoceptor; however, stimulation of β_2 -adrenoceptors with isoprenaline did not enhance this β -galactosidase activity (data not shown; $n = 4$), despite a considerable stimulation by isoprenaline of [³H]cAMP accumulation in these cells (Baker et al., 2003a). Furthermore, direct stimulation of cAMP accumulation with forskolin (3 μ M) did not alter β -galactosidase activity in either CHO-K1 cells (data not shown; $n = 4$) or in C2C12 cells (expressing both β -gal fusion proteins; Fig. 1b). These data confirm that the endogenous β -galactosidase activity is not regulated by β_2 -adrenoceptor stimulation or elevation in cAMP levels.

Isoprenaline stimulated cAMP accumulation in both C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins and in native C2C12 cells (Fig. 1c). This latter effect indicates that C2C12 cells express endogenous β_2 -adrenoceptors, but they are also not able to regulate endogenous β -galactosidase activity (Fig. 1a). The log EC₅₀ for isoprenaline for this cAMP response in native C2C12 cells was -5.6 ± 0.0 and the maximum level of cyclic AMP generated was 1240 ± 297 fmol cAMP/well ($n = 3$). In the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ -expressing cells, the log EC₅₀ was an order of magnitude more sensitive (-6.7 ± 0.3) and the maximal response was larger (2387 ± 373 fmol/well; $n = 4$). Transient expression of the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein in CHO-K1 cells that do not endogenously express a β_2 -adrenoceptor (Baker et al., 2003a) confirmed that this receptor fusion protein was able to respond normally to β -agonists and to stimulate cAMP accumulation (Fig. 1d).

In the host C2C12 cells expressing the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein, but not the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, no high-affinity specific binding of [³H]CGP 12177 was detectable (Fig. 2), indicating a very low level of endogenous β_2 -adrenoceptor expression in C2C12 cells. Indeed, the binding that was sensitive to inhibition by 1 μ M ICI 118551 in C2C12 cells expressing the endogenous β_2 -adrenoceptor was best fit to a straight line (Fig. 2). Furthermore, no displacement of this binding of [³H]CGP 12177 (1 nM) by β_2 -agonists was detectable in these cells, suggesting that it is nonspecific ($n = 4$). In C2C12 cells also expressing the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, however, saturation binding analysis confirmed the expression of detectable levels of β_2 -adrenoceptors (38.2 ± 2.7 fmol \cdot mg protein⁻¹; $n = 4$; K_D for [³H]CGP 12177 was 0.2 ± 0.1 nM; $n = 4$; Fig. 2). The β_2 -adrenoceptor-selective antagonist ICI 118551 was able to potently inhibit this specific binding of [³H]CGP 12177 with a $-\log K_D$ of 9.17 ± 0.04 ($n = 4$). Interestingly, the linear low-affinity nonspecific component of ICI 118551-displaceable binding was also detected in the binding curve obtained in cells transfected with the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein (Fig. 2).

β -Galactosidase complementation in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins was also stimulated by adrenaline and noradrenaline, yielding very similar log EC₅₀ values and maximal responses to that obtained with isoprenaline (Table 1). Isoprenaline responses were detectable as early as 5 min




Fig. 1. β_2 -Adrenoceptor/ β -arrestin 2 complementation (β -galactosidase activity) and cAMP accumulation stimulated by isoprenaline in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins. a, β -galactosidase activity [in relative light units (RLU)] obtained in a single experiment after 1-h incubation with agonist in native C2C12 cells or C2C12 cells expressing the β_2 -adrenoceptor and β -arrestin 2 fusion proteins. Similar data were obtained in five (native C2C12 cells) or nine (fusion protein-containing cells) separate experiments. Data points are mean \pm S.E.M. of triplicate determinations. b, β -galactosidase activity in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins in response to isoprenaline and forskolin. Data are the mean values \pm S.E.M. of four separate experiments. In each experiment, determinations were made in triplicate. c, cyclic AMP accumulation in response to isoprenaline in native C2C12 cells or in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins. Values are mean \pm S.E.M. of six replicates in a single experiment. Similar data were obtained in three (C2C12 cells) or four (fusion protein-containing cells) separate experiments. cAMP was measured using the DiscoverX assay as described under *Materials and Methods*. In the cells expressing the β_2 -adrenoceptor and β -arrestin 2 fusion proteins, parallel studies were made of the effect of isoprenaline on the cAMP activity in the absence of the DiscoverX cAMP antibody and cAMP ED reagents. This background galactosidase activity contributed less than 6000 RLU to the overall signal and was not influenced by agonist concentration. d, effect of isoprenaline and salmeterol on cAMP accumulation (DiscoverX) in CHO-K1 cells transiently transfected with the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein. Data are expressed as a percentage of the response to 3 μ M forskolin (measured in each individual experiment). Values are mean \pm S.E.M. of 16 (isoprenaline) and 18 (salmeterol) separate experiments.

(Fig. 3) after agonist stimulation (1.86 ± 0.09 -fold over basal, $-\log EC_{50} = 5.5 \pm 0.2$; $n = 7$) and maintained for up to an hour after addition of isoprenaline (2.1 ± 0.1 -fold over basal, $-\log EC_{50} = 5.7 \pm 0.2$; $n = 7$). All three agonists were also able to elicit comparable stimulations of [3 H]cAMP accumulation in the same cell line, although the EC_{50} values were substantially lower than those obtained for β -galactosidase complementation (Table 2).

To determine the extent to which the agonist-stimulated β -galactosidase complementation was maintained after removal of agonist, cells were stimulated for either 10 or 30 min

(Fig. 4) with isoprenaline. After this period of time, the agonist was washed out, the media were replaced, and the assay continued for a further 50 or 30 min in the absence of agonist. Compared with a standard isoprenaline response measured over 1 h, responses were markedly reduced after washout of isoprenaline (Fig. 4). The decrease in the maximal response to isoprenaline after this treatment was $52 \pm 2\%$ (10 min, $-\log EC_{50} = 5.4 \pm 0.4$; $n = 5$) and $60 \pm 1\%$ (30 min, $-\log EC_{50} = 5.3 \pm 0.3$; $n = 6$) respectively. The residual, but reduced, maximal responses may well reflect internalization of receptor-arrestin complexes, but this requires further investigation.




Fig. 2. Specific binding of [³H]CGP 12177 to C2C12 cells expressing β -arrestin 2- β -gal $\Delta\omega$ -galactosidase and either the endogenous murine β_2 -AR or this endogenous receptor and the human β_2 -adrenoceptor- $\Delta\alpha$ - β -galactosidase fusion protein (β_2 -AR- β -gal $\Delta\alpha$). Specific binding was taken as that displaceable by 1 μ M ICI 118551. Values represent mean \pm S.E.M. from eight (β_2 -AR) or four (β_2 -AR- β -gal $\Delta\alpha$) separate experiments. In each individual experiment, quadruplicate determinations were made of total and nonspecific binding at each [³H]CGP 12177 concentration. The combined endogenous β_2 -AR data were fitted by linear regression (slope, 1.06 \pm 0.04). The mean data from the β_2 -AR- β -gal $\Delta\alpha$ cells were fitted to an equation based on a single binding site with an additional linear component (slope, 1.06). The B_{MAX} and K_D values from this analysis were 38.2 \pm 2.7 fmol \cdot mg protein⁻¹ and 0.09 \pm 0.01 nM, respectively.

TABLE 1

Concentration-response parameters for agonist-stimulated complementation between β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins in C2C12 cells

Values are mean \pm S.E.M. E_{max} is the maximum response expressed as a percentage of the response to 0.1 mM isoprenaline, which was measured in each experiment. The response to salmeterol was only sufficiently large to obtain an EC_{50} value in seven of the eight experiments.

	$-\log EC_{50}$	E_{max} %	n
Isoprenaline	5.9 \pm 0.1	100	8
Adrenaline	5.7 \pm 0.1	118 \pm 8	9
Noradrenaline	5.5 \pm 0.0	92 \pm 5	8
Terbutaline	5.6 \pm 0.1	64 \pm 6	8
Salbutamol	6.3 \pm 0.1	40 \pm 2	9
Salmeterol	7.6 \pm 0.2	14 \pm 3	8

The β_2 -adrenoceptor antagonists timolol and ICI 118551 (Fig. 5a) shifted the isoprenaline concentration-response curve for galactosidase complementation to the right, but also reduced the maximum response. This was consistent with a hemi-equilibrium between slowly dissociating antagonists and an agonist response with a low receptor reserve (Christopoulos et al., 1999; Baker et al., 2003a). Estimation of $-\log K_D$ values according to the method of Christopoulos et al. (1999) produced $-\log K_D$ values of 9.7 \pm 0.2 ($n = 6$) and 8.7 \pm 0.2 ($n = 5$) for timolol and ICI 118551, respectively. In contrast, low concentrations of the β -adrenoceptor antagonist propranolol (Fig. 5b) produced shifts in the concentration-response curve to isoprenaline that were more consistent with competitive antagonism ($-\log K_D = 9.0 \pm 0.1$; $n = 6$) and its more rapid dissociation from the receptor (Motulsky and Mahan, 1984). The lower affinity β -adrenoceptor antagonist sotalol ($-\log K_D = -6.0 \pm 0.1$; $n = 9$; Fig. 5c) also produced a shift that was more compatible with competitive




Fig. 3. Comparison of isoprenaline-induced β_2 -adrenoceptor/ β -arrestin 2 complementation obtained after different times of agonist stimulation in C2C12 cells. Data are expressed as a percentage of the response to 0.1 mM isoprenaline in each experiment and represent mean \pm S.E.M. from seven separate experiments.

antagonism as would be predicted by its more rapid dissociation. CGP 12177 was without any agonist effects, but it proved to be a potent antagonist of the isoprenaline response ($-\log K_D = -9.3 \pm 0.2$; $n = 6$; Fig. 5d).

Partial Agonist-Induced β -Galactosidase Complementation. Partial agonists of the β_2 -adrenoceptor have previously been reported to induce only a small desensitization and internalization of the β_2 -adrenoceptor via GRK phosphorylation and β -arrestin recruitment (January et al., 1997; Clark et al., 1999; Baker et al., 2003b). In these C2C12 cells expressing low levels of the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, terbutaline and salbutamol were able to elicit [³H]cAMP responses of a similar size to those obtained with isoprenaline (Table 2). However, in the case of β_2 -adrenoceptor- β -arrestin 2 galactosidase complementation responses, the maximum responses to these low-efficacy agonists were markedly lower than that obtained with isoprenaline (Table 1; Fig. 6a). Salmeterol is a long-acting β_2 -agonist that has been proposed to interact additionally with an excite on the receptor via which its action is prolonged (Green et al., 1996). This agonist was very weak at producing an effective interaction between the β_2 -adrenoceptor and β -arrestin 2. The maximum response obtained accounted for only 14% of the maximal response to isoprenaline (0.1 mM) measured in the same experiment (Fig. 6b; Table 1). In those experiments in which it could be determined, the mean $-\log EC_{50}$ value for salmeterol was 7.6 \pm 0.2 ($n = 7$). However, salmeterol was able to elicit a full cAMP response after activation of the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ conjugate in the same cells (Table 2). In keeping with its very low efficacy for activation of β_2 -adrenoceptor/ β -arrestin 2 complexes, salmeterol was able to antagonize galactosidase responses to the more efficacious agonist isoprenaline (Fig. 7). Thus, salmeterol was able to shift isoprenaline concentration-response curves to higher agonist concentrations if added simultaneously with isoprenaline (Fig. 7b) or if applied 30 min before isoprenaline (Fig. 7a). Estimation of $-\log K_D$ values according to the method of Stephenson (1956) produced values of 8.3 \pm 0.2 ($n = 7$) and 8.2 \pm 0.3 ($n = 9$) for simultaneous addition or preincubation with salmeterol, respectively. The effect of pretreatment (30 min) with different concentrations of salmeterol on the β -galactosidase complementation response to a fixed concentration (10 μ M) of isoprenaline was

TABLE 2

Concentration-response parameters for agonist-stimulated [³H]cAMP accumulation in C2C12 cellsValues are mean ± S.E.M. E_{max} is the maximum response expressed as a percentage of the response to 0.1 mM isoprenaline, which was measured in each experiment.

Agonist	Human β_2 -Adrenoceptor- β -gal- $\Delta\alpha$ - β -arrestin 2- β -gal- $\Delta\omega$ C2C12 Cells			β -Arrestin 2- β -gal- $\Delta\omega$ C2C12 Cells		
	$-\log EC_{50}$	E_{max} %	n	$-\log EC_{50}$	E_{max} %	n
Isoprenaline	7.6 ± 0.2	100	3	6.9 ± 0.0	100	6
Adrenaline	7.7 ± 0.1	88 ± 5	3	6.5 ± 0.3	96 ± 18	5
Noradrenaline	6.3 ± 0.1	102 ± 1	3	4.5 ± 0.0	73 ± 3	4
Terbutaline	7.3 ± 0.3	89 ± 6	4	5.9 ± 0.1	52 ± 3	4
Salbutamol	7.3 ± 0.1	94 ± 5	3	5.2 ± 0.1	60 ± 6	4
Salmeterol	9.2 ± 0.2	90 ± 5	6	8.2 ± 0.2	42 ± 5	5




Fig. 4. β_2 -Adrenoceptor/ β -arrestin 2 complementation (β -galactosidase activity) after washout (w/o) of isoprenaline in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins. Cells were initially stimulated for 10 (a) or 30 (b) min with isoprenaline, after which time all media were removed, the cells were washed in fresh media, and then incubations were continued for a further 50 or 30 min in the absence of agonist. Control cells were maintained in the presence of isoprenaline for 1 h. Data points are expressed as a percentage of the response to 0.1 mM isoprenaline measured after 1-h continuous stimulation in each experiment. Data points represent mean ± S.E.M. (triplicate determinations in each individual experiment) from six separate experiments.

also tested (Fig. 8). The $-\log IC_{50}$ value for this inhibitory effect was 6.9 ± 0.2 ($n = 21$) and yielded an estimated $-\log K_D$ value of 7.8 ± 0.1 .

[³H]CGP 12177 Whole Cell Binding. To establish the binding affinities of the various agonists studied for the β_2 -

adrenoceptor- β -gal- $\Delta\alpha$ fusion protein in these cells, whole cell binding experiments were performed using [³H]CGP 12177 (which did not detect any endogenous β_2 -adrenoceptors in the C2C12 cell line). The β_2 -adrenoceptor selective antagonist ICI 118551 was able to potently inhibit the specific binding of [³H]CGP 12177 with a $\log -K_D$ of 9.2 ± 0.0 ($n = 4$). As expected, the majority of the agonists used (isoprenaline, salbutamol, and terbutaline) had low binding affinities (Table 3). However, the exception was salmeterol with a $-\log K_D$ of 8.65 ± 0.03 ($n = 4$).

[³H]cyclic AMP Accumulation in Cells Expressing only the β -Arrestin 2- β -Gal- $\Delta\omega$ Fusion Protein. Despite the very low expression level of endogenous murine β_2 -adrenoceptors, these receptors were able to stimulate [³H]cAMP accumulation in cells expressing only the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein (Table 2). A notable feature of these data was that cAMP responses were only elicited by high agonist concentrations and that noradrenaline, salbutamol, terbutaline, and salmeterol were all partial agonists (Table 2). These data were in marked contrast to higher potencies and apparent efficacies obtained with these ligands in cells where the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein was also expressed (Table 2). In the cells expressing only the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein, the maximum response to isoprenaline was $164.6 \pm 14.8\%$ ($n = 4$) of the response to 3 μ M forskolin. In the C2C12 cells also expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, the maximum response to isoprenaline was $210.9 \pm 7.5\%$ ($n = 4$) of the 3 μ M forskolin response. The endogenous receptor was, however, of the β_2 -adrenoceptor subtype since the cAMP responses to isoprenaline was potently antagonized by ICI 118551 ($-\log K_D = 9.6 \pm 0.1$; $n = 4$). In marked contrast, no significant antagonism was obtained with the selective β_1 -adrenoceptor antagonist CGP 20712A (100 nM; $n = 6$).

Discussion

Recruitment of β -arrestin to cell surface β_2 -adrenoceptors is associated with the desensitization and internalization of this receptor (Barak et al., 1997; Ferguson et al., 1998; Kahout and Lefkowitz, 2003). However, only a few investigations have used technologies capable of detecting direct protein-protein interactions to probe specifically and quantitatively the characteristics of the interaction between GPCRs and β -arrestin 2. These have included the use of bioluminescence resonance energy transfer (Berglund et al., 2003) and study of the translocation of fluorescently labeled arrestins to the plasma membrane (Oakley et al., 2002). However, at the present time very little information is available concerning




Fig. 5. β -Galactosidase complementation induced by isoprenaline in the presence and absence of 10 nM, 100 nM, or 1 μ M ICI 118551 (a); 1, 10, or 100 nM propranolol (b); 100 μ M sotalol (c); or 100 nM CGP 12177 (d). Antagonists were added 30 min prior to isoprenaline. Values are expressed as a percentage of the control response to 1 mM isoprenaline (a–c) or 0.1 mM isoprenaline (d) measured in each experiment. Data points are mean \pm S.E.M. from five (a), six (b), three (c), and six (d) separate experiments.

the pharmacological characteristics of receptor-arrestin interactions in terms of both agonist (efficacy and potency) and antagonist (affinity) properties. In this study we have used β -galactosidase complementation and a simple luminescence detection method to investigate the pharmacology of these interactions.

Isoprenaline was able to stimulate a rapid association between the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins expressed in C2C12 cells. However, an unexpected feature of these studies was the high basal galactosidase activity detected in mammalian cells. This was due to endogenous mammalian galactosidase activity since it could be detected in cells (CHO-K1 and C2C12) that had not been transfected with galactosidase-containing constructs. Endogenous galactosidase activity has been detected previously in mammalian cells and tissues (Fisher et

al., 1967; Hendriks et al., 1994; Weiss et al., 1997). However, in CHO cells expressing high levels of the human β_2 -adrenoceptor (Baker et al., 2003a, 2004), no isoprenaline-stimulated increase in this endogenous galactosidase activity was detected. This was also true of the native C2C12 cells that express endogenous murine β_2 -adrenoceptors at low levels, but can nevertheless stimulate a cAMP response. A similar observation was made with C2C12 cells expressing the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein. It was only when the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein was also present that agonist-stimulated β -galactosidase activity was detected.

A similar sized β -galactosidase response to isoprenaline was observed with adrenaline and noradrenaline as agonists. It was notable, however, that the EC_{50} values for β_2 -adrenoceptor-stimulated cAMP accumulation for all three agonists




Fig. 6. β_2 -Adrenoceptor/ β -arrestin 2 complementation (β -galactosidase activity) stimulated by salbutamol (a) or salmeterol (b) in C2C12 cells expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins. Data points are mean \pm S.E.M. (triplicate determinations) from a single experiment and represent nine (a) or eight (b) separate experiments. Bars represent the β -galactosidase complementation from unstimulated cells and from cells stimulated with 0.1 mM isoprenaline (Iso), respectively.

were 2 orders of magnitude lower than those required for β -galactosidase complementation. This reflects greater signal amplification for cAMP accumulation and the need for higher agonist occupancy for receptor-arrestin interactions (January et al., 1997). These data also confirm that the C-terminal modification to the β_2 -adrenoceptor to create the β -galactosidase- $\Delta\alpha$ fusion protein did not interfere with its ability to signal to Gs and adenylyl cyclase. This was also demonstrated by expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein in CHO-K1 cells and generating potent isoprenaline and salmeterol-stimulated cAMP accumulation. It should be noted that the cell surface expression level of the




Fig. 7. β_2 -Adrenoceptor/ β -arrestin 2 complementation induced by isoprenaline in the presence and absence of 1 μ M salmeterol. In a, salmeterol was preincubated for 30 min before addition isoprenaline, whereas in b, the two agents were added simultaneously. Data points are mean \pm S.E.M. (triplicate determinations) from a single experiment and represent seven (a) and nine (b) separate experiments, respectively. The bar represents the β -galactosidase complementation from unstimulated and cells stimulated with 1 mM salmeterol, respectively.

β_2 -adrenoceptor- β -galactosidase fusion protein was low in the transfected C2C12 cells at \approx 38.2 fmol/mg protein and was therefore very similar to the endogenous levels found in native airway smooth muscle cells (70 fmol/mg protein; Green et al., 1995).

In C2C12 cells transfected with the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein, but not the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, the levels of endogenous murine β_2 -adrenoceptors were not detectable by [3 H]CGP 12177 binding. However, a poorly coupled [3 H]cAMP response was measurable that was characterized by submaximal (relative to isoprenaline) agonist responses to noradrenaline, salbutamol, terbutaline, and salmeterol. The expression in these cells of the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion protein, however, produced a leftward shift in the cAMP concentration-response curves to all agonists and increased their maximal responses to the same level as that obtained with isoprenaline.




Fig. 8. β_2 -Adrenoceptor/ β -arrestin 2 complementation induced by isoprenaline in the presence of increasing concentrations of salmeterol. The two ligands were added simultaneously. Data points are mean \pm S.E.M. (triplicate determinations) from a single experiment representative of 21 separate experiments.

TABLE 3

–Log K_D values determined from inhibition of [3 H]CGP 12177 binding to intact C2C12 cells

Values represent mean \pm S.E.M. of –log K_D values obtained from inhibition of the specific binding of [3 H]CGP12177 (1 nM). Nonspecific binding was determined with 1 μ M ICI 118551. *n* gives the number of separate experiments.

Ligand	–log K_D	<i>n</i>
Terbutaline	5.9 \pm 0.1	7
Salbutamol	6.4 \pm 0.1	8
Noradrenaline	5.5 \pm 0.0	8
Isoprenaline	7.0 \pm 0.1	8
Adrenaline	7.2 \pm 0.1	8
Salmeterol	8.7 \pm 0.0	4

The isoprenaline-stimulated β_2 -adrenoceptor/ β -arrestin 2 galactosidase complementation was antagonized effectively by a number of different β_2 -adrenoceptor antagonists. However, although low concentrations of propranolol seemed to produce a parallel rightward shift of the isoprenaline concentration-response curves consistent with competitive antagonism, timolol, ICI 118551, and higher concentrations of propranolol also substantially decreased the maximal agonist responses. It is known that both ICI 118551 and timolol dissociate slowly from the receptor due to their lipophilic nature (Becker and Porzig, 1984; Baker et al., 2003a) and it is likely that this, combined with the low β_2 -adrenoceptor expression and the lack of signal amplification of the receptor-arrestin interactions in these cells, is the reason for the antagonist behavior. ICI 118551 and timolol are thus effectively removing receptor from the agonist accessible pool needed for activation of the arrestin recruitment. Similar behavior of ICI 118551 has been reported for cAMP accumulation (Baker et al., 2003a). In contrast, it is well documented that propranolol dissociates more quickly from the β_2 -adrenoceptor (Motulsky and Mahan, 1984), which explains why this antagonist effect was surmountable at low concentrations. In keeping with this proposal, the low-affinity β -antagonist sotalol (which should also dissociate more rapidly) produced shifts with a better maintained maximal response. Calculation of the antagonist K_D values for ICI 118551 and timolol by the method of Christopoulos et al. (1999), however, provided values in keeping with values obtained previously

for β_2 -adrenoceptor-mediated responses (Baker et al., 2003a).

Low-efficacy agonists of the β_2 -adrenoceptor such as terbutaline, salbutamol, and salmeterol have previously been described as having little effect on both GRK phosphorylation and internalization of the β_2 -adrenoceptor (January et al., 1997). In the present study, the use of a galactosidase complementation assay has enabled the pharmacological characteristics of β_2 -adrenoceptor- β -arrestin 2 interactions induced by low efficacy agonists to be directly compared. At low physiological levels of receptor expression, both salbutamol and terbutaline were able to act as partial agonists of β_2 -adrenoceptor-mediated arrestin complementation. However, the most striking finding was that, at these levels of receptor expression, the long-acting β_2 -agonist salmeterol had very little effect on arrestin complementation but nevertheless was able to act as a potent and full agonist of cAMP-mediated responses. This is despite the similarity of its efficacy to that of both salbutamol and terbutaline for the stimulation of cAMP accumulation via the endogenous β_2 -adrenoceptor in C2C12 cells expressing only the β -arrestin 2- β -gal- $\Delta\omega$ fusion protein (Table 2). Furthermore, in the C2C12 cells also expressing the human β_2 -adrenoceptor, salmeterol was able to act as an antagonist of the arrestin complementation response to more efficacious β_2 -agonists (e.g., isoprenaline). The log dissociation constant for salmeterol obtained from this interaction was ca. –8.0 (average of the mean values obtained in the three different assay formats). This is an order of magnitude higher than the log EC_{50} value for salmeterol-stimulated cAMP accumulation (–9.2) and higher than the log K_D value determined from inhibition of [3 H]CGP 12177 binding (–8.7). These data suggest that salmeterol may be able to distinguish between the receptor conformations required for G_s -protein activation and that required for β -arrestin 2 binding both in terms of efficacy and affinity. The data obtained with [3 H]CGP 12177 binding confirm that salmeterol has a higher affinity (compared with the other agonists measured) for the β_2 -adrenoceptor in intact C2C12 cells. The values obtained from [3 H]CGP 12177 binding probably represent those for the low-affinity agonist resting state (compared with the high-affinity G_s -coupled active state) of the receptor, since the intracellular surfaces of the receptor in intact cells will be exposed to normal physiological levels of GTP.

As mentioned above, low-efficacy agonists such as salmeterol, salbutamol, and terbutaline do not readily phosphorylate the β_2 -adrenoceptor via GRK (January et al., 1997). This may be because partial agonists stabilize receptor conformations that differ in their ability to act as substrates for GRKs (January et al., 1997). It is known that arrestins possess recognition domains that discriminate between active and inactive receptors as well as between phosphorylated and unphosphorylated forms of the receptor (Gurevich and Gurevich, 2004). As a consequence, the reduced receptor phosphorylation by GRK obtained with low-efficacy agonists may significantly contribute to the low potency of these ligands to activate receptor-arrestin association. Alternatively, the nature of the receptor-arrestin interaction may not be entirely dependent upon receptor phosphorylation and the pharmacology of the β_2 -adrenoceptor-arrestin interaction may differ from that of the β_2 -adrenoceptor- G_s protein interaction in a manner similar to that described for agonist

trafficking between different G proteins (Kenakin, 1995). However, the phosphorylation status of the β_2 -adrenoceptor- β -gal- $\Delta\alpha$ fusion is unknown in this study, and it remains to be established whether the extent of receptor phosphorylation by each agonist correlates with their efficacy in stimulating receptor-arrestin interactions. The study of receptor-arrestin interactions using enzyme complementation does, however, provide a means by which some of these important questions can now be directly addressed.

In the present study, the use of a galactosidase complementation assay has also provided clear evidence that the long-acting β_2 -adrenoceptor agonist salmeterol (Green et al., 1996) has a very low efficacy for stimulating β_2 -adrenoceptor- β -arrestin 2 complementation at physiological levels of receptor expression. These properties are clearly beneficial for an agonist that is designed to provide long-acting agonism of G_s -mediated responses but without substantial receptor internalization. Furthermore, these data suggest that salmeterol may be able to discriminate between receptor- G_s protein and receptor-arrestin 2 complexes (in terms of efficacy and affinity) in a way that is favorable for its long duration of action.

Acknowledgments

We thank Applied Biosystems for provision of the C2C12 (mouse myoblast) cells stably expressing the human β_2 -adrenoceptor- β -gal- $\Delta\alpha$ and β -arrestin 2- β -gal- $\Delta\omega$ fusion proteins (InteraX system) and Dr. Melissa Gee (Applied Biosystems) for technical discussion. We also thank GlaxoSmithKline for financial support.

References

- Alto N, Carlisle MJ, Dodge KL, Langeberg LK, and Scott JD (2002) Intracellular targeting of protein kinases and phosphatases. *Diabetes* **51**:S385–S388.
- Alvarez R and Daniels DV (1992) A separation method for the assay of adenylylase, intracellular cyclic AMP, and cyclic-AMP phosphodiesterase using tritium-labeled substrates. *Anal Biochem* **203**:76–82.
- Azzi M, Charest PG, Angers S, Rousseau G, Kohout T, Bouvier M, and Pineyro G (2003) β -Arrestin-mediated activation of MAPK by inverse agonists reveals distinct active conformations for G protein-coupled receptors. *Proc Natl Acad Sci USA* **100**:11406–11411.
- Baker JG, Hall IP, and Hill SJ (2003a) Agonist and inverse agonist actions of “ β -blockers” at the human β_2 -adrenoceptor provide evidence for agonist-directed signalling. *Mol Pharmacol* **64**:1357–1369.
- Baker JG, Hall IP, and Hill SJ (2003b) Influence of agonist efficacy and receptor phosphorylation on antagonist affinity measurements: differences between second messenger and reporter gene responses. *Mol Pharmacol* **64**:679–688.
- Baker JG, Hall IP, and Hill SJ (2004) Temporal characteristics of CRE-mediated gene transcription: requirement for sustained cAMP production. *Mol Pharmacol* **65**:986–998.
- Barak LS, Ferguson SSG, Zhang J, and Caron MG (1997) A β -arrestin/green fluorescent protein biosensor for detecting G protein-coupled receptor activation. *J Biol Chem* **272**:27497–27500.
- Becker C and Porzig H (1984) Recovery of β -adrenoceptors and cyclic AMP response after long term treatment of intact heart cells with β -blockers. *Br J Pharmacol* **82**:745–755.
- Benovic JL (2002) Novel β_2 -adrenergic receptor signaling pathways. *J Allergy Clin Immunol* **110**:S229–S235.
- Berglund MM, Schober DA, Statnick MA, McDonald PH, and Gehlert DR (2003) The use of bioluminescence resonance energy transfer 2 to study neuropeptide Y receptor agonist-induced β -arrestin 2 interaction. *J Pharmacol Exp Ther* **306**:147–156.
- Carman CV and Benovic JL (1998) G-protein-coupled receptors: turn-ons and turn-offs. *Curr Opin Neurobiol* **8**:335–344.
- Christopoulos A, Parsons AM, Lew MJ, and El-Fakahany EE (1999) The assessment of antagonist potency under conditions of transient response kinetics. *Eur J Pharmacol* **382**:217–227.
- Clark RB, Knoll BJ, and Barber R (1999) Partial agonists and G protein-coupled receptor desensitization. *Trend Pharmacol Sci* **20**:279–286.
- Ferguson SSG, Zhang J, Barak LS, and Caron MG (1998) Molecular mechanisms of

- G protein-coupled receptor desensitization and resensitization. *Life Sci* **62**:1561–1565.
- Fisher D, Whitehouse MW, and Kent PW (1967) β -Xylosidase and β -galactosidase activities of mammalian connective tissues and other sources. *Nature (Lond)* **214**:204–205.
- Green SA, Spasoff AP, Coleman RA, Johnson M, and Liggett SB (1996) Sustained activation of a G protein-coupled receptor via “anchored” agonist binding. Molecular localization of the salmeterol exosite within the β_2 -adrenergic receptor. *J Biol Chem* **271**:24029–24035.
- Green SA, Turki J, Bejarano P, Hall IP, and Liggett SB (1995) Influence of β_2 -adrenergic receptor genotypes on signal transduction in human airway smooth muscle cells. *Am J Respir Cell Mol Biol* **13**:25–33.
- Gurevich VV and Gurevich EV (2004) The molecular acrobatics of arrestin activation. *Trends Pharmacol Sci* **25**:105–111.
- Hendriks PJ, Martens ACM, Visser JWM, and Hagenbeek A (1994) Differential suppression of background mammalian lysosomal β -galactosidase increases the detection sensitivity of LacZ-marked leukemic cells. *Anal Biochem* **222**:456–460.
- January B, Seibold A, Whaley R, Hipkin W, Lin D, Schonbrunn A, Barber R, and Clark RB (1997) β_2 -Adrenergic receptor desensitization, internalization and phosphorylation in response to full and partial agonists. *J Biol Chem* **272**:23871–23879.
- Kahout TA and Lefkowitz RJ (2003) Regulation of G-protein-coupled receptor kinases and arrestins during receptor desensitization. *Mol Pharmacol* **63**:9–18.
- Kenakin T (1995) Agonist-receptor efficacy II: agonist trafficking of receptor signals. *Trends Pharmacol Sci* **16**:232–238.
- Kobilka B (1992) Adrenergic mechanisms as models for G-protein-coupled receptors. *Annu Rev Neurosci* **15**:87–114.
- Luttrell LM and Lefkowitz RJ (2002) The role of β -arrestins in the termination and transduction of G-protein-coupled receptor signals. *J Cell Sci* **115**:455–465.
- Luttrell LM, Roudabush FL, Choy EW, Miller WE, Field ME, Pierce KL, and Lefkowitz RJ (2001) Activation and targeting of extracellular-signal-regulated kinases by β -arrestin scaffolds. *Proc Natl Acad Sci USA* **98**:2449–2454.
- Marchese A, Chen C, Kim Y-M, and Benovic JL (2003) The ins and outs of G-protein-coupled receptor trafficking. *Trends Biochem Sci* **28**:369–376.
- Mohler WA and Blau HM (1996) Gene expression and cell fusion analyzed by LacZ complementation in mammalian cells. *Proc Natl Acad Sci USA* **93**:12423–12427.
- Motulsky HJ and Mahan LC (1984) The kinetics of competitive radioligand binding predicted by the law of mass action. *Mol Pharmacol* **25**:1–9.
- Oakley RH, Hudson CC, Cruikshank RD, Meyers DM, Payne REJ, Rhem SM, and Loomis CR (2002) The cellular distribution of fluorescently labeled arrestins provides a robust, sensitive and universal assay for screening G protein-coupled receptors. *Assay Drug Dev Technol* **1**:21–30.
- Rossi F, Charlton CA, and Blau HM (1997) Monitoring protein-protein interactions in intact eukaryotic cells by β -galactosidase complementation. *Proc Natl Acad Sci USA* **94**:8405–8410.
- Rossi FMV, Blakely BT, and Blau HM (2000) Interaction blues: protein interactions monitored in live mammalian cells by β -galactosidase complementation. *Trends Cell Biol* **10**:119–122.
- Seibold A, Williams B, Huang F, Friedman J, Moore RH, Knoll BJ, and Clark RB (2000) Localisation of the sites mediating desensitization of the β_2 -adrenergic receptor by the GRK pathway. *Mol Pharmacol* **58**:1162–1173.
- Seta K, Nanamori M, Modrall JG, Neubig RR, and Sadoshima J (2002) AT1 receptor mutant lacking heterotrimeric G protein coupling activates the Src-Ras-ERK pathway without nuclear translocation of ERKs. *J Biol Chem* **277**:9268–9277.
- Stephenson RP (1956) A modification of receptor theory. *Br J Pharmacol* **11**:5109–5116.
- Terrillon S and Bouvier M (2004) Receptor activity-independent recruitment of β -arrestin 2 reveals specific signalling modes. *EMBO (Eur Mol Biol Organ) J* **23**:3950–3961.
- Tohgo A, Pierce KL, Choy EW, Lefkowitz RJ, and Luttrell (2002) β -Arrestin scaffolding of the ERK cascade enhances cytosolic ERK activity but inhibits ERK-mediated transcription following angiotensin AT1a receptor stimulation. *J Biol Chem* **277**:9429–9436.
- Wei H, Ahn S, Shenoy SK, Karnik SS, Hunyadi L, Luttrell LM, and Lefkowitz RJ (2003) Independent β -arrestin 2 and G-protein-mediated pathways for angiotensin II activation of extracellular signal regulated kinases 1 and 2. *Proc Natl Acad Sci USA* **100**:10782–10787.
- Weiss DJ, Liggitt S, and Clark JG (1997) In situ histochemical detection of β -galactosidase activity in lung: assessment of X-gal reagent in distinguishing LacZ gene expression and endogenous β -galactosidase activity. *Hum Gene Ther* **8**:1545–1554.
- Yan YX, Boldt-Houle DM, Tillotson BP, Gee MAA, D'Eon BJ, Chang XJ, Olesen CEM, and Palmer MAJ (2002) Cell-based high-throughput screening assay system for monitoring G protein-coupled receptor activation using β -galactosidase enzyme complementation technology. *J Biomol Screen* **7**:451–459.
- Yuan N, Friedman J, Whaley BS, and Clark RB (1994) cAMP-dependent protein kinase and protein kinase C consensus site mutations of the β -adrenergic receptor. Effect on desensitization and stimulation of adenylyl cyclase. *J Biol Chem* **269**:23032–23038.

Address correspondence to: Professor Stephen J. Hill, Institute of Cell Signaling, Medical School, Queen's Medical Centre, Nottingham NG7 2UH, UK. E-mail: stephen.hill@nottingham.ac.uk.