

Inhibiting Protein Arginine Deiminases Has Antioxidant Consequences

Erin E. Witalison, Xiangli Cui, Anne B. Hofseth, Venkataraman Subramanian, Corey P. Causey, Paul R. Thompson, and Lorne J. Hofseth

Department of Drug Discovery and Biomedical Sciences, South Carolina College of Pharmacy, University of South Carolina, Columbia, South Carolina (E.E.W., X.C., A.B.H., L.J.H.); Shanxi Medical University, Taiyuan, Shanxi, China (X.C.); Department of Chemistry, The Scripps Research Institute, Scripps Florida, Jupiter, Florida (V.S.); Department of Chemistry, University of North Florida, Jacksonville, Florida (C.P.C.); and Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, Worcester, Massachusetts (P.R.T.)

Received January 7, 2015; accepted January 28, 2015

ABSTRACT

Ulcerative colitis is a dynamic, idiopathic, chronic inflammatory condition that carries a high colon cancer risk. We previously showed that Cl-amidine, a small-molecule inhibitor of the protein arginine deiminases, suppresses colitis in mice. Because colitis is defined as inflammation of the colon associated with infiltration of white blood cells that release free radicals and citrullination is an inflammation-dependent process, we asked whether Cl-amidine has antioxidant properties. Here we show that colitis induced with azoxymethane via intraperitoneal injection + 2% dextran sulfate sodium in the drinking water is suppressed by Cl-amidine (also given in the drinking water).

Inducible nitric oxide synthase, an inflammatory marker, was also downregulated in macrophages by Cl-amidine. Because epithelial cell DNA damage associated with colitis is at least in part a result of an oxidative burst from overactive leukocytes, we tested the hypothesis that Cl-amidine can inhibit leukocyte activation, as well as subsequent target epithelial cell DNA damage in vitro and in vivo. Results are consistent with this hypothesis, and because DNA damage is a pro-carcinogenic mechanism, our data predict that Cl-amidine will not only suppress colitis, but we hypothesize that it may prevent colon cancer associated with colitis.

Introduction

Ulcerative colitis is a heterogeneous, chronic, relapsing inflammatory bowel disease (IBD) that has a significant impact on quality of life. The millions of people who have this disease have an increased colon cancer risk. Despite varying causes (e.g., environmental factors, genetic susceptibility, imbalanced enteric bacteria), the end result is an abnormal immune response with repeated episodes of colonic inflammation. Although not everyone with colitis develops colon cancer, risk increases when disease duration exceeds 10 years, on the order of 0.5–1.0% annually (Itzkowitz and Yio, 2004). Conventional colitis treatments can reduce periods of active disease and help maintain remission, but these treatments often bring marginal results and patients become refractory and develop serious side effects. Hence, we continue to look for less toxic and more efficacious drugs to suppress colitis and prevent colon cancer.

Protein arginine deiminases (PADs) are an enzyme family that converts peptidyl-arginine to peptidyl-citrulline (Arg→Cit) (Jones et al., 2009), a process called "citrullination." Mammals encode 5 isozymes within a single evolutionarily conserved gene

cluster located on human chromosome 1 (1p35–36) (Vossenaar et al., 2003). Mammalian PAD family members (PAD1–4 and 6) are highly related enzymes within and between individual species. PAD-mediated citrullination post-translationally modifies target proteins, which affects their function (Vossenaar et al., 2003). Recently, protein citrullination has received increased attention because of its role in the pathogenesis of various inflammatory conditions, such as rheumatoid arthritis, multiple sclerosis, psoriasis, chronic obstructive pulmonary disease, neurodegenerative diseases and, because of its emerging role in various human and animal cancers (Mohan et al., 2012). Biochemical and genetic evidence suggests that dysregulated PADs also contribute to the onset and progression of colitis and colon cancer. For example, increased PAD levels are observed in colonic inflammatory lesions in Crohn's disease (Struyf et al., 2009). We confirmed that PADs are also overexpressed in the colons of colitis patients and in mouse colitis (Chumanevich et al., 2011). PAD levels have also been reported to be increased in tumors, including colon adenocarcinomas (Chang and Han, 2006; Mohan et al., 2012).

Importantly, we showed that Cl-amidine (Fig. 1), a novel small-molecule PAD inhibitor, delivered by oral gavage, suppresses mouse colitis (Chumanevich et al., 2011). It should be noted that Cl-amidine irreversibly inhibits PADs through the covalent modification of Cys645 in the active site of

This work was supported by the National Institutes of Health National Cancer Institute [Grant 5R01-CA151304].
dx.doi.org/10.1124/jpet.115.222745.

ABBREVIATIONS: AOM, azoxymethane; Cox-2, cyclooxygenase-2; DSS, dextran sulfate sodium; GPx1, glutathione peroxidase 1; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; IBD, inflammatory bowel disease; HED, human equivalent dose; IFN- γ , interferon- γ ; iNOS, inducible nitric oxide synthase; PAD, protein arginine deiminase; SOD1, superoxide dismutase 1.




Fig. 1. Structure of the pan-PAD inhibitor, Cl-amidine.

the enzymes (Knuckley et al., 2010), and because of steric hindrance, can only inhibit activated PADs that have undergone the calcium-dependent conformational change at the active site (Luo et al., 2006).

Because citrullination is inflammation-dependent (Makrygiannakis et al., 2006), PADs cause citrullination, PAD inhibition by Cl-amidine suppresses inflammation, and inflammation is characterized by infiltrating leukocytes releasing free radicals ("oxidative stress") (Kim et al., 2012); it is possible that Cl-amidine also has antioxidant properties. To better understand the mechanisms by which Cl-amidine suppresses colitis and potentially protects against colitis-associated colon cancer, we explored the antioxidant consequences of PAD inhibition as well as its ability to protect against DNA damage in target epithelial cells *in vitro* and *in vivo*.

Materials and Methods

Cl-Amidine. The synthesis of Cl-amidine has been described previously (Luo et al., 2006; Causey and Thompson, 2008).

Cell Culture and Treatment. ANA-1 mouse macrophages or HCT116 human colon cancer cell lines were maintained in Dulbecco's modified Eagle's media (Hyclone, Logan, UT) supplemented with 10% fetal bovine serum (Biofluids, Rockville, MD), penicillin (10 U/ml) and streptomycin (10 μ g/ml; Biofluids). Experiments with Cl-amidine were carried out by preincubating cells with 0–10 μ g/ml (0–25 μ M) Cl-amidine for 12 hours. To activate ANA-1 cells, Cl-amidine was washed off, and cells were exposed to 100 U/ml interferon- γ (IFN- γ ; R&D Systems, Minneapolis, MN).

Coculture Conditions. Coculture experiments were carried out as previously described (Hofseth et al., 2003) with modifications. HCT116 colon cancer cells were seeded at 2.5×10^6 cells per 150 mm culture dish 24 hours before exposure to Cl-amidine (10 μ g/ml) for 12 hours. Cl-amidine was washed off, then HCT116 cells were exposed to IFN- γ -activated or nonactivated ANA-1 cells. ANA-1 cells were added to the actively growing colon cancer cells at a 3:1 ratio (ANA-1: HCT116 cells). The coculture was incubated for 4 hours before harvest. After harvest, HCT116 and ANA-1 cells were separated with MACS mini-column separators and CD45⁺ microbeads. HCT116 cells were examined for DNA damage by Comet analysis. ANA-1 cells were examined for oxidative bursts by chemiluminescence.

Dextran Sulfate Sodium Mouse Model of Colitis. An outline of the colitis mouse model can be found in Fig. 2A. For this, dextran sulfate sodium (DSS; MP Biomedicals, Solon, OH; mol. wt. 36,000–50,000) mouse model, forty 8-week-old C57BL/6 male mice were injected with azoxymethane (AOM; 10 mg/kg *i.p.*) and divided into four groups. AOM was used because this experiment was carried out in parallel with a colon cancer study (using AOM/DSS); thus, we wanted to replicate each animal model as close as possible. Mice in group 1 were given drinking water ad




Fig. 2. (A) Outline of the AOM/DSS mouse model of colitis used in this study. (B) Effects of Cl-amidine on the colon histology score in the acute AOM/DSS colitis model. Six mice from each group described in *Materials and Methods* were euthanized on day 14, and colons were measured and harvested. Then the histology score was determined. Weight was recorded every 48 hours during the 14-day experiment. Values represent the mean \pm S.E. of the mean. Representative H&E-stained colons are shown for each group. *Significant difference from the AOM + DSS group ($P < 0.01$).

libitum throughout the experiment. One week after AOM injection, 2% DSS was added to the drinking water for groups 2, 3, and 4. Groups 3 and 4 were given 0.25 and 0.5 mg/ml Cl-amidine, respectively, in the drinking water when 2% DSS was added to the water. Fresh doses of DSS and Cl-amidine were added to the drinking water every 48 hours, ensuring compound stability throughout the experiment. Colitis induction was evaluated during this study by monitoring body weight changes, bloody and loose stool, and rectal bleeding with daily observations. We determined that colitis was being sufficiently induced in group 2 compared with groups 1, 3, and 4 (data not shown). We did not examine colon cancer as an endpoint for these experiments because we wanted to determine whether Cl-amidine administered in the drinking water at our doses was effective in suppressing DSS-induced colitis before embarking on a lengthy, expensive mouse model of colitis-associated colon cancer.

Cl-amidine (0.5 mg/ml) is the equivalent to approximately 486 mg/day for humans. Our calculation of the human equivalent amount of Cl-amidine consumed by mice uses the body surface area normalization method (Reagan-Shaw et al., 2008), with the following assumptions: a typical mouse drinks 5 ml water daily and weighs 25 g; the average adult human weighs 60 kg. Therefore, 0.5 mg/ml \times 5 ml drinking water/day = 2.5 mg Cl-amidine/day. An average mouse weighs 25 g, therefore 2.5 mg/25 g \times 1000 g/1 kg = 100 mg/kg per day. As discussed by Reagan-Shaw et al. (2008), the human equivalent dose (HED; mg/kg) = animal dose (mg/kg) \times [animal K_m /human K_m]. As such, HED (mg/kg) for mouse = 100 mg/kg \times [3/37] = 8.1 mg/kg. If an average human adult weighs 60 kg, this equates to 8.1 mg/kg \times 60 kg = 486 mg/day for humans. Correspondingly, 0.25 mg/ml is approximately the HED of 243 mg/day.

After 14 days, mice were euthanized. For 4 mice/group, colons were flushed out with $1 \times$ phosphate-buffered saline, opened longitudinally, and then incubated in 10% fetal bovine serum/5 mM EDTA/ Ca^{2+} /Mg $^{2+}$ free phosphate-buffered saline for 15 minutes. Colons were shaken




Fig. 3. iNOS levels are reduced in the colons of mice treated with Cl-amidine. Six mice from each of the indicated groups were euthanized on day 14, and colons were harvested from each animal and stained with iNOS as described in *Materials and Methods*. (A) Immunoreactivity score (IRS) for each group. Values represent the mean \pm S.E. *Significant difference from the AOM + DSS only group ($P < 0.01$). (B) Representative sections of indicated group. Positive staining is brown colored. (400 \times magnification.)

gently for 10 seconds, and the single-cell suspension consisting of epithelial and inflammatory cells was collected in the supernatant. Trypan blue staining revealed >95% viable cells by microscopic observation. Epithelial and inflammatory cells were separated using CD45⁺ magnetic cell sorting technology, according to kit instructions (Miltenyi BioTec, Auburn, CA). Small aliquots of cells from each group were centrifuged, and a dry pellet was frozen at -80°C for Western blot analysis. CD45⁻ epithelial cells were counted, equalized to 1×10^6 cells, centrifuged (1500 rpm, 5 minutes). Pellets were resuspended in freezing media and frozen at -80°C until Comet analysis. CD45⁺ inflammatory cells were counted, equalized to 1×10^6 cells, centrifuged (1500 rpm, 5 minutes), and examined for oxidative bursts by chemiluminescence. For the remaining 6 mice/group, colons were removed and measured. Colons were cut longitudinally, Swiss-rolled, and fixed in 10% buffered formalin overnight and then processed for histopathology/immunohistochemistry by paraffin embedding and sectioning.

Quantification of Inflammation to Examine Effects on Colitis. Paraffin embedded tissues were serially sectioned, and one section from each mouse was stained with H&E. Sections were microscopically examined for histopathologic changes, as we did previously (Jin et al., 2010).

Western Blot Analysis and Antibodies. Western blots were carried out as described previously (Ying et al., 2005). Antibodies used include inducible nitric oxide synthase (iNOS; rabbit polyclonal, diluted 1:1000; Cayman Chemicals, Ann Arbor, MI), cyclooxygenase-2 (Cox-2; rabbit polyclonal, diluted 1:2000; Cayman Chemicals), superoxide dismutase 1 (SOD1; rabbit polyclonal, diluted 1:2000; Abcam,

Cambridge, MA), glutathione peroxidase 1 (GPx1; rabbit polyclonal, diluted 1:500; Abcam), catalase (rabbit polyclonal, diluted 1:500; Abcam), and GAPDH (glyceraldehyde 3-phosphate dehydrogenase; rabbit monoclonal, clone 14C10, diluted 1:1000; Cell Signaling Technology, Danvers, MA).

Immunohistochemical Staining. For immunohistochemical staining, serial sections of mouse colon tissues (processed as described above) were incubated with anti-iNOS (rabbit polyclonal, diluted 1:2000; Cayman Chemicals), catalase (rabbit polyclonal, diluted 1:1000; Abcam), GPx1 (rabbit polyclonal, diluted 1:1000; Abcam), and SOD1 (rabbit polyclonal, diluted 1:500; Abcam). To ensure even staining and reproducibility, sections were incubated by slow rocking overnight in primary antibodies (4°C) using the antibody amplifier (ProHisto, LLC, Columbia, SC). Sections were processed with EnVision+ System-HRP kit according to kit protocols (DakoCytomation, Carpinteria, CA). The positive control was colon cancer tissue and the negative control lacked primary antibody incubation. Immunohistochemistry was quantified as described previously (Poudyal et al., 2012).




Fig. 4. Cl-amidine attenuates the activation of macrophages and protects from DNA damage in target epithelial cells *in vitro*. (A) iNOS and Cox-2 induction after treatment of ANA-1 mouse macrophages with IFN- γ . Numbers below each blot represent the GAPDH-adjusted density of each band, with the control (0 hour, no treatment) being a baseline of 1.0. The observation that for both markers (iNOS and Cox-2) density is lower in unstimulated cells exposed to Cl-amidine (0 hour, +10 $\mu\text{g/ml}$ Cl-amidine, 5th lane) suggests Cl-amidine inhibits basal activity of macrophages. Accordingly, it also inhibits the activation of macrophages. (B) An oxidative burst in ANA-1 mouse macrophages is attenuated by pretreatment with Cl-amidine (10 $\mu\text{g/ml}$). Chemiluminescence was measured as described in *Materials and Methods*. Results were compared with no Cl-amidine control (\pm S.E.). (C) In the presence of an oxidative burst, target epithelial cells (HCT116 colon cancer cells) pretreated with 10 $\mu\text{g/ml}$ Cl-amidine are protected from DNA damage. Results are represented as the mean Comet tail moment \pm S.E., scoring 50 comets/treatment group. Representative images of Comets in each treatment group are shown above each bar graph. *Significant difference from the untreated (No Cl-amidine) macrophages that were cocultured for 4 hours ($P < 0.01$).




Fig. 5. Cl-amidine induces antioxidant enzymes (catalase, GPx1, SOD1) in IFN- γ -stimulated ANA-1 mouse macrophages and in vivo. (A) ANA-1 mouse macrophage cells were pretreated with 0–10 μ g/ml Cl-amidine for 12 hours and then cells were stimulated with IFN- γ for 8 hours. Antioxidant enzymes of interest were suppressed in activated cells pretreated with 10 μ g/ml (lane 4). (B–D) Mice from each of the indicated groups were euthanized on day 14, and colons were harvested from each

Measuring Oxidative Bursts. Oxidative bursts from inflammatory cells were measured by chemiluminescence according to kit directions (World Precision Instruments, Sarasota, FL) and described by our group previously (Jin et al., 2008).

Comet Analysis. A Comet assay was performed according to kit instructions (CometAssay; Trevigen, Gaithersburg, MD). Cells treated with hydrogen peroxide (200 μ M, 20 minutes) were positive controls. Fifty comets/treatment were captured and quantified with the Automated Comet Assay Analysis System (Loats Associates, Inc., Westminster, MD). The percentage of DNA damage is defined as the percentage of DNA in the tail. The percentage of DNA in the tail is defined as integrated tail intensity \times 100/total integrated cell intensity = normalized measure of the % total cell DNA in the tail.

Statistical Analysis. Mean differences between groups were compared by one-way analysis of variance with Scheffe multiple comparison tests. The *P* value chosen for significance in this study was 0.05.

Results

Cl-Amidine Delivered through the Drinking Suppresses DSS-Induced Colitis. We previously showed that Cl-amidine, delivered daily by oral gavage, suppresses DSS-induced colitis in mice (Chumanevich et al., 2011). Because Cl-amidine has a relatively short half-life in mice (37 minutes) (Bicker et al., 2012a), we wanted to determine whether adding Cl-amidine in the drinking water would enable a slower delivery of similar dosages that had worked previously (Chumanevich et al., 2011). In our previous study, mice were dosed with Cl-amidine at 75 mg/kg per day by oral gavage. Here, we estimate that 0.25 and 0.5 mg/ml are equivalent to 50 and 100 mg/kg daily (see calculations in *Materials and Methods*). Figure 2B shows that delivering Cl-amidine through the water effectively suppresses colitis induced by AOM + DSS. The addition of both 0.25 and 0.5 mg/ml are equally effective, with the histology score nearly that of the AOM only control group. Similarly, Cl-amidine increased weight gain compared with AOM + DSS-treated mice. It is important to note that under such conditions Cl-amidine does suppress protein citrullination (Bicker et al., 2012b).

Because mouse colon length shrinks with stress, inflammation, and ulceration, colon lengths were measured upon euthanasia. Compared with the AOM only group (group 1: 8.7 ± 0.15 cm), the length was significantly reduced in the AOM + DSS group (group 2: 7.7 ± 0.27 cm). Mice consuming AOM + DSS + 0.25 mg/ml Cl-amidine (group 3) had a statistically significant increase in colon length (9.0 ± 0.15 cm) compared with the AOM + DSS group. Mice consuming AOM + DSS + 0.5 mg/ml Cl-amidine (group 4) had a statistically similar colon length (8.6 ± 0.26 cm) to that of groups 1 and 3, indicative of healthier, less inflamed colon in groups 1, 3, and 4.

To further verify the in vivo immunosuppressive effects of Cl-amidine on colon inflammation, we probed colon tissues for iNOS by immunohistochemistry. Figure 3 shows representative sections of stained tissues and quantification of iNOS. Overall, iNOS levels were elevated in AOM + DSS-treated

animal and stained with catalase (B), GPx1 (C), and SOD1 (D) as described in *Materials and Methods*. Immunoreactivity scores (IRS) are shown for each group. Values represent the mean \pm S.E. Significance is compared with the AOM + DSS only group: **P* < 0.05; ***P* < 0.01; ****P* < 0.005. Representative sections of each group were taken at 400 \times magnification, and positive staining is brown colored.

mice, with staining appearing mostly in epithelial cells. iNOS staining was statistically significantly reduced in the AOM + DSS + Cl-amidine-treated mice. Such results both confirm and complement our H&E pathology results.

Cl-Amidine Inhibits Inflammatory Cell Activation and Associated DNA Damage in Target Epithelial Cells In Vitro and In Vivo. Mucosal and DNA damage associated with colitis, in part, is a result of an oxidative burst from overactive inflammatory cells (Luhrs et al., 2002; Sartor, 2006). Because we found that preincubating ANA-1 mouse macrophages with Cl-amidine suppressed iNOS and Cox-2 induction by IFN- γ (Fig. 4A), we tested the hypothesis that PAD inhibition by Cl-amidine can inhibit leukocyte activation with release of free radicals through an oxidative burst and resultant epithelial cell DNA damage. Figure 4B shows that Cl-amidine (10 μ g/ml) inhibits oxidative bursts in cultured macrophages. Cells pretreated with Cl-amidine for 12 hours have 60% less oxidative burst capacity than cells not treated with Cl-amidine. This indicates Cl-amidine blunts basal oxidative bursts. One hour after activation with IFN- γ , Cl-amidine-treated cells have 60% of the oxidative burst capacity of untreated cells, indicating that Cl-amidine also protects from an induced oxidative burst. Thereafter, cells begin regaining their oxidative burst capacity, presumably because of the depletion of Cl-amidine.

Because an oxidative burst from macrophages can induce DNA damage in target epithelial cells, we cocultured ANA-1 macrophages with HCT116 colon cancer cells (see *Materials and Methods*). Figure 4C shows a time-dependent increase in DNA damage, as assessed by Comet assay. Cells preincubated with Cl-amidine were significantly protected from DNA damage at 4 hours after the initiation of cocultivation ($P < 0.01$).

We next asked whether Cl-amidine can activate antioxidant enzymes. Therefore, we measured the levels of key antioxidant enzymes, catalase, GPx1, and SOD1, which are shown to be reduced in models of IBD (Cetinkaya et al., 2006; Sakthivel and Guruvayoorappan, 2013; Ren et al., 2014). Figure 5A shows that the level of each enzyme increases maximally when Cl-amidine is dosed at 5 μ g/ml (12 μ M) in ANA-1 murine macrophages. Also, we demonstrated that Cl-amidine treatment upregulates antioxidant enzymes in our mouse model of colitis compared with AOM + DSS-treated mice (Fig. 5, B–D). As expected, the AOM only-treated mice had low basal levels of catalase and GPx1 due to no induction of inflammation. Likewise, the AOM + DSS-treated mice had attenuated levels of catalase, GPx1, and SOD1 in the presence of significant inflammation. To note, the AOM only group did have a higher basal level of SOD1; however, SOD1 levels were restored upon Cl-amidine treatment (Fig. 5D).

To test whether Cl-amidine inhibits an oxidative burst and associated DNA damage in vivo, isolated colon epithelial and inflammatory cells from mice treated with and without Cl-amidine (see *Materials and Methods*). Figure 6A shows that AOM + DSS-treated mice have increased levels of iNOS in CD45⁺ and CD45⁻ cells. iNOS induction is attenuated in mice consuming Cl-amidine. CD45⁺ leukocytes were examined for an oxidative burst by chemiluminescence and we found that Cl-amidine suppresses CD45⁺ cell activity in the colon (Fig. 6B). CD45-colon epithelial cells from the same mice were examined for DNA damage by Comet analysis. Figure 6C shows DNA damage is blunted in mice consuming Cl-amidine. Interestingly, but not surprisingly, there was appreciable DNA damage in the AOM + DSS group. Such damage is attributed to AOM, a




Fig. 6. Cl-amidine attenuates the activation of white blood cells and protects from DNA damage in target epithelial cells in vivo. Mice were injected with AOM (10 mg/kg), then 1 week later they were given either water ad libitum or 2% DSS in the drinking water for 14 days as described in *Materials and Methods* and in Fig. 2. (A) Protein lysates from scraped mucosa of the colon (4 mice per group; lysates were combined) were examined for iNOS and GAPDH (internal control). Mice consuming 2% DSS had activation of only iNOS in both CD45⁺ and CD45⁻ cells. Mice consuming Cl-amidine + 2% DSS had iNOS attenuated in both cell types. (B) After column separation of inflammatory cells from mucosal cells, we examined an oxidative burst of CD45⁺ inflammatory cells (4 mice per group). Mice consuming Cl-amidine + DSS exhibit CD45⁺ inflammatory cells with attenuated activity compared with mice on 2% DSS only. Chemiluminescence was measured as described in *Materials and Methods* and expressed as mean (\pm S.E.) relative light units (RLU) per 1×10^6 cells. *Significant difference from the AOM + DSS-only group ($P < 0.05$). (C) Mucosal epithelial cells were examined for DNA damage by Comet analysis. Results are presented as the mean (\pm S.E.) tail moment from 200 Comets taken from 4 mice per group. *Significant difference from the AOM + DSS-only group ($P < 0.01$). Representative Comets for each group are shown.

carcinogen capable of causing DNA damage in the gastrointestinal tract (Petzold and Swenberg, 1978; Hong et al., 2001). These results indicate that Cl-amidine attenuates inflammatory cell activation and protects from colon epithelial cell damage in vivo. Such results provide mechanistic reasoning for the ability of Cl-amidine to attenuate colitis-associated mucosal damage, and the potential for protection against colon cancer development.

Discussion

Here we have shown that the pan-PAD inhibitor Cl-amidine, given to mice in drinking water and shown by us to block

protein citrullination in mice fed Cl-amidine (Bicker et al., 2012b), suppresses colitis (Fig. 2). Additionally, we have shown that Cl-amidine suppresses an oxidative burst in leukocytes and protects target epithelial cells from DNA damage in vitro (Fig. 4) and in vivo (Fig. 6). Increasing evidence shows that PADs are involved in the citrullination of multiple target proteins. Although there are some overlapping target proteins, each isozyme (PAD1–4 and PAD6) appears to target a unique set of cellular proteins (Knuckley et al., 2007; Darrah et al., 2012). For instance, PAD4 (the most well studied PAD isozyme) is involved in gene regulation and apoptosis, acting as a transcriptional coregulator for p21, p53, p300, CIP1, ELK1, ING4, and nucleophosmin (Li et al., 2008; Yao et al., 2008; Tanikawa et al., 2009; Guo and Fast, 2011; Zhang et al., 2011). Because Cl-amidine is a pan-PAD inhibitor (Knuckley et al., 2010), it is likely that it suppresses the citrullination of many of these PAD target proteins.

Because citrullination is an inflammation-dependent event (Makrygiannakis et al., 2006) and we previously showed (and show here) that Cl-amidine suppresses colon inflammation (Fig. 2) (Chumanevich et al., 2011), it is likely that citrullination of proteins specifically involved in colitis drives inflammation. Accordingly, we have shown here that PAD inhibition by Cl-amidine suppresses reactive oxygen species release by inflammatory cells. This brings up the intriguing possibility that enzymes/proteins directly involved in reactive oxygen species production are affected by citrullination. Such enzymes/proteins include not only TNF- α (a major target for treatment of IBD in humans that has been shown to be citrullinated, resulting in a change in activity; Moelants et al., 2013), but also downstream targets, such as iNOS and Cox-2, both of which were suppressed by Cl-amidine (Figs. 3, 4, and 6). Our finding that catalase, GPx1, and SOD1 are all elevated by Cl-amidine in vitro and in vivo (Fig. 5) is also consistent with the hypothesis that these antioxidant enzymes are mechanistically involved in the crossroads of Cl-amidine and its ability to suppress an oxidative burst. Because Cl-amidine induces p53 (Li et al., 2008; Cui et al., 2013) and p53 drives the expression of antioxidant enzymes (Tan et al., 1999; Hussain et al., 2004; Yoon et al., 2004; Bensaad et al., 2006; Popowich et al., 2010; Kang et al., 2013), potentially p53 repression by PADs (either by direct or upstream citrullination) is suppressing the ability of p53 to activate antioxidant enzymes. This hypothesis, though, would have to be tested.

In summary, we have shown that the pan-PAD inhibitor Cl-amidine through the drinking water is a viable treatment strategy for colitis. Our data reveal that Cl-amidine directly inhibits leukocyte activation and target epithelial DNA damage within the colon. Because of the strong link between chronic DNA damage and increased cancer risk, we are carrying out separate, long-term studies to explore the hypothesis that Cl-amidine protects mice from inflammation-driven colon cancer. Further studies will also explore whether Cl-amidine works upstream of the colon in peripheral blood cells or lymphoid tissues. Indeed, other biologic therapies can cause cellular apoptosis within spleens of treated mice (Fuss et al., 1999). Here, we present the first line of evidence that a pan-PAD small-molecule inhibitor (Cl-amidine) has anti-inflammatory properties in a colitis mouse model with the ability to suppress leukocyte activation and prevent colon epithelial DNA damage both in vitro and in vivo.

Authorship Contributions

Participated in research design: Witalison, Cui, L. J. Hofseth, Subramanian, Thompson.

Conducted experiments: Witalison, Cui, A. B. Hofseth.

Contributed new reagents or analytic tools: Subramanian, Causey, Thompson.

Performed data analysis: Witalison, Cui.

Wrote or contributed to the writing of the manuscript: L. J. Hofseth, Witalison.

References

- Bensaad K, Tsuruta A, Selak MA, Vidal MN, Nakano K, Bartrons R, Gottlieb E, and Vousden KH (2006) TIGAR, a p53-inducible regulator of glycolysis and apoptosis. *Cell* **126**:107–120.
- Bicker KL, Anguish L, Chumanevich AA, Cameron MD, Cui X, Witalison E, Subramanian V, Zhang X, Chumanevich AP, Hofseth LJ, et al. (2012a) D-amino acid based protein arginine deiminase inhibitors: Synthesis, pharmacokinetics, and in cellulo efficacy. *ACS Med Chem Lett* **3**:1081–1085.
- Bicker KL, Subramanian V, Chumanevich AA, Hofseth LJ, and Thompson PR (2012b) Seeing citrulline: development of a phenylglyoxal-based probe to visualize protein citrullination. *J Am Chem Soc* **134**:17015–17018.
- Causey CP and Thompson PR (2008) An improved synthesis of haloacetamidase-based inactivators of protein arginine deiminase 4 (PAD4). *Tetrahedron Lett* **49**:4383–4385.
- Cetinkaya A, Bulbuloglu E, Kantarceken B, Ciralik H, Kurutas EB, Buyukbese MA, and Gumusalan Y (2006) Effects of L-carnitine on oxidant/antioxidant status in acetic acid-induced colitis. *Dig Dis Sci* **51**:488–494.
- Chang X and Han J (2006) Expression of peptidylarginine deiminase type 4 (PAD4) in various tumors. *Mol Carcinog* **45**:183–196.
- Chumanevich AA, Causey CP, Knuckley BA, Jones JE, Poudyal D, Chumanevich AP, Davis T, Matesic LE, Thompson PR, and Hofseth LJ (2011) Suppression of colitis in mice by Cl-amidine: a novel peptidylarginine deiminase inhibitor. *Am J Physiol Gastrointest Liver Physiol* **300**:G929–G938.
- Cui X, Witalison EE, Chumanevich AP, Chumanevich AA, Poudyal D, Subramanian V, Schetter AJ, Harris CC, Thompson PR, and Hofseth LJ (2013) The induction of microRNA-16 in colon cancer cells by protein arginine deiminase inhibition causes a p53-dependent cell cycle arrest. *PLoS ONE* **8**:e53791.
- Darrah E, Rosen A, Giles JT, and Andrade F (2012) Peptidylarginine deiminase 2, 3 and 4 have distinct specificities against cellular substrates: novel insights into autoantigen selection in rheumatoid arthritis. *Ann Rheum Dis* **71**:92–98.
- Fuss IJ, Marth T, Neurath MF, Pearlstein GR, Jain A, and Strober W (1999) Anti-interleukin 12 treatment regulates apoptosis of Th1 T cells in experimental colitis in mice. *Gastroenterology* **117**:1078–1088.
- Guo Q and Fast W (2011) Citrullination of inhibitor of growth 4 (ING4) by peptidylarginine deiminase 4 (PAD4) disrupts the interaction between ING4 and p53. *J Biol Chem* **286**:17069–17078.
- Hofseth LJ, Hussain SP, Wogan GN, and Harris CC (2003) Nitric oxide in cancer and chemoprevention. *Free Radic Biol Med* **34**:955–968.
- Hong MY, Chapkin RS, Morris JS, Wang N, Carroll RJ, Turner ND, Chang WC, Davidson LA, and Lupton JR (2001) Anatomical site-specific response to DNA damage is related to later tumor development in the rat azoxymethane colon carcinogenesis model. *Carcinogenesis* **22**:1831–1835.
- Hussain SP, Amstad P, He P, Robles A, Lupold S, Kaneko I, Ichimiya M, Sengupta S, Mechanic L, Okamura S, et al. (2004) p53-induced up-regulation of MnSOD and GPx but not catalase increases oxidative stress and apoptosis. *Cancer Res* **64**:2350–2356.
- Itzkowitz SH and Yio X (2004) Inflammation and cancer IV. Colorectal cancer in inflammatory bowel disease: the role of inflammation. *Am J Physiol Gastrointest Liver Physiol* **287**:G7–G17.
- Jin Y, Hofseth AB, Cui X, Windust AJ, Poudyal D, Chumanevich AA, Matesic LE, Singh NP, Nagarkatti M, Nagarkatti PS, et al. (2010) American ginseng suppresses colitis through p53-mediated apoptosis of inflammatory cells. *Cancer Prev Res (Phila)* **3**:339–347.
- Jin Y, Kotakadi VS, Ying L, Hofseth AB, Cui X, Wood PA, Windust A, Matesic LE, Pena EA, Chiuhan C, et al. (2008) American ginseng suppresses inflammation and DNA damage associated with mouse colitis. *Carcinogenesis* **29**:2351–2359.
- Jones JE, Causey CP, Knuckley B, Slack-Noyes JL, and Thompson PR (2009) Protein arginine deiminase 4 (PAD4): Current understanding and future therapeutic potential. *Curr Opin Drug Discov Devel* **12**:616–627.
- Kang MY, Kim HB, Piao C, Lee KH, Hyun JW, Chang IY, and You HJ (2013) The critical role of catalase in prooxidant and antioxidant function of p53. *Cell Death Differ* **20**:117–129.
- Kim YJ, Kim EH, and Hahm KB (2012) Oxidative stress in inflammation-based gastrointestinal tract diseases: challenges and opportunities. *J Gastroenterol Hepatol* **27**:1004–1010.
- Knuckley B, Bhatia M, and Thompson PR (2007) Protein arginine deiminase 4: evidence for a reverse protonation mechanism. *Biochemistry* **46**:6578–6587.
- Knuckley B, Causey CP, Jones JE, Bhatia M, Dreyton CJ, Osborne TC, Takahara H, and Thompson PR (2010) Substrate specificity and kinetic studies of PADs 1, 3, and 4 identify potent and selective inhibitors of protein arginine deiminase 3. *Biochemistry* **49**:4852–4863.
- Li P, Yao H, Zhang Z, Li M, Luo Y, Thompson PR, Gilmour DS, and Wang Y (2008) Regulation of p53 target gene expression by peptidylarginine deiminase 4. *Mol Cell Biol* **28**:4745–4758.
- Lührs H, Gerke T, Müller JG, Melcher R, Schaubert J, Boxberger F, Scheppach W, and Menzel T (2002) Butyrate inhibits NF- κ B activation in lamina propria macrophages of patients with ulcerative colitis. *Scand J Gastroenterol* **37**:458–466.

- Luo Y, Arita K, Bhatia M, Knuckley B, Lee YH, Stallcup MR, Sato M, and Thompson PR (2006) Inhibitors and inactivators of protein arginine deiminase 4: functional and structural characterization. *Biochemistry* **45**:11727–11736.
- Makrygiannakis D, af Klint E, Lundberg IE, Löfberg R, Ulfgrén AK, Klareskog L, and Catrina AI (2006) Citrullination is an inflammation-dependent process. *Ann Rheum Dis* **65**:1219–1222.
- Moelants EA, Mortier A, Grauwen K, Ronsse I, Van Damme J, and Proost P (2013) Citrullination of TNF- α by peptidylarginine deiminases reduces its capacity to stimulate the production of inflammatory chemokines. *Cytokine* **61**:161–167.
- Mohan S, Cherrington BD, Horibata S, McElwee JL, Thompson PR, and Coonrod SA (2012) Potential role of peptidylarginine deiminase enzymes and protein citrullination in cancer pathogenesis. *Biochem Res Int* **2012**:895343.
- Petzold GL and Swenberg JA (1978) Detection of DNA damage induced in vivo following exposure of rats to carcinogens. *Cancer Res* **38**:1589–1594.
- Popowich DA, Vavra AK, Walsh CP, Bhikapurwala HA, Rossi NB, Jiang Q, Aalami OO, and Kibbe MR (2010) Regulation of reactive oxygen species by p53: implications for nitric oxide-mediated apoptosis. *Am J Physiol Heart Circ Physiol* **298**:H2192–H2200.
- Poudyal D, Le PM, Davis T, Hofseth AB, Chumanevich A, Chumanevich AA, Wargovich MJ, Nagarkatti M, Nagarkatti PS, Windust A, et al. (2012) A hexane fraction of American ginseng suppresses mouse colitis and associated colon cancer: anti-inflammatory and proapoptotic mechanisms. *Cancer Prev Res (Phila)* **5**:685–696.
- Reagan-Shaw S, Nihal M, and Ahmad N (2008) Dose translation from animal to human studies revisited. *FASEB J* **22**:659–661.
- Ren W, Yin J, Wu M, Liu G, Yang G, Xion Y, Su D, Wu L, Li T, Chen S, et al. (2014) Serum amino acids profile and the beneficial effects of L-arginine or L-glutamine supplementation in dextran sulfate sodium colitis. *PLoS ONE* **9**:e88335.
- Sakthivel KM and Guruvayoorappan C (2013) Amentoflavone inhibits iNOS, COX-2 expression and modulates cytokine profile, NF- κ B signal transduction pathways in rats with ulcerative colitis. *Int Immunopharmacol* **17**:907–916.
- Sartor RB (2006) Mechanisms of disease: pathogenesis of Crohn's disease and ulcerative colitis. *Nat Clin Pract Gastroenterol Hepatol* **3**:390–407.
- Struyf S, Noppen S, Loos T, Mortier A, Gouwy M, Verbeke H, Huskens D, Luangsay S, Parmentier M, Geboes K, et al. (2009) Citrullination of CXCL12 differentially reduces CXCR4 and CXCR7 binding with loss of inflammatory and anti-HIV-1 activity via CXCR4. *J Immunol* **182**:666–674.
- Tan M, Li S, Swaroop M, Guan K, Oberley LW, and Sun Y (1999) Transcriptional activation of the human glutathione peroxidase promoter by p53. *J Biol Chem* **274**:12061–12066.
- Tanikawa C, Ueda K, Nakagawa H, Yoshida N, Nakamura Y, and Matsuda K (2009) Regulation of protein Citrullination through p53/PADI4 network in DNA damage response. *Cancer Res* **69**:8761–8769.
- Vossenaar ER, Zendman AJ, van Venrooij WJ, and Pruijn GJ (2003) PAD, a growing family of citrullinating enzymes: genes, features and involvement in disease. *BioEssays* **25**:1106–1118.
- Yao H, Li P, Venters BJ, Zheng S, Thompson PR, Pugh BF, and Wang Y (2008) Histone Arg modifications and p53 regulate the expression of OKL38, a mediator of apoptosis. *J Biol Chem* **283**:20060–20068.
- Ying L, Marino J, Hussain SP, Khan MA, You S, Hofseth AB, Trivers GE, Dixon DA, Harris CC, and Hofseth LJ (2005) Chronic inflammation promotes retinoblastoma protein hyperphosphorylation and E2F1 activation. *Cancer Res* **65**:9132–9136.
- Yoon KA, Nakamura Y, and Arakawa H (2004) Identification of ALDH4 as a p53-inducible gene and its protective role in cellular stresses. *J Hum Genet* **49**:134–140.
- Zhang X, Gamble MJ, Stadler S, Cherrington BD, Causey CP, Thompson PR, Roberson MS, Kraus WL, and Coonrod SA (2011) Genome-wide analysis reveals PADI4 cooperates with Elk-1 to activate c-Fos expression in breast cancer cells. *PLoS Genet* **7**:e1002112.

Address correspondence to: Dr. Lorne J. Hofseth, Department of Drug Discovery and Biomedical Sciences, South Carolina College of Pharmacy, 770 Sumter St., Coker Life Sciences, Rm. 513C, University of South Carolina, Columbia, SC 29208. E-mail: hofseth@cop.sc.edu
