

Investigation of Diacylglycerol Lipase Alpha Inhibition in the Mouse Lipopolysaccharide Inflammatory Pain Model^[S]

Jenny L. Wilkerson, Giulia Donvito, Travis W. Grim, Rehab A. Abdullah, Daisuke Ogasawara, Benjamin F. Cravatt, and Aron H. Lichtman

Department of Pharmacology and Toxicology, Virginia Commonwealth University, Richmond, Virginia (J.L.W., G.D., T.W.G., R.A.A., A.H.L.); and The Skaggs Institute for Chemical Biology and Department of Chemical Physiology, The Scripps Research Institute, La Jolla, California (D.O., B.F.C.)

Received July 5, 2017; accepted September 20, 2017

ABSTRACT

Diacylglycerol lipase (DAGL) α and β , the major biosynthetic enzymes of the endogenous cannabinoid (endocannabinoid) 2-arachidonoylglycerol (2-AG), are highly expressed in the nervous system and immune system, respectively. Genetic deletion or pharmacological inhibition of DAGL- β protects against lipopolysaccharide (LPS)-induced inflammatory responses in mouse peritoneal macrophages and reverses LPS-induced allodynia in mice. To gain insight into the contribution of DAGL- α in LPS-induced allodynia, we tested global knockout mice as well as DO34, a dual DAGL- α/β inhibitor. Intraperitoneal administration of DO34 (30 mg/kg) significantly decreased whole-brain levels of 2-AG (~83%), anandamide (~42%), and arachidonic acid (~58%). DO34 dose-dependently reversed mechanical and cold allodynia, and these antinociceptive effects did not undergo tolerance after 6 days of repeated

administration. In contrast, DO34 lacked acute thermal antinociceptive, motor, and hypothermic pharmacological effects in naive mice. As previously reported, DAGL- β (-/-) mice displayed a protective phenotype from LPS-induced allodynia. However, DAGL- α (-/-) mice showed full allodynic responses, similar to their wild-type littermates. Interestingly, DO34 (30 mg/kg) fully reversed LPS-induced allodynia in DAGL- α (+/+) and (-/-) mice, but did not affect the antinociceptive phenotype of DAGL- β (-/-) mice in this model, indicating a DAGL- α -independent site of action. These findings suggest that DAGL- α and DAGL- β play distinct roles in LPS-induced nociception. Whereas DAGL- α appears to be dispensable for the development and expression of LPS-induced nociception, DAGL- β inhibition represents a promising strategy to treat inflammatory pain.

Introduction

Diacylglycerol lipase (DAGL)- α and DAGL- β (Bisogno et al., 2003; Gao et al., 2010; Tanimura et al., 2010) transform diacylglycerols into 2-arachidonoylglycerol (2-AG), the most highly expressed endocannabinoid in the central nervous system (Mechoulam et al., 1995; Sugiura et al., 1995). 2-AG plays critical roles in maintaining proper neuronal function (Goncalves et al., 2008; Tanimura et al., 2010), mediating neuronal axonal growth (Williams et al., 2003) and retrograde suppression of synaptic transmission (Kreitzer and Regehr, 2001; Ohno-Shosaku et al., 2001; Wilson and Nicoll, 2001; Pan et al., 2009). These enzymes are differentially expressed within cells in the nervous system and peripheral tissue (Hsu et al., 2012). DAGL- α is expressed on postsynaptic neurons within various brain regions (Katona et al., 2006; Yoshida et al., 2006; Lafourcade et al., 2007; Uchigashima et al., 2007), and its genetic deletion results in marked decreases of 2-AG, anandamide (AEA), and arachidonic acid (AA) in the

brain (Gao et al., 2010; Tanimura et al., 2010; Shonesy et al., 2014) and spinal cord (Gao et al., 2010). Accordingly, DAGL- α (-/-) mice display impaired depolarization-induced suppression of inhibition and excitation in the brain (Gao et al., 2010; Tanimura et al., 2010; Yoshino et al., 2011). These mice also show an increased mortality rate (Sugaya et al., 2016), display increased spontaneous seizures in the kainate model of status epilepticus (Sugaya et al., 2016), and exhibit an anxiogenic phenotype (Shonesy et al., 2014). In contrast, DAGL- β is most highly expressed on macrophages, and, although its relative brain expression is sparse, it is highly expressed on microglia (Hsu et al., 2012). This distribution pattern suggests that DAGL- β activity contributes to inflammatory responses. Importantly, DAGL- β deletion does not affect endocannabinoid-mediated forms of retrograde synaptic suppression (Gao et al., 2010). However, DAGL- β blockade reduces lipopolysaccharide (LPS)-induced inflammatory responses in peritoneal macrophages from C57BL/6 mice by decreasing levels of 2-AG, AA, prostanoids, and proinflammatory cytokines (Hsu et al., 2012). Similarly, DAGL- α inhibition leads to protection from the neuroinflammatory effects of 20 mg/kg systemic LPS (Ogasawara et al., 2016).

A wide scope of evidence supports inflammatory as well as neuronal signaling contributions to many forms of pathologic pain. Immune cell signaling plays a critical role in the

Research was supported by the following grants from the National Institutes of Health: [DA009789], [DA017259], [DA032933], [DA033934-01A1], [DA035864], and [DA038493-01A1].

<https://doi.org/10.1124/jpet.117.243808>

[S] This article has supplemental material available at jpet.aspetjournals.org.

ABBREVIATIONS: AA, arachidonic acid; ABHD6, alpha beta hydrolase domain-containing protein 6; AEA, anandamide; 2-AG, 2-arachidonoylglycerol; ANOVA, analysis of variance; CI, confidence interval; DAGL, diacylglycerol lipase; LPS, lipopolysaccharide.

development and maintenance of neuropathic pain (Watkins et al., 2001; De Leo et al., 2006; Beggs and Salter, 2013). Likewise, increased neuronal signaling underlies pathologic inflammatory pain and can contribute to a positive pain feedback loop (De Leo et al., 2006; Chen et al., 2015). For example, in LPS-stimulated neurons, neuronal signaling leads to further inflammatory signaling and immune cell activation (Chen et al., 2015). Determining the antecedents of pathologic pain and the subsequent identification of potential therapeutic targets remain important areas of research. Accordingly, DAGL- α and DAGL- β represent provocative targets to treat pathologic pain conditions.

Complementary approaches of pharmacological agents and genetically modified mice demonstrate that DAGL- β blockade reduces nociceptive behavior in the LPS model of inflammatory pain (Wilkerson et al., 2016). The DAGL- β inhibitor KT109 reverses nociceptive behavior in models of neuropathic pain (Wilkerson et al., 2016). These findings strongly implicate inhibition of this enzyme as a viable approach to treat inflammatory and neuropathic pain. However, it remains to be determined whether DAGL- α inhibition or deletion produces antinociceptive effects in pathologic pain models. The present study attempted to investigate the role of this enzyme in LPS-induced allodynia, using the DAGL inhibitor DO34, which disrupts depolarization-induced suppression of excitation and depolarization-induced suppression of inhibition in the cerebellum and hippocampus and reduces LPS-induced anapyrexia in vitro responses (Ogasawara et al., 2016), providing a useful tool for in vitro and in vivo studies. Thus, the present study examined DAGL- α ($-/-$) and DAGL- β ($-/-$) mice in the LPS model of inflammatory pain.

In initial experiments, we quantified brain levels of endogenous cannabinoids and AA in mice administered vehicle or DO34 (30 mg/kg) and tested DO34 in assays of locomotor behavior, catalepsy, body temperature, and acute thermal antinociceptive responses. We then evaluated the dose-response relationship and time course of acute DO34 administration in attenuating LPS-induced mechanical and cold allodynia. In addition, we examined whether the antiallodynic effects would undergo tolerance after repeated DO34 administration. Finally, we tested DO34 in DAGL- α ($-/-$) and DAGL- β ($-/-$) mice, and respective wild-type littermates in the LPS model of inflammatory pain. Because DO34 also inhibits the serine hydrolase ABHD6 (Ogasawara et al., 2016), a 2-AG hydrolytic enzyme expressed on postsynaptic neurons (Blankman et al., 2007; Marrs et al., 2010), we used the selective ABHD6 inhibitor DO53, which lacks DAGL activity (Ogasawara et al., 2016), for comparison.

Materials and Methods

Animals. Adult male C57BL/6J and ICR mice (23–40 g; The Jackson Laboratory, Bar Harbor, ME) served as subjects in these experiments. DAGL- α ($-/-$) and DAGL- β ($-/-$) mice were generated in the Cravatt laboratory on a mixed C57BL/6J and 129/SvEv background, as previously described (Hsu et al., 2012), and breeding pairs were transferred to Virginia Commonwealth University. A total of 84 DAGL- α ($-/-$) and 36 DAGL- β ($-/-$) mice were used in these studies. Mice were housed four per cage in a temperature- (20–22°C), humidity- (55% \pm 10%), and light-controlled (12-hour light/dark; lights on at 6:00 AM) Association for Assessment and Accreditation of Laboratory Animal Care International

(Frederick, MD)-approved facility, with standard rodent chow and water available ad libitum.

All tests were conducted during the light phase. The sample size selected for each treatment group in each experiment was based on previous studies from our laboratory and complied with power analyses.

All animal protocols were approved by the Institutional Animal Care and Use Committee at Virginia Commonwealth University and were carried out in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals (National Research Council, 2011). After testing was completed, mice were euthanized via CO₂ asphyxia, followed by rapid cervical dislocation.

Drugs. The DAGL inhibitor DO34 and the selective ABHD6 inhibitor DO53 were synthesized by the Cravatt laboratory according to previous methods (Ogasawara et al., 2016). All drugs were dissolved in a vehicle solution consisting of a mixture of ethanol, alkamuls-620 (Sanofi-Aventis, Bridgewater, NJ), and saline (0.9% NaCl) in a 1:1:18 ratio. All drugs were administered in an injection volume of 10 μ l/g body mass. Each drug was given via the intraperitoneal route of administration. The dose range of DO34 was selected based on results reported by Ogasawara et al. (2016), indicating that acute administration of 30 mg/kg DO34 in mice treated with LPS was sufficient to produce inhibition of DAGL- α , as well as measured decreases in peritoneal macrophages of AA and proinflammatory cytokines.

Extraction and Quantification of Endocannabinoids by Liquid Chromatography-Tandem Mass Spectrometry. 2-AG, AA, and AEA levels were quantified from the whole brain of ICR mice, after acute administration of DO34 (30 mg/kg, i.p.) or 1:1:18 vehicle. Brains were collected and processed for quantification of 2-AG, AA, and AEA. Because equivalent doses of DO34 significantly attenuated allodynia associated with LPS at 2 hours after injection, mice were euthanized via rapid decapitation at this time point. Brains were rapidly harvested, snap frozen in dry ice, and stored at -80°C until the time of processing. Tissues were further processed according to methods described previously (Ignatowska-Jankowska et al., 2015). See Supplemental Material for details.

Evaluation of Acute Pharmacological Effects of DO34. Mice (counterbalanced Latin square within subject design) were housed individually overnight. The behavioral testing was conducted in the following order: bar test (catalepsy), tail withdrawal test, rectal temperature, and locomotor activity. Testing was performed according to previously described procedures (Long et al., 2009; Schlosburg et al., 2010). Catalepsy was assessed on a bar 0.7 cm in diameter placed 4.5 cm off of the ground. The mouse was placed with its front paws on the bar and a timer (Timer #1) was started. A second timer (Timer #2) was turned on only when the mouse was immobile on the bar, with the exception of respiratory movements. If the mouse moved off the bar, it was placed back on in the original position. The assay was stopped either when Timer #1 reached 60 seconds or after the fourth time the mouse moved off the bar, and the cataleptic time was scored as the amount of time on Timer #2. Nociception was then assessed in the tail immersion assay. The mouse was placed head first into a small bag fabricated from absorbent underpads (4 cm diameter, 11 cm length; VWR Scientific Products, Radnor, PA) with the tail out of the bag. Each mouse was hand held, and 1 cm of the tail was submerged in a 52°C water bath. The latency for the mouse to withdraw its tail within a 10-second cutoff time was scored. Rectal temperature was assessed by inserting a thermocouple probe 2 cm into the rectum, and temperature was determined by thermometer (BAT-10 Multipurpose Thermometer; Protech International Inc., Boerne, TX). Locomotor activity was assessed 120 minutes after treatment for a 60-minute period in a Plexiglas cage (42.7 \times 21.0 \times 20.4 cm) and Anymaze software (Stoelting, Wood Dale, IL) was used to determine the percentage of time spent immobile, mean speed, and distance traveled.

LPS Inflammatory Pain Model. Mice were given an injection of 2.5 μg of LPS from *Escherichia coli* 026:B6 (Sigma-Aldrich, St. Louis, MO) in 20 μl of physiologic sterile saline (Hospira Inc., Lake Forest, IL) into the plantar surface of the right hind paw. As previously reported, this is the minimally effective dose of LPS that elicits mechanical allodynia, but not measurable increases in paw thickness (Booker et al., 2012). After LPS administration, mice were returned to their home cages. At 22 hours, mice were given the appropriate injection of drug or vehicle and tested at 24 hours for allodynia. In the time course study, allodynia was assessed at 40 minutes and 1, 3, 5, 8, and 24 hours after the intraperitoneal injection.

To determine whether repeated administration of DO34 would produce sustained antinociceptive effects, mice were given injections of vehicle or DO34 (30 mg/kg, i.p.) once a day for 5 days. On day 5, each mouse received its appropriate intraperitoneal injection of vehicle or DO34, and 2 hours later all mice were given an intraplantar injection of LPS. On day 6 (22 hours after LPS administration), each mouse received its final intraperitoneal injection. The vehicle-treated mice were divided into two groups. The first group received another injection of vehicle (vehicle control group), and the second group was given 30 mg/kg DO34 (the acute DO34 group). The mice that had been given repeated injections of drug received their final injection of DO34 (the repeated DO34 group). All mice were tested for mechanical and cold allodynia 2 hours after the final intraperitoneal injection.

Behavioral Assessment of Nociception. Baseline responses to light mechanical touch were assessed using the von Frey test after habituation to the testing environment, as described previously (Murphy et al., 1999). In brief, mice were placed atop a wire mesh screen, with spaces 0.5 mm apart and habituated for approximately 30 min/day for 4 days. Mice were unrestrained, and were singly placed under an inverted wire mesh basket to allow for unrestricted airflow. The von Frey test utilizes a series of calibrated monofilaments (2.83–4.31 log stimulus intensity; North Coast Medical, Morgan Hills, CA) applied randomly to the left and right plantar surface of the hind paw for 3 seconds. Lifting, licking, or shaking the paw was considered a response. After the completion of allodynia testing for LPS experiments, cold allodynia testing was performed with the application of acetone (Decosterd and Woolf, 2000). In this assay, 10 μl of acetone (99% high-performance liquid chromatography grade; Thermo Fisher Scientific, Waltham, MA) was projected via a 100- μl pipette (Rainin Instruments, Woburn, MA) onto the plantar surface of each hind paw. Acetone was propelled from below via air burst by expressing the pipette, thereby avoiding mechanical stimulation of the paw with the pipette. Total time lifting/clutching each hind paw was recorded with an arbitrary maximum cutoff time of 60 seconds. For all behavioral testing, threshold assessment was performed in a blinded fashion.

Data Analysis. Data were analyzed using Student's *t* test (evaluation of endocannabinoid and AA levels) or one-way or two-way analysis of variance (ANOVA). Tukey's test was used for post hoc analysis after a significant one-way ANOVA. Multiple comparisons after two-way ANOVA were conducted with a Bonferroni post hoc comparison. A *P* value of <0.05 was considered statistically significant. The computer program GraphPad Prism version 4.03 (GraphPad Software Inc., San Diego, CA) was used in all statistical analyses. All data are expressed as the mean \pm S.E.M.

Results

DO34 (30 mg/kg) significantly decreased whole-brain levels of 2-AG ($P < 0.0001$) (Fig. 1A), AEA ($P < 0.05$) (Fig. 1B), and AA ($P < 0.0001$) (Fig. 1C).

To examine whether DO34 produces overt pharmacological effects, we assessed whether it affects spontaneous locomotor behavior, elicits cataleptic effects in the bar test,




Fig. 1. Endocannabinoid levels in whole brain tissue are altered 2 hours after 30 mg/kg DO34. DO34 decreases 2-AG (A), AEA (B), and AA (C) compared with vehicle (Veh). **** $P < 0.0001$; * $P < 0.05$ vs. vehicle. Data reflect the mean \pm S.E.M. $n = 12$ mice/group.

produces antinociception in the warm water tail withdrawal assay, or alters body temperature. Naive mice given vehicle or 1, 3, 10, 30, 50, or 100 mg/kg DO34. DO34 did not display differences of treatment in catalepsy (Fig. 2A), hypothermia ($P = 0.60$) (Fig. 2B), thermal antinociception ($P = 0.13$) (Fig. 2C), or locomotor alterations (defined as time spent immobile; $P = 0.57$) (Fig. 2D).

Having confirmed that the DAGL inhibitor DO34 significantly reduces endocannabinoids and AA in whole brain but does not affect overt motor or sensory behavior, the next set of experiments investigated this compound in the LPS model of inflammatory pain. The dose-response evaluation of the antiallodynic effects of DO34 (1, 3, 10, and 30 mg/kg) at 2 hours postinjection in the von Frey and acetone-induced flinching assays are shown in Fig. 3, A and B, respectively. DO34 dose-dependently reversed LPS-induced mechanical allodynia [$F(3,20) = 14.12$; $P < 0.0001$] and cold allodynia, [$F(3,20) = 15.99$; $P < 0.0001$]. The respective ED₅₀ values [95% confidence interval (CI)] of DO34 in reversing LPS-induced mechanical allodynia and cold allodynia were 3.8 mg/kg (95% CI, 2.8–5.3 mg/kg) and 6.0 mg/kg (95% CI, 4.0–9.0 mg/kg). The potency ratio (95% CI) of DO34 for mechanical versus cold




Fig. 2. Assessment of behavior reveals DO34 treatment does not produce common cannabimimetic effects in naive mice. DO34 does not produce catalepsy (A), antinociception (B), body temperature (C), or change in locomotion (D). Data reflect the mean \pm S.E.M. $n = 6$ mice/group.

allodynia was 1.6 (95% CI, 1.0–2.6), indicating equipotency in the two nociceptive assays. As depicted in Fig. 3C, DO34 significantly reversed allodynia within 30 minutes of intraperitoneal administration, and the antinociceptive effect of 30 mg/kg DO34 persisted for at least 8 hours [interaction between treatment and time: $F(20,125) = 3.58$; $P < 0.0001$].

To evaluate whether the antinociceptive effects of DO34 undergo tolerance, we evaluated von Frey thresholds and acetone-induced flinching in mice that received vehicle or DO34 (30 mg/kg) after one injection or 6 days of repeated administration. As shown in Fig. 4, DO34 retained its anti-allodynic effects after 6 days of repeated administration in the von Frey assay [$F(1,20) = 22.55$; $P < 0.0001$] (Fig. 4A), as well as in the acetone-induced flinching assay [$F(1,20) = 46.34$; $P < 0.0001$] (Fig. 4B).

In the next series of experiments, we evaluated LPS-induced allodynia in DAGL- α ($-/-$) mice or DAGL- β ($-/-$) mice given an intraperitoneal injection of vehicle, DO34 (30 mg/kg), or DO53 (30 mg/kg), a selective ABHD6 inhibitor that served as a control for this off-target of DO34. As previously reported (Wilkerson et al., 2016), DAGL- β ($-/-$) mice displayed an antiallodynic phenotype in the von Frey assay [$F(1,10) = 27.9$; $P < 0.001$] (Fig. 5A). In addition, these mice showed a reduction in acetone-induced flinching [$F(1,10) = 225$; $P < 0.0001$] (Fig. 5B) 24 hours after intraplantar LPS administration. DO34 (30 mg/kg, i.p.) administered at 22 hours after LPS injection reversed mechanical ($P < 0.001$) (Fig. 5A) and cold ($P < 0.0001$) (Fig. 5B) allodynia in DAGL- β ($+/+$) mice but did not alter the antiallodynic phenotypes of the DAGL- β ($-/-$) mice. DO53 (30 mg/kg, i.p.) did not produce significant effects in either mechanical ($P = 0.97$) or cold ($P = 0.49$)

allodynia and did not alter the DAGL- β ($-/-$) antiallodynic phenotype.

In the final experiment, we examined the dose-response relationship of DO34 (0, 1, 3, 10, or 30 mg/kg) in LPS-treated DAGL- α ($-/-$) and ($+/+$) mice. LPS elicited similar magnitudes of mechanical (Fig. 6A) and cold (Fig. 6B) allodynia regardless of genotype. DO34 dose-relatedly reversed LPS-induced mechanical allodynia in DAGL- α ($+/+$) mice [$F(3,20) = 29.42$; $P < 0.0001$] and DAGL- α ($-/-$) mice [$F(3,20) = 4.45$; $P < 0.05$] (Fig. 6A). The respective ED₅₀ (95% CI) values of DO34 in reversing LPS-induced mechanical allodynia in DAGL- α ($+/+$) and ($-/-$) mice were 8.6 mg/kg (95% CI, 6.4–11.5 mg/kg) and 5.9 mg/kg (95% CI, 3.9–8.8 mg/kg). The potency ratio of DAGL- α ($+/+$) to DAGL- α ($-/-$) for mechanical allodynia was 1.4 (95% CI, 0.8–2.5). Likewise, DO34 reversed cold allodynia in DAGL- α ($+/+$) [$F(3,20) = 48.47$; $P < 0.0001$] and DAGL- α ($-/-$) mice [$F(3,20) = 30.99$; $P < 0.0001$] (Fig. 6B). The respective ED₅₀ values (95% CI) of DO34 in reversing LPS-induced cold allodynia in DAGL- α ($+/+$) and ($-/-$) mice were 6.1 mg/kg (95% CI, 4.7–7.9 mg/kg) and 4.5 mg/kg (95% CI, 3.4–6.0 mg/kg). The potency ratio for DAGL- α ($+/+$) to DAGL- α ($-/-$) for cold allodynia was 1.35 (95% CI, 0.9–1.9). Finally, DO53 (30 mg/kg) administered at 22 hours after LPS injection did not produce antinociceptive effects in either genotype (Supplemental Fig. 1).

Discussion

The present study used complementary pharmacological and genetic approaches to test whether blockade of the 2-AG biosynthetic enzymes DAGL- β and DAGL- α produces antinociceptive effects in the LPS model of inflammatory pain. As




Fig. 3. Pharmacological inhibition of DAGL with DO34 reverses LPS-induced mechanical and cold allodynia. (A) DO34 reverses LPS-induced mechanical allodynia in a dose- and time-dependent manner. (B) DO34 reverses LPS-induced cold allodynia in a dose-dependent manner 2 hours after intraperitoneal administration. (C) DO34 significantly reversed allodynia in a time- and dose-dependent manner, within reversal onset at 30 minutes, lasting beyond 8 hours, after intraperitoneal administration. Data reflect the mean \pm S.E.M. $n = 6$ mice/group. *** $P < 0.0001$; ** $P < 0.001$ vs. LPS + vehicle. Filled circles = $P < 0.05$ vs. LPS + vehicle. Veh, vehicle.

previously reported (Wilkerson et al., 2016), DAGL- β ($-/-$) mice were resistant to the development of LPS-induced mechanical allodynia. Moreover these mice were also resistant to the development of LPS-induced cold allodynia. In contrast, DAGL- α ($-/-$) mice displayed full development of LPS-induced mechanical and cold allodynia. These findings suggest that DAGL- α and DAGL- β play differential roles in the development of LPS-induced hyperalgesic states. Thus, whereas DAGL- α is dispensable for the development of LPS-induced allodynia, DAGL- β plays a necessary role in the increased nociceptive behavior after endotoxin treatment.

The disparate roles that DAGL- α and DAGL- β play in LPS-induced allodynia are consistent with the differential expression of these enzymes on cells in the nervous system and peripheral tissue (Hsu et al., 2012). Specifically, DAGL- α is expressed on postsynaptic neurons within the hippocampus, cerebellum, prefrontal cortex, and striatum (Katona et al., 2006; Yoshida et al., 2006; Lafourcade et al., 2007; Uchigashima et al., 2007), and its genetic deletion results in marked decreases in 2-AG, AEA, and AA in the brain (Gao et al., 2010; Tanimura et al., 2010; Shonesy et al., 2014)

and spinal cord (Gao et al., 2010). In contrast, the relative expression of DAGL- β throughout the brain is generally sparse. DAGL- β ($-/-$) mice express wild-type levels of 2-AG in the whole brain (Hsu et al., 2012), and endocannabinoid-mediated forms of retrograde synaptic suppression in these mice are spared (Gao et al., 2010). However, the high expression of DAGL- β on microglia in the central nervous system (Viader et al., 2016) and on macrophages in the periphery (Hsu et al., 2012) is consistent with its role within the innate immune system. Specifically, pharmacological inhibition or genetic deletion of DAGL- β leads to decreased levels of endocannabinoids, AA, prostanoids, and proinflammatory cytokines in LPS-treated peritoneal macrophage cell cultures from C57BL/6 mice (Hsu et al., 2012).

Here we show that acute administration of DO34, at a dose that produced reversal of mechanical and cold allodynia (30 mg/kg), produced significant reductions of 2-AG (~83%), AEA (~42%), and AA (~58%) in naive mice. These findings are in agreement with previous work showing that DO34 reduces 2-AG, AEA, and AA levels in mouse whole brains (Ogasawara et al., 2016). However, the use of whole brain precludes insight of whether DO34 differentially affects lipid levels in discrete brain regions.




Fig. 4. Repeated administration of DO34 prevents LPS-induced mechanical and cold allodynia. (A) Acute or repeated administration of DO34 (30 mg/kg) prevents the expression of LPS-induced mechanical allodynia. (B) Acute or repeated administration of DO34 (30 mg/kg) prevents the expression of LPS-induced cold allodynia. Data reflect the mean \pm S.E.M. $n = 6$ mice/group. ** $P < 0.001$ vs. LPS + vehicle. Veh, vehicle.

Here, we also report that the DAGL inhibitor DO34 dose-dependently reversed LPS-induced mechanical allodynia and cold allodynia. These antinociceptive effects are DAGL- α dispensable, as genetic deletion of this enzyme did not alter the dose-response curves of DO34 for both measures. DO34 also did not alter the antinociceptive phenotype DAGL- β (-/-) mouse response but completely reversed the LPS-induced allodynic responses in DAGL- β (+/+) mice. Additionally, the DO34 time course experiment demonstrates that its antiallodynic effects persist for at least 8 hours, which is consistent with its long duration of action in inhibiting DAGL- β activity (Ogasawara et al., 2016). Besides its actions in DAGL- α and DAGL- β , DO34 also inhibits ABHD6 (Ogasawara et al., 2016), a serine hydrolase that hydrolyzes 2-AG, but to a much lesser extent than monoacylglycerol lipase (Blankman et al., 2007; Marrs et al., 2010). Thus, we tested DO53, a structurally similar compound that inhibits ABHD6 without actions at either DAGL- β or DAGL- α (Ogasawara et al., 2016) in the LPS model of inflammatory pain. DO53 did not reverse LPS-induced allodynia in either DAGL- α (-/-) or (+/+) mice, suggesting that ABHD6 inhibition alone or in combination with DAGL- α inhibition does not elicit antinociceptive effects in this assay. The observation that another ABHD6 inhibitor, KT195, lacked efficacy in reversing LPS-induced mechanical allodynia (Wilkerson et al., 2016) further excludes consideration of the involvement of this enzyme in the results reported here. Although the indirect evidence offered here is consistent with the idea that DAGL- β mediates the

antiallodynic effects of DO34, it does not rule out the possibility of another target.

Another relevant finding in the present study is that repeated administration of DO34 (30 mg/kg) for 6 days continued to prevent the expression of LPS-induced allodynia. Similarly, the antiallodynic effects of the preferential DAGL- β inhibitor KT109 in the LPS model of inflammatory pain did not undergo tolerance. This apparent lack of tolerance is consistent with the observation that DAGL- β (-/-) mice displayed an antiallodynic phenotype in the LPS model of inflammatory pain. It will be important in future studies to ascertain whether repeated administration DAGL inhibitors also reverses nociceptive behavior in chronic models of inflammatory or neuropathic pain. Additionally, given the previously mentioned caveats of target selectivity, there is a need for more selective inhibitors for DAGL- α and DAGL- β , and further studies with these inhibitors are needed to verify our proposed mechanism of action.

The underlying mechanisms for the antinociceptive effects of DO34 remain to be determined but may be related to a reduction of AA and its bioactive metabolites in macrophages expressed in the LPS-treated paw. In particular, KT109 as well as DAGL- β deletion resulted in decreased levels of a variety of proinflammatory lipids and proteins in LPS-stimulated peritoneal macrophage (Hsu et al., 2012). Specifically, prostaglandins are crucial for the development and maintenance of inflammatory pain (Ulmann et al., 2010; Endo et al., 2014; Sugita et al., 2016). DAGL- β inhibition is also protective from microglial activation in the brains of mice repeatedly administered LPS (Viader et al., 2016) and specifically produces reversal of LPS-stimulated proinflammatory cytokine release (Hsu et al., 2012) as well as reverses allodynia and thermal sensitivity in inflammatory, chronic constriction injury of the sciatic nerve, and chemotherapy-induced peripheral neuropathy pain models (Wilkerson et al., 2016). In these models of neuropathic pain, although these analgesic effects appear to be due to DAGL- β inhibition, the relative contribution of additional modulation of AA metabolites remains unclear. However, AA can act as a direct modulator of neuronal activity through its mechanosensory mediatory effects on lipid-sensitive ion channels (Meves, 2008; Brohawn et al., 2014).

Given that these studies represent one of the first in vivo evaluations of DO34, we assessed whether it would produce overt behavioral effects. Accordingly, we examined whether DO34 would produce changes in spontaneous activity or body temperature as well as assess whether it would elicit acute cataleptic or thermal antinociceptive effects. DO34 did not alter spontaneous activity or body temperature and was inactive in a warm water tail withdrawal test for acute thermal antinociception and in the bar test for catalepsy. Other studies have shown that DO34 decreases food intake (Deng et al., 2017). It is important to note here that DAGL- α (-/-) mice display increased anxiogenic behavior in multiple assays used to infer anxiety (i.e., the open-field, light/dark box, and novelty-induced hypophagia test) (Shonesy et al., 2014). Thus, it will be important to assess the effects of DO34 in fear and anxiety assays in future studies.

The results of the present study provide proof of principle that DAGL- β plays a necessary role in the expression of




Fig. 5. Genetic inhibition of DAGL-β in the mediation of LPS-induced mechanical and cold allodynia. (A) LPS-treated DAGL-β (-/-) mice do not develop mechanical allodynia. The DAGL-α and DAGL-β inhibitor DO34 (30 mg/kg) reverses LPS-induced mechanical allodynia in DAGL-β (+/+) mice and does not further alter the antinociceptive effects in DAGL-β (-/-) mice. The ABHD6 inhibitor DO53 did not produce reversal of mechanical allodynia in DAGL-β (+/+) mice. (B) LPS-treated DAGL-β (-/-) mice do not develop cold allodynia. DO34 (30 mg/kg) reverses LPS-induced cold allodynia in DAGL-β (+/+) mice and does not further alter the antinociceptive effects in DAGL-β (-/-) mice. The ABHD6 inhibitor DO53 did not produce reversal of cold allodynia in DAGL-β (+/+) mice. *N* = 6 mice/group. Data reflect the mean ± S.E.M. *****P* < 0.0001 vs. respective genotype control paw.

nociceptive behavior in the LPS model of inflammatory pain. The observation that DAGL-α (-/-) mice display LPS-induced allodynia indicates that this enzyme is dispensable for these effects. Moreover, these null mice served as a useful tool in showing that the antiallodynic effects of DO34 are

independent of its actions on DAGL-β. Taken together, the present study and our previous work (Wilkerson et al., 2016) suggest that DAGL-β represents a potential therapeutic target to relieve pain elicited by the activation of proinflammatory events.




Fig. 6. The role of DAGL-α inhibition in the mediation of DO34 reversal of LPS-induced mechanical and cold allodynia. (A) LPS-treated DAGL-α (-/-) mice normally develop mechanical allodynia. DO34 (30 mg/kg) reverses LPS-induced mechanical allodynia in DAGL-α (+/+) and (-/-) mice. There is no significant shift in the dose-response curve due to genotype. (B) LPS-treated DAGL-α (-/-) mice develop cold allodynia. DO34 (30 mg/kg) reverses LPS-induced cold allodynia in DAGL-α (+/+) and (-/-) mice. There is no significant shift in the dose-response curve due to genotype. C = Non-LPS injected control paw, V = LPS-injected paw with vehicle treatment. *N* = 6 mice/group. Data reflect the mean ± S.E.M. Filled circles = *P* < 0.05 vs. LPS + vehicle in the respective genotype paw.

Authorship Contributions

Participated in research design: Wilkerson, Donvito, Grim, and Lichtman.

Conducted experiments: Wilkerson, Grim, and Abdullah.

Contributed new reagents or analytic tools: Ogasawara and Cravatt.

Performed data analysis: Wilkerson, Grim, and Abdullah.

Wrote or contributed to the writing of the manuscript: Wilkerson and Lichtman.

References

- Beggs S and Salter MW (2013) The known knowns of microglia-neuronal signalling in neuropathic pain. *Neurosci Lett* **557**:37–42.
- Bisogno T, Howell F, Williams G, Minassi A, Cascio MG, Ligresti A, Matias I, Schiano-Moriello A, Paul P, Williams EJ, et al. (2003) Cloning of the first sn1-DAG lipases points to the spatial and temporal regulation of endocannabinoid signaling in the brain. *J Cell Biol* **163**:463–468.
- Blankman JL, Simon GM, and Cravatt BF (2007) A comprehensive profile of brain enzymes that hydrolyze the endocannabinoid 2-arachidonoylglycerol. *Chem Biol* **14**:1347–1356.
- Booker L, Kinsey SG, Abdullah RA, Blankman JL, Long JZ, Ezzili C, Boger DL, Cravatt BF, and Lichtman AH (2012) The fatty acid amide hydrolase (FAAH) inhibitor PF-3845 acts in the nervous system to reverse LPS-induced tactile allodynia in mice. *Br J Pharmacol* **165**:2485–2496.
- Brohawn SG, Su Z, and MacKinnon R (2014) Mechanosensitivity is mediated directly by the lipid membrane in TRAAK and TREK1 K⁺ channels. *Proc Natl Acad Sci USA* **111**:3614–3619.
- Chen S, Xiong J, Zhan Y, Liu W, and Wang X (2015) Wogonin inhibits LPS-induced inflammatory responses in rat dorsal root ganglion neurons via inhibiting TLR4-MYD88-TAK1-mediated NF- κ B and MAPK signaling pathway. *Cell Mol Neurobiol* **35**:523–531.
- Decosterd I and Woolf CJ (2000) Spared nerve injury: an animal model of persistent peripheral neuropathic pain. *Pain* **87**:149–158.
- De Leo JA, Tawfik VL, and LaCroix-Fralish ML (2006) The tetrapartite synapse: path to CNS sensitization and chronic pain. *Pain* **122**:17–21.
- Deng H, Kooijman S, van den Nieuwendijk AM, Ogasawara D, van der Wel T, van Dalen F, Baggelaar MP, Janssen FJ, van den Berg RJ, den Dulk H, et al. (2017) Triazole ureas act as diacylglycerol lipase inhibitors and prevent fasting-induced refeeding. *J Med Chem* **60**:428–440.
- Endo Y, Blinova K, Romantseva T, Golding H, and Zaitseva M (2014) Differences in PGE2 production between primary human monocytes and differentiated macrophages: role of IL-1 β and TRIF/IRF3. *PLoS One* **9**:e98517.
- Gao Y, Vasilyev DV, Goncalves MB, Howell FV, Hobbs C, Reisenberg M, Shen R, Zhang MY, Strassle BW, Lu P, et al. (2010) Loss of retrograde endocannabinoid signaling and reduced adult neurogenesis in diacylglycerol lipase knock-out mice. *J Neurosci* **30**:2017–2024.
- Goncalves MB, Suetterlin P, Yip P, Molina-Holgado F, Walker DJ, Oudin MJ, Zentar MP, Pollard S, Yáñez-Muñoz RJ, Williams G, et al. (2008) A diacylglycerol lipase-CB2 cannabinoid pathway regulates adult subventricular zone neurogenesis in an age-dependent manner. *Mol Cell Neurosci* **38**:526–536.
- Hsu KL, Tsuboi K, Adibekian A, Pugh H, Masuda K, and Cravatt BF (2012) DAGL β inhibition perturbs a lipid network involved in macrophage inflammatory responses. *Nat Chem Biol* **8**:999–1007.
- Ignatowska-Jankowska B, Wilkerson JL, Mustafa M, Abdullah R, Niphakis M, Wiley JL, Cravatt BF, and Lichtman AH (2015) Selective monoacylglycerol lipase inhibitors: antinociceptive versus cannabimimetic effects in mice. *J Pharmacol Exp Ther* **353**:424–432.
- Katona I, Urbán GM, Wallace M, Ledent C, Jung KM, Piomelli D, Mackie K, and Freund TF (2006) Molecular composition of the endocannabinoid system at glutamatergic synapses. *J Neurosci* **26**:5628–5637.
- Kinsey SG, Long JZ, O'Neal ST, Abdullah RA, Poklis JL, Boger DL, Cravatt BF, and Lichtman AH (2009) Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. *J Pharmacol Exp Ther* **330**:902–910.
- Kretzner AC and Regehr WG (2001) Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* **29**:717–727.
- Lafourcade M, Elezgarai I, Mato S, Bakiri Y, Grandes P, and Manzoni OJ (2007) Molecular components and functions of the endocannabinoid system in mouse prefrontal cortex. *PLoS One* **2**:e709.
- Long JZ, Li W, Booker L, Burston JJ, Kinsey SG, Schlosburg JE, Pavón FJ, Serrano AM, Selley DE, Parsons LH, et al. (2009) Selective blockade of 2-arachidonoylglycerol hydrolysis produces cannabinoid behavioral effects. *Nat Chem Biol* **5**:37–44.
- Marrs WR, Blankman JL, Horne EA, Thomazeau A, Lin YH, Coy J, Bodor AL, Muccioli GG, Hu SS, Woodruff G, et al. (2010) The serine hydrolase ABHD6 controls the accumulation and efficacy of 2-AG at cannabinoid receptors. *Nat Neurosci* **13**:951–957.
- Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminski NE, Schatz AR, Gopher A, Almog S, Martin BR, Compton DR, et al. (1995) Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochem Pharmacol* **50**:83–90.
- Meves H (2008) Arachidonic acid and ion channels: an update. *Br J Pharmacol* **155**:4–16.
- Murphy PG, Ramer MS, Borthwick L, Gaudie J, Richardson PM, and Bisby MA (1999) Endogenous interleukin-6 contributes to hypersensitivity to cutaneous stimuli and changes in neuropeptides associated with chronic nerve constriction in mice. *Eur J Neurosci* **11**:2243–2253.
- National Research Council (2011) *Guide for the Care and Use of Laboratory Animals*, National Academies Press, Washington, DC.
- Ogasawara D, Deng H, Viader A, Baggelaar MP, Berman A, den Dulk H, van den Nieuwendijk AM, Soethoudt M, van der Wel T, Zhou J, et al. (2016) Rapid and profound rewiring of brain lipid signaling networks by acute diacylglycerol lipase inhibition. *Proc Natl Acad Sci USA* **113**:26–33.
- Ohno-Shosaku T, Maejima T, and Kano M (2001) Endogenous cannabinoids mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* **29**:729–738.
- Pan B, Wang W, Long JZ, Sun D, Hillard CJ, Cravatt BF, and Liu Q-s (2009) Blockade of 2-arachidonoylglycerol lipase inhibitor 4-nitrophenyl 4-(dibenzo[d][1,3]dioxol-5-yl(hydroxyl)methyl)piperidine-1-carboxylate (JZL184) enhances retrograde endocannabinoid signaling. *J Pharmacol Exp Ther* **331**:591–597.
- Schlosburg JE, Blankman JL, Long JZ, Nomura DK, Pan B, Kinsey SG, Nguyen PT, Ramesh D, Booker L, Burston JJ, et al. (2010) Chronic monoacylglycerol lipase blockade causes functional antagonism of the endocannabinoid system. *Nat Neurosci* **13**:1113–1119.
- Shonesy BC, Bluett RJ, Ramikie TS, Baldi R, Hermanson DJ, Kingsley PJ, Marnett LJ, Winder DG, Colbran RJ, and Patel S (2014) Genetic disruption of 2-arachidonoylglycerol synthesis reveals a key role for endocannabinoid signaling in anxiety modulation. *Cell Reports* **9**:1644–1653.
- Sugaya Y, Yamazaki M, Uchigashima M, Kobayashi K, Watanabe M, Sakimura K, and Kano M (2016) Crucial roles of the endocannabinoid 2-arachidonoylglycerol in the suppression of epileptic seizures. *Cell Reports* **16**:1405–1415.
- Sugita R, Kuwabara H, Kubota K, Sugimoto K, Kihō T, Tengeji A, Kawakami K, and Shimada K (2016) Simultaneous inhibition of PGE2 and PG12 signals is necessary to suppress hyperalgesia in rat inflammatory pain models. *Mediators Inflamm* **2016**:9847840.
- Sugiura T, Kondo S, Sukagawa A, Nakane S, Shinoda A, Itoh K, Yamashita A, and Waku K (1995) 2-Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochem Biophys Res Commun* **215**:89–97.
- Tanimura A, Yamazaki M, Hashimoto Y, Uchigashima M, Kawata S, Abe M, Kita Y, Hashimoto K, Shimizu T, Watanabe M, et al. (2010) The endocannabinoid 2-arachidonoylglycerol produced by diacylglycerol lipase alpha mediates retrograde suppression of synaptic transmission. *Neuron* **65**:320–327.
- Uchigashima M, Narushima M, Fukaya M, Katona I, Kano M, and Watanabe M (2007) Subcellular arrangement of molecules for 2-arachidonoylglycerol-mediated retrograde signaling and its physiological contribution to synaptic modulation in the striatum. *J Neurosci* **27**:3663–3676.
- Ulmann L, Hirbec H, and Rassendren F (2010) P2X4 receptors mediate PGE2 release by tissue-resident macrophages and initiate inflammatory pain. *EMBO J* **29**:2290–2300.
- Viader A, Ogasawara D, Joslyn CM, Sanchez-Alavez M, Mori S, Nguyen W, Conti B, and Cravatt BF (2016) A chemical proteomic atlas of brain serine hydrolases identifies cell type-specific pathways regulating neuroinflammation. *eLife* **5**:e12345.
- Watkins LR, Milligan ED, and Maier SF (2001) Glial activation: a driving force for pathological pain. *Trends Neurosci* **24**:450–455.
- Wilkerson JL, Ghosh S, Bagdas D, Mason BL, Crowe MS, Hsu KL, Wise LE, Kinsey SG, Damaj MI, Cravatt BF, et al. (2016) Diacylglycerol lipase β inhibition reverses nociceptive behaviour in mouse models of inflammatory and neuropathic pain. *Br J Pharmacol* **173**:1678–1692.
- Williams EJ, Walsh FS, and Doherty P (2003) The FGF receptor uses the endocannabinoid signaling system to couple to an axonal growth response. *J Cell Biol* **160**:481–486.
- Wilson RI and Nicoll RA (2001) Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* **410**:588–592.
- Yoshida T, Fukaya M, Uchigashima M, Miura E, Kamiya H, Kano M, and Watanabe M (2006) Localization of diacylglycerol lipase- α around postsynaptic spine suggests close proximity between production site of an endocannabinoid, 2-arachidonoylglycerol, and presynaptic cannabinoid CB1 receptor. *J Neurosci* **26**:4740–4751.
- Yoshino H, Miyamae T, Hansen G, Zambrowicz B, Flynn M, Pedicord D, Blat Y, Westphal RS, Zaczek R, Lewis DA, et al. (2011) Postsynaptic diacylglycerol lipase mediates retrograde endocannabinoid suppression of inhibition in mouse prefrontal cortex. *J Physiol* **589**:4857–4884.

Address correspondence to: Dr. Jenny L. Wilkerson, P.O. Box 980613, Department of Pharmacology and Toxicology, Virginia Commonwealth University, Richmond, VA 23298-0613. E-mail: jenny.wilkerson@vcuhealth.org


Supplementary methods

Measurement of brain endocannabinoid levels

On the day of processing, tissues were weighed and homogenized with 1.4 ml of chloroform/methanol (2:1 v/v containing 0.0348 g of phenylmethanesulfonyl fluoride/ml) after the addition of internal standards to each sample (8 pmol of AEA-d8, 1 nmol of 2-AG-d8, 3.3 nmol and 1 nmol AA-d8, Cayman Chemical). Homogenates were then mixed with 0.3 ml of 0.73% w/v NaCl, vortexed, and then centrifuged for 10 min at $3220 \times g$ (4°C). The aqueous phase plus debris were collected and extracted two more times with 0.8 ml of chloroform. The organic phases from the three extractions were pooled and the organic solvents were evaporated under nitrogen gas. Dried samples were reconstituted with 0.1 ml of chloroform and mixed with 1 ml of ice-cold acetone. The mixtures were then centrifuged for 5 min at $1811 \times g$ and 4°C to precipitate the proteins. The upper layer of each sample was collected and evaporated under nitrogen. Dried samples were reconstituted with 0.1 ml of methanol and placed in autosample vials for analysis.

Liquid chromatography-tandem mass spectrometry with an electrospray ionization source was used to identify and quantify the endocannabinoids and AA. The chromatographic separation was performed using a Discovery HS C18, 2.1×15 cm, $3 \mu\text{m}$ (Supelco, Bellefonte, PA). The mobile phase consisted of water/methanol (10:90) with 0.1% ammonium acetate and 0.1% formic acid. The following ions were monitored in a multiple-reaction-monitoring positive mode: (348 > 62) and (348 > 91) for AEA; (356 > 62) for AEA-d8; (379 > 287) and (379 > 269) for 2-AG; (387 > 96) for 2AG-d8; (300 > 62) and in negative mode: (303 > 259) and (303 > 59) for AA and (311 > 267) for AA-d8. A calibration curve was constructed for each assay based on linear regression with use of the peak area ratios of the calibrators. The extracted standard curves ranged from 0.039 to 40 pmol for AEA and from 0.0625 to 64 nmol for 2-AG, and from 1 nmol to 32 nmol for AA.

Supplementary Results



Supplemental Figure 1. Pharmacological inhibition of ABHD6 in DAGL- α (-/-) mice with LPS-induced mechanical and cold allodynia. (A) The ABHD6 inhibitor DO53 did not produce reversal of mechanical allodynia in either DAGL- α (+/+) or (-/-) mice. (B) DO53 did not produce reversal of cold allodynia in either DAGL- α (+/+) or (-/-) mice. N = 6 mice/group. Data reflect mean \pm SEM, **** p < 0.0001 vs. respective genotype control paw.