

JPET/2002/45195

SIGNIFICANT ROLE OF CERAMIDE PATHWAY IN EXPERIMENTAL GASTRIC ULCER FORMATION IN RATS

Keita Uehara, Soichiro Miura, Tetsu Takeuchi, Takao Taki, Manabu Nakashita,
Masayuki Adachi, Toshiaki Inamura, Toshiko Ogawa, Yasutada Akiba, Hidekazu Suzuki,
Hiroshi Nagata and Hiromasa Ishii

Department of Internal Medicine, School of Medicine, Keio University, Tokyo (K.U.,
Te.T., M.N., M.A., T.I., T.O., Y.A., H.S., H.N., H.I.); Second Department of Internal
Medicine, National Defense Medical College, Saitama (S.M.); and Molecular Medical
Science Institute Otsuka Pharmaceutical Co., Ltd, Japan (Ta.T.)

JPET/2002/45195

Running Title: Ceramide in gastric ulcer

Address for correspondence and galley proofs: Soichiro Miura, M.D.

Professor, Second Department of Internal Medicine, National Defense Medical College,
3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan

Tel: 81-429-95-1211, ext 2367, Fax: 81-429-96-5201, E-mail: miura@me.ndmc.ac.jp

The number of text pages: 26

The number of tables: 0

The number of figures: 11

The number of references: 40

The number of words in the Abstract: 241

The number of words in the Introduction: 648

The number of words in the Discussion: 821

ABBREVIATIONS:

PMA, phorbol-12-myristate-13-acetate. EMSA, electrophoretic mobility shift assay. TNF- α , tumor necrosis factor α . PDTC, pyrrolidine dithiocarbamate. HPTLC, high-performance thin layer chromatography. EDTA, ethylenediamine-tetraacetic acid. EGTA, ethylene glycol-bis- β -aminoethyl ether-N,N,N',N'-tetraacetic acid. DTT, dithiothreitol. PMSF, phenylmethylsulfonyl fluoride. FITC, fluorescein isothiocyanate. BSA, bovine serum albumin. TBS, tris-buffered saline. ANOVA, analysis of variance. DAG, diacylglycerol. PKC, protein kinase C.

Section options is Gastrointestinal, Hepatic, Pulmonary, & Renal

ABSTRACT

Ceramides have emerged as key participants in the signaling pathway of cytokines and apoptosis. We previously revealed that phorbol-12-myristate-13-acetate (PMA) induced experimental ulcers in rat gastric mucosa. In this study, we investigated the role of ceramide in ulcer formation and its relation to the activation of transcription factors and apoptosis. PMA was subserosally injected to rat glandular stomach. Fumonisin B1 (FB1), an inhibitor of ceramide synthase, was administered together with the PMA. The time course of ceramide content was quantified using thin-layer chromatography and the number of apoptotic cells was determined by immunohistochemistry. The activation of transcription factor NF- κ B (nuclear factor kappa B) or AP-1 was evaluated using an electrophoretic mobility shift assay (EMSA). The administration of FB1 attenuated PMA-induced gastric ulcer formation in a dose-dependent manner. Before the appearance of the ulcers became obvious, the ceramide content (C18 and C24 ceramide) increased significantly in the gastric wall. The activation of NF- κ B and AP-1 and an increase in the number of apoptotic cells were also observed. Both of these were significantly inhibited by the coadministration of FB1. However, NF- κ B inhibitors attenuated gastric ulcer formation without affecting the ceramide content or the number of apoptotic cells. Ceramide formation in the stomach significantly contributes to PMA-induced tissue damage, possibly via the activation of transcription factors and an increase in apoptosis in the gastric mucosa. However, after the increase in ceramide levels, the NF- κ B and apoptosis pathways may be separately involved in ulcer formation.

Sphingolipids are derivatives of a number of common chemical backbones known as long-chain (sphingoid) bases (Spiegel and Merrill, 1996). These lipids appear to be ubiquitous among eukaryotic organisms and have long been recognized as having roles in membrane structure. Recent discoveries have revealed that sphingolipids (ceramide, sphingosine, etc.) are highly bioactive compounds that are involved in diverse cell processes, including cell-cell interactions, adhesion, cell proliferation, differentiation, and oncogenic transformation (Hakomori and Igarashi, 1995). In particular, ceramide has received attention as an important bioeffector molecule that may participate in the mediation of some of the actions of extracellular agents, such as tumor necrosis factor α (TNF- α) (Dbaibo et al., 1993; Tepper et al., 1995) and interferon- γ (Kim et al., 1991). Ceramide has also been reported to induce IL-6 production in fibroblasts (Laulederkind et al., 1995), IL-2 secretion in lymphocytes (Mathias et al., 1993), and IL-1-induced E selectin expression in endothelial cells (Masamune et al., 1996), suggesting that ceramide is involved in the regulation of immune function and inflammatory responses.

Several stimuli that induce the generation of ceramide also activate transcription factor NF- κ B (nuclear factor kappa B). The activation of NF- κ B involves the proteasomal degradation of its cytoplasmic inhibitors, I κ Bs (Verma et al., 1995; Baldwin, 1996). This pathway allows the translocation of free, active NF- κ B complexes into the nucleus, where they bind to cognate DNA sequences in the promoter/enhancer regions of a large number of target genes involved in immune responses and inflammation. However, the role of ceramide in NF- κ B activation remains controversial. In several reports, ceramide did not induce the transcription of NF- κ B-dependent genes (Westwick et al., 1995; Latinis and Koretzky, 1996). In others, the ceramide response could not be dissociated from NF- κ B activation (Dbaibo et al., 1993; Betts et al., 1994; Higuchi et al., 1996).

Nevertheless, arguments in favor of a role for ceramide in NF- κ B activation also exist. In several cell lines, including HL-60 leukemia cells, Jurkat cells, and human umbilical vein endothelial cells, the activation of NF- κ B by ceramide has been demonstrated (Schutze et al., 1992; Yang et al., 1993; Masamune et al., 1996). Masamune et al. recently described the formation of NF- κ B-specific DNA-protein complexes consisting of nuclear proteins from Kato III cells (a gastric cancer cell line) that had been treated with C2-ceramide (Masamune et al., 1999). However, no information has been reported on the interaction between ceramide and NF- κ B activation in the gastric mucosa.

Ceramide has also been suggested to have a role in signaling apoptosis induced by the addition of extracellular agents, such as TNF- α (Dbaibo et al., 1993; Kolesnick and Golde, 1994) or the anti-Fas antibody (Cifone et al., 1994), and analogues of ceramide have been reported to induce apoptosis (Obeid et al., 1993; Jarvis et al., 1994; Sweeney et al., 1996). In addition to its acute proinflammatory potency, TNF- α has been shown to have a direct cytotoxic effect on gastric epithelial cells (Fiorucci et al., 1998). However, the relationship between ceramide formation and the induction of apoptosis and gastric mucosal damage during the process of gastric ulcer formation has not been investigated.

We previously reported that the subserosal injection of phorbol-12-myristate-13-acetate (PMA) resulted in the formation of gastric ulcers in the rat gastric mucosa (Takeuchi et al., 2002). In that study, activation of NF- κ B in the gastric mucosa corresponding to the PMA injection sites was observed, and the ulcer formation was significantly inhibited by the inhibitors of NF- κ B or an antibody against TNF- α , suggesting that both NF- κ B activation and the following TNF- α release may contribute to tissue damage in PMA-induced gastric ulcer formation. Using an experimental model, the aims of this study were to 1) investigate whether the activation of ceramide occurs in the gastric mucosa and is involved in the induction of apoptosis and experimental ulcer formation

JPET/2002/45195

and 2) determine whether or not ceramide formation influences the activation of transcription factors, including NF- κ B, in the process of gastric ulcer formations.

METHODS

Animals and ulcer induction

Male Wistar rats, weighing 200–250 g and maintained on standard laboratory chow (Oriental Yeast Mfg., Ltd., Tokyo, Japan), were used in all the experiments. All animals were handled according to the guidelines of Keio University School of Medicine, Animal Research Committee. The rats were not given any food for 24 hours prior to the experiments but were allowed access to tap water ad libitum. Under anesthesia with 30 mg/kg of pentobarbital sodium, the abdomen was opened with a midline incision. The stomach was exposed, and 50 μ l of either PMA (Sigma Chemical Co., St. Louis, MO, USA) (at a dose of 50 μ g) or its vehicle (20% ethanol in saline) were injected into the subserosal layer of the anterior wall of the glandular stomach using a microsyringe. The abdomen was then closed.

At different time intervals (between 0.5 hours and 48 hours) after the injection of PMA or its vehicle, the rats were killed using an overdose of sodium pentobarbital (500 mg/kg). Their stomachs were rapidly removed, opened along the greater curvature, and rinsed with cold normal saline. The surface area of each lesion in the gastric mucosa was assessed macroscopically. Mucosal injury was also histologically evaluated using paraffin sections stained with hematoxylin and eosin and scored using a 0–4 scale based on the following criteria: 0 = normal; 1 = patches of superficial necrosis; 2 = vasocongestion and focal necrosis of less than one third of the mucosa; 3 = vasocongestion and focal necrosis of more than one third of the mucosa, but not reaching the full thickness; and 4 = extensive vasocongestion and necrosis involving the full thickness of the mucosa.

Administration of various inhibitors

To examine ceramide activation in this model, an inhibitor of sphingolipid biosynthesis, fumonisin B1 (FB1) 0.01–10 μ M (10 μ l of 0.05–50 μ M FB1 equals 0.5–500 pmols)(Sigma), was co-administered along with the PMA. To determine the role of NF- κ B in PMA-induced ulcer formation, pyrrolidine dithiocarbamate (PDTC) 100 mM

(10 μ l of 500 mM PDTC equals 5 μ moles) (Sigma) and an NF- κ B decoy 15 mM (10 μ l of 75 μ M decoy of NF- κ B equals 0.75 μ moles) were also locally injected into the stomach in combination with the PMA. The sequence of the synthetic double-stranded oligodeoxynucleotides (NF- κ B decoy) was as follows:

5'-CCTTGAAGGGATTTCCCTCC-3'

3'-GGAACTTCCCTAAAGGGAGG-5'.

We chose these NF- κ B inhibitor concentrations, because there is evidence that these doses sufficiently inhibited the activation of NF- κ B as well as ulcer formation in our previous experiments (Takeuchi et al., 2002). To assess the apoptotic effects in PMA-induced ulcer formation, Z-DEVD-FMK, a caspase 3/7 inhibitor 10–100 nM (10 μ l of 50–500 nM caspase 3/7 inhibitors equals 0.5–5 pmoles)(Calbiochem, U.S. and Canada) and Ac-Tyr-Val-Ala-Asp-H, a caspase1/4 inhibitor 10–100 nM (10 μ l of 50–500 nM caspase 1/4 inhibitor equals 0.5–5 pmoles) (Peptide Institute, Osaka, Japan) were also locally injected into the stomach in combination with the PMA.

Determination of ceramide content in the stomach

The time course of the ceramide content in the stomach was quantified. The excised stomachs were cut along the greater curvature and rinsed with physiological saline. Approximately 0.5 g of the tissue sample, including the ulcer lesions, was removed and minced, and lipid extraction was performed by a modified version of the method described by Bligh and Dyer (Berg et al., 1997). After the extraction of the major lipids, the neutral lipids, including the ceramides, were separated using high-performance thin layer chromatography (HPTLC) (Silicagel 60, Merck, Germany) (Kasama et al., 1996). The dried lipids were then resolved by thin-layer chromatography using petroleum ether and diethyl ether (7:3) as the first solvent and chloroform and methanol (95:5) as the second solvent. After lipid separation, the HPTLC plate was sprayed with a primulin reagent until thoroughly wet and then air dried completely. The lipids were visualized under UV light at 365 nm and analyzed with a densitometer (FluorchemTM 8000, Alpha Innotech Co., San Leandro, CA, USA).

Activation of NF- κ B and AP-1

The activation of NF- κ B and AP-1 was assessed in the PMA-injected gastric mucosa using an electrophoretic mobility shift assay (EMSA). The stomachs were excised under pentobarbital anesthesia at different time intervals (between 15 minutes and 3 hours) after the PMA administration. Glandular stomach was homogenized on ice in 3.5 μ l/mg of buffer containing 10 mM HEPES (pH: 7.9), 10 mM KCl, 0.1 mM ethylenediamine tetraacetic acid (EDTA), 0.1 mM ethylene glycol-bis- β -aminoethyl ether-N,N,N',N'-tetraacetic acid (EGTA), 1 mM dithiothreitol (DTT), 0.5 mM phenylmethylsulfonyl fluoride (PMSF), and 2 μ g/ml each of antipain, chymostatin, bestatin, pepstatin, and leupeptin. The lysate was centrifuged at 8000 g for 2 minutes at 4°C, and the cytoplasmic supernatant was removed and frozen. The nuclear pellet was reconstituted in 1.5 μ l/mg of buffer containing 20 mM HEPES (pH: 7.9), 0.4 M NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM DTT, 1 mM PMSF, and 2 μ g/ml each of antipain, chymostatin, bestatin, pepstatin, and leupeptin, followed by vigorous vortexing for 20 minutes at 4°C. The nuclear lysate was centrifuged at 14,000 g for 5 minutes, and the nuclear extracts were divided into aliquots, frozen, and stored at -80°C. A gel shift assay was performed using a fluorescein isothiocyanate (FITC)-labeled NF- κ B synthetic double-stranded oligonucleotide (AGTTGAGGGGACTTTCCCAGGC) and an AP-1 synthetic double-stranded oligonucleotide (CCAAAGTGCTGAGTCACTAAT). Equal amounts of nuclear proteins (20 μ g) were incubated for 30 minutes at room temperature with 1 pg of labeled NF- κ B and AP-1 consensus oligonucleotide. After incubation, the samples were electrophoresed through nondenaturing 6% polyacrylamide gels at 4°C, 160 V for 2 hours. For the competition analysis, a 100-fold molar excess of unlabeled consensus oligonucleotide was added to the samples. The fluorescence intensity of the

JPET/2002/45195

gels was quantified using a fluorescence laser scanning system equipped with a computer-assisted image analyzer (FluorImager 575; Molecular Dynamics, Sunnyvale, CA, USA).

Determination of apoptosis in the gastric mucosa

Apoptosis in the stomach after PMA administration was immunohistochemically examined using a polyclonal antibody to ss-DNA (Korkolopoulou et al., 2001). The area of the stomach containing the ulcer was rapidly excised and processed using routine techniques before being embedded in paraffin. Sections (4 μ m thick) were then prepared and mounted on glass slides. Deparaffinized sections were treated with 3% hydrogen peroxide for 20 minutes to block endogenous peroxide. After blocking in 10% non-immune serum for 10 minutes at room temperature, sections were incubated for 40 minutes at room temperature with primary antibody (anti-ss-DNA, polyclonal rabbit, DAKO, Carpinteria, CA, USA) diluted 1:100 with 0.1% bovine serum albumin (BSA) in 0.05 M tris-buffered saline (TBS). The slides were washed three times with 0.05 M TBS-Tween for 5 minutes followed by incubation for 30 minutes with rabbit peroxidase (DAKO). After being washed for 5 minutes in TBS-Tween, the sections were stained using a Diaminobenzidine Reagent set (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, MD, USA) and observed by microscopy (Nikon ECLIPSE-E-600, Tokyo, Japan). Negative controls containing nonimmune rabbit serum or the omission of the primary antibody were also prepared. Staining for all antibodies was assessed in a blind manner by the same observer.

Statistical analysis

All results were expressed as the means \pm SEM. Differences among groups were evaluated using a one-way analysis of variance (ANOVA) and Fisher's post hoc test. The

JPET/2002/45195

histological damage scores were analyzed by Mann-Whitney's U test. Statistical significance was set at $p < 0.05$.

RESULTS

Figure 1A shows representative pictures of a PMA-induced gastric mucosal lesion 48 hours after the subserosal injection of PMA and the inhibitory effect of different concentrations of FB1 (0.1 and 10 μM) on ulcer formation. The ulcers produced by the PMA injection arose at the injection site in the stomach, but were significantly inhibited by FB1; a fumonisin dose of 10 μM almost completely blocked the PMA-induced ulcer formation. **Figure 1B** shows the area of the mucosal lesions and the histological damage scores. The size of ulcers and histological damage induced by PMA were dose-dependently inhibited by the fumonisin treatment. FB1 doses of 0.1 μM or more significantly attenuated the area of the mucosal lesions and the histological damage score.

Figure 2A shows the time course for the changes in C18 and C24 ceramide contents in gastric mucosal lesions after the injection of 50 μg of PMA. In the rat gastric mucosa, the main ceramides induced by PMA were C18 and C24. **Figure 2B** shows a quantitative analysis of the changes in C18 (left panel) and C24 (right panel) ceramide contents. The amounts of both ceramides significantly increased 3 hours after PMA injection and reached their maximal values at 3–6 hours. Thereafter, the ceramide contents began to decrease, but remained at elevated levels, compared with the controls at 48 hours.

Figure 3A shows the inhibitory effect of FB1 (10 μM) and NF- κB inhibitors (PDTC and NF- κB decoy) on ceramide activation 3 hours after PMA administration, while **Figure 3B** shows the quantitative analysis of the changes in C18 (left panel) and C24 (right panel) ceramide contents caused by these treatments. The activation of both ceramides in response to PMA was significantly attenuated by FB1. However, these activations were not significantly inhibited by the NF- κB inhibitors, PDTC, or the NF- κB decoy.

Figure 4 shows the effect of FB1 on the activation of NF- κB and AP-1 in the gastric mucosa 3 hours after the injection of PMA, as determined by EMSA using FITC-labeled NF- κB and an AP-1 consensus oligonucleotide. PMA treatment induced a significant

increase in NF- κ B, and AP-1 binding activity compared with the vehicle treatment (control). The increase in NF- κ B binding activity at 3 hours after PMA administration was significantly attenuated by treatment with FB1 (0.01–10 μ M). The increase in AP-1 binding activity at 3 hours after PMA administration was also dose-dependently attenuated by treatment with FB1. Although not shown in this figure, the activation of NF- κ B binding activity was also inhibited by the NF- κ B inhibitors, PDTC, and the NF- κ B decoy.

Figure 5 shows the effect of caspase inhibitors on PMA-induced gastric ulcer formation and the histological damage scores. Caspase 3/7 and caspase 1/4 inhibitors both attenuated the PMA-induced gastric mucosal lesions and histological damage score in a dose-dependent manner when assessed 48 hours after treatment.

Figure 6 presents representative microscopic findings showing apoptotic cells in the gastric mucosa after PMA administration, as determined by immunohistochemistry using a polyclonal antibody against ss-DNA. In control ethanol-treated mucosa specimens, only a few apoptotic cells were observed in the gastric mucosa. In contrast, a few apoptotic cells were observed at the tip of the mucosa 0.5 hours after PMA treatment, and their number significantly increased at 3 hours after treatment. Six hours after treatment, the mucosal surface has been destroyed and the diffuse pattern of apoptotic cells was observed in the deeper layers of the stomach. After 24 hours, the distribution of these apoptotic cells extended deeply toward the submucosa in the area corresponding to the lesion and in its vicinity.

Figure 7 shows the time course for the changes in the number of apoptotic cells in the gastric mucosa after PMA administration, as immunohistochemically assessed using microscopy, and the effects of FB1, NF- κ B inhibitors, and caspase 3/7 and caspase 1/4 inhibitors. A significant increase in apoptotic cells was observed at 3 hours after administration, reaching maximum value at 24 hours after administration. Thereafter, the number of apoptotic cells began to decrease somewhat with the development of mucosal defects. FB1, the caspase 3/7 inhibitor, and the caspase 1/4 inhibitor significantly inhibited the PMA-induced increase in the number of apoptotic cells at 24 hours after

JPET/2002/45195

administration (**Figs. 6F and 7**). In contrast, neither PDTC nor the NF- κ B decoy significantly attenuated the increase in the number of apoptotic cells after PMA administration, although in our previous paper we demonstrated that the same concentrations of PDTC or NF- κ B significantly attenuated the ulcer area and tissue damage (Takeuchi et al., 2002).

DISCUSSION

Fumonisin are a family of mycotoxins that are produced by some strains of *Fusarium moniliforme* (Sydenham et al., 1990). FB1 is a potent inhibitor of sphinganine (sphingosine) N-acetyltransferase (ceramide synthase) *in vivo*, and it exhibits competitive-type inhibition with respect to both substrates of this enzyme (Wang et al., 1991). Our present results showing that the blockade of ceramide synthase by FB1 successfully attenuates experimentally induced gastric ulcer formation suggest the importance of *de novo* ceramide synthesis in this process. Ceramide synthase is responsible for the acylation of sphinganine in the *de novo* biosynthetic pathway for sphingolipids as well as the reutilization of sphingosine derived from sphingolipid turnover (Wang et al., 1991; Merrill et al., 1993). Although ceramide derived from the hydrolysis of sphingomyelin in response to extracellular signals appears to be important in most pathways (Ballou et al., 1996; Hannun, 1996), ceramide production from *de novo* synthesis via ceramide synthase may also largely account for its bioactive roles as a messenger in the stimulation of a variety of cellular functions. Ohta et al. demonstrated that HL60 cells treated with PMA exhibited an increase in both ceramide and sphingosine, which agrees with our present results (Ohta et al., 1995). In the case of PMA-injected gastric mucosa, the excess production of ceramide via ceramide synthase plays a critical role in the development of gastric tissue damage, although the exact downstream processes remain unknown.

In an earlier study, we presented data indicating that the activation of NF- κ B in the gastric mucosa after PMA injection generally corresponded to the location of the gastric gland cells (Takeuchi et al., 2002). In that study, we also revealed the critical role of NF- κ B activation in the formation of PMA-induced gastric ulcer. The present study shows that the activation of NF- κ B and AP-1 induced by PMA was significantly inhibited by treatment with FB1, suggesting that ceramide synthesis is involved in the activation of these transcription factors. Johns et al. demonstrated that the ceramide pathway additionally activates proteins that bind to an AP-1 consensus site, suggesting that the ceramide pathway may serve more globally to induce genes that are responsive to both

NF- κ B and AP-1 (Johns et al., 1994). Since we previously found that PMA-induced acute gastric ulcer formation is mediated by the strong activation of NF- κ B in situ, the blocking effect of fumonisin B1 may arise from significant attenuation in transcription factors, especially NF- κ B. Phorbol esters, such as PMA, stimulate PKC by mimicking the effects of diacylglycerol (DAG) and also induce NF- κ B activation. Fernandez and Dobbelaere showed that ceramide and PMA have a synergistic effect on the degradation of I κ B in primary lymph node T-cells as well as in transformed T-cells (Fernandez and Dobbelaere, 1999). Thus, ceramide may potentiate NF- κ B activation induced in PMA-injected gastric mucosa. The activation of NF- κ B leads to the activation of various pro-inflammatory molecules, including TNF- α , IL-2, IL-6, inducible nitric oxide synthase (Jourdeuil et al., 1997), and adhesion molecules responsible for leukocyte-endothelial interactions (Dhawan et al., 1997). These inflammatory mediators could in turn activate NF- κ B, initiating a vicious inflammatory cycle that is likely to lead to tissue damage (Barnes and Karin, 1997).

In the present study, we clearly demonstrated that the number of apoptotic cells was increased 3 hours after the injection of PMA. This increase in apoptotic cells initially occurred mainly in the upper part of the glandular mucosa and thereafter in the deeper layer at 6 hours, and extended deeply toward the submucosa at 24 hours. The increase in apoptotic cells in the gastric mucosa was similar to a previously reported time course for ulcer formation (Takeuchi et al., 2002). In addition, both caspase 3/7 and caspase 1/4 inhibitors significantly attenuated PMA-induced ulcer formation in this study. This suggests that apoptosis is involved in the formation of gastric ulcers. The increase in apoptosis in the gastric mucosa corresponded with the increase in C18 and C24 ceramide content in the stomach wall. We also observed a significant reduction in PMA-induced apoptosis in the gastric mucosa after treatment with FB1. Ceramide is known to be involved in signaling apoptosis, and ceramide analogues have been reported to induce apoptosis in vitro in using gastric mucosal cell lines (Johal and Hanson, 2000; Shimada et al., 2000). These findings suggest that the ceramide pathway may be strongly associated with the increase in apoptotic cell death in the PMA-injected mucosa, although the

JPET/2002/45195

involvement of other pathways, including ICE-like proteases, cannot be ruled out (Sweeney et al., 1996). However, it should also be noted that the inhibition of NF- κ B activation did not successfully prevent the increase in apoptosis in this ulcer model. Even though ceramide can activate NF- κ B in the gastric mucosa and NF- κ B activation contributes to PMA-induced ulcer formation, the NF- κ B and apoptosis pathways may be separately involved in the ulcer formation (Fig. 8). Further investigation of the downstream signals in ceramide activation is necessary to elucidate the exact role of this molecule in the formation of gastric ulcers.

References

- Baldwin AS Jr (1996) The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol* **14**: 649-683.
- Ballou LR, Lauderkind SJ, Rosloniec EF, and Raghow R (1996) Ceramide signalling and the immune response. *Biochim Biophys Acta* **1301**: 273-287.
- Barnes PJ and Karin M (1997) Nuclear factor-kappa B: a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med* **336**: 1066-1071.
- Berg H, Magard M, Johansson G, and Mathiasson L (1997) Development of a supercritical fluid extraction method for determination of lipid classes and total fat in meats and its comparison with conventional methods. *J Chromatogr A* **785**: 345-352.
- Betts JC, Agranoff AB, Nabel GJ, and Shayman JA (1994) Dissociation of endogenous cellular ceramide from NF-kappa B activation. *J Biol Chem* **269**: 8455-8458.
- Cifone MG, De Maria R, Roncaioli P, Rippo MR, Azuma M, Lanier LL, Santoni A, and Testi R (1994) Apoptotic signaling through CD95 (Fas/Apo-1) activates an acidic sphingomyelinase. *J Exp Med* **180**: 1547-1552.
- Dbaibo GS, Obeid LM, and Hannun YA (1993) Tumor necrosis factor-alpha (TNF-alpha) signal transduction through ceramide. Dissociation of growth inhibitory effects of TNF-alpha from activation of nuclear factor-kappa B. *J Biol Chem* **268**: 17762-17766.
- Dhawan S, Singh S, and Aggarwal BB (1997) Induction of endothelial cell surface adhesion molecules by tumor necrosis factor is blocked by protein tyrosine phosphatase inhibitors: role of the nuclear transcription factor NF-kappa B. *Eur J Immunol* **27**: 2172-2179.

JPET/2002/45195

- Fernandez PC and Dobbelaere DA (1999) Ceramide synergizes with phorbol ester or okadaic acid to induce IkappaB degradation. *Biochem Biophys Res Commun* **263**: 63-67.
- Fiorucci S, Antonelli E, Migliorati G, Santucci L, Morelli O, Federici B, Morelli A (1998) TNF alpha processing enzyme inhibitors prevent aspirin-induced TNF alpha release and protect against gastric mucosal injury in rats. *Aliment Pharmacol Ther* **12**: 1139-1153.
- Hakomori S and Igarashi Y (1995) Functional role of glycosphingolipids in cell recognition and signaling. *J Biochem (Tokyo)* **118**:1091-1103.
- Hannun YA (1996) Functions of ceramide in coordinating cellular responses to stress. *Science* **274**: 1855-1859.
- Higuchi M, Singh S, Jaffrezou JP, and Aggarwal BB (1996) Acidic sphingomyelinase-generated ceramide is needed but not sufficient for TNF-induced apoptosis and nuclear factor-kappa B activation. *J Immunol* **157**: 297-304.
- Jarvis WD, Kolesnick RN, Fornari FA, Traylor RS, Gewirtz DA, and Grant S (1994) Induction of apoptotic DNA damage and cell death by activation of the sphingomyelin pathway. *Proc Natl Acad Sci U S A* **91**: 73-77.
- Johal K and Hanson PJ (2000) Opposite effects of flurbiprofen and the nitroxybutyl ester of flurbiprofen on apoptosis in cultured guinea-pig gastric mucous cells. *Br J Pharmacol* **130**: 811-818.
- Johns LD, Sarr T, and Ranges GE (1994) Inhibition of ceramide pathway does not affect ability of TNF-alpha to activate nuclear factor-kappa B. *J Immunol* **152**: 5877-5882.
- Jourd'heuil D, Morise Z, Conner EM, Kurose I, and Grisham MB (1997) Oxidant-regulation of gene expression in the chronically inflamed intestine. *Keio J*

JPET/2002/45195

Med **46**: 10-15.

Kasama T, Hisano Y, Nakajima M, Handa S, and Taki T (1996) Microscale analysis of glycosphingolipids by TCL blotting/secondary ion mass spectrometry: a novel blood group A-active glycosphingolipid and changes in glycosphingolipid expression in rat mammary tumor cells with different metastatic potentials. *Glycoconj J* **13**: 461-469.

Kim MY, Linardic C, Obeid L, and Hannun Y (1991) Identification of sphingomyelin turnover as an effector mechanism for the action of tumor necrosis factor alpha and gamma-interferon. Specific role in cell differentiation. *J Biol Chem* **266**: 484-489.

Kolesnick RN and Golde DW (1994) The sphingomyelin pathway in tumor necrosis factor and interleukin-1 signaling. *Cell* **77**: 325-328.

Korkolopoulou PA, Konstantinidou AE, Patsouris ES, and Christodoulou PN (2001) Detection of apoptotic cells in archival tissue from diffuse astrocytomas using a monoclonal antibody to single stranded DNA. *J Pathol* **193**: 377-382.

Latinis KM and Koretzky GA (1996) Fas ligation induces apoptosis and Jun kinase activation independently of CD45 and Lck in human T cells. *Blood* **87**: 871-875.

Laulederkind SJ, Bielawska A, Raghov R, Hannun YA, and Ballou LR (1995) Ceramide induces interleukin 6 gene expression in human fibroblasts. *J Exp Med* **182**: 599-604.

Masamune A, Igarashi Y, and Hakomori S (1996) Regulatory role of ceramide in interleukin (IL)-1 beta-induced E-selectin expression in human umbilical vein endothelial cells. Ceramide enhances IL-1 beta action, but is not sufficient for E-selectin expression. *J Biol Chem* **271**: 9368-9375.

Masamune A, Shimosegawa T, Masamune O, Mukaida N, Koizumi M, and Toyota T (1999) Helicobacter pylori-dependent ceramide production may mediate increased

JPET/2002/45195

interleukin 8 expression in human gastric cancer cell lines. *Gastroenterology* **116**: 1330-1341.

Mathias S, Younes A, Kan CC, Orlow I, Joseph C, and Kolesnick RN (1993) Activation of the sphingomyelin signaling pathway in intact EL4 cells and in a cell-free system by IL-1 beta. *Science* **259**: 519-522.

Merrill AH Jr, van Echten G, Wang E, and Sandhoff K (1993) Fumonisin B1 inhibits sphingosine (sphinganine) N-acyltransferase and de novo sphingolipid biosynthesis in cultured neurons in situ. *J Biol Chem* **268**: 27299-27306.

Obeid LM, Linardic CM, Karolak LA, and Hannun YA (1993) Programmed cell death induced by ceramide. *Science* **259**: 1769-1771.

Ohta H, Sweeney EA, Masamune A, Yatomi Y, Hakomori S, and Igarashi Y (1995) Induction of apoptosis by sphingosine in human leukemic HL-60 cells: a possible endogenous modulator of apoptotic DNA fragmentation occurring during phorbol ester-induced differentiation. *Cancer Res* **55**: 691-697.

Schutze S, Potthoff K, Machleidt T, Berkovic D, Wiegmann K, and Kronke M (1992) TNF activates NF-kappa B by phosphatidylcholine-specific phospholipase C-induced "acidic" sphingomyelin breakdown. *Cell* **71**: 765-776.

Shimada T, Hiraishi H, and Terano A (2000) Hepatocyte growth factor protects gastric epithelial cells against ceramide-induced apoptosis through induction of cyclooxygenase-2. *Life Sci* **68**: 539-546.

Spiegel S and Merrill AH Jr (1996) Sphingolipid metabolism and cell growth regulation. *FASEB J* **10**:1388-1397.

Sweeney EA, Sakakura C, Shirahama T, Masamune A, Ohta H, Hakomori S, and Igarashi Y (1996) Sphingosine and its methylated derivative N,N-dimethylsphingosine (DMS) induce apoptosis in a variety of human cancer cell

JPET/2002/45195

lines. *Int J Cancer* **66**: 358-366.

Sydenham EW, Gelderblom WCA, Thiel PG, and Marasas WFO (1990) Evidence of the natural occurrence of fumonisin B1 a mycotoxin produced by *Fusarium moniliforme* in corn. *J Agric Food Chem* **38**: 285-290.

Takeuchi T, Miura S, Wang L, Uehara K, Mizumori M, Kishikawa H, Hokari R, Higuchi H, Adachi M, Nakamizo H, and Ishii H (2002) Nuclear factor kappa B and TNF-alpha mediate gastric ulceration induced by phorbol myristate acetate. *Dig Dis Sci* **47**: 2070-2078.

Tepper CG, Jayadev S, Liu B, Bielawska A, Wolff R, Yonehara S, Hannun YA, and Seldin MF (1995) Role for ceramide as an endogenous mediator of Fas-induced cytotoxicity. *Proc Natl Acad Sci U S A* **92**:8443-8447.

Verma IM, Stevenson JK, Schwarz EM, Van Antwerp D, and Miyamoto S (1995) Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes Dev* **9**:2723-2735.

Wang E, Norred WP, Bacon CW, Riley RT, and Merrill AH Jr (1991) Inhibition of sphingolipid biosynthesis by fumonisins. Implications for diseases associated with *Fusarium moniliforme*. *J Biol Chem* **266**: 14486-14490.

Westwick JK, Bielawska AE, Dbaibo G, Hannun YA, and Brenner DA (1995) Ceramide activates the stress-activated protein kinases. *J Biol Chem* **270**: 22689-22692.

Yang Z, Costanzo M, Golde DW, and Kolesnick RN (1993) Tumor necrosis factor activation of the sphingomyelin pathway signals nuclear factor kappa B translocation in intact HL-60 cells. *J Biol Chem* **268**: 20520-20523.

Figure captions

Figure 1. (A) Representative photographs of the gross appearance of rat gastric mucosa 48 hours after subserosal injection of phorbol myristate acetate (PMA). PMA (50 μg) injection revealed an active ulcer (arrow) (left panel). Fumonisin B1 (FB1) (0.1 μM) decreased the area of gastric ulcer (middle panel), and fumonisin B1 at the dose of 10 μM completely inhibited the ulcer formation macroscopically (right panel). (B) Effect of different doses of FB1 on PMA-induced macroscopically assessed mucosal lesions (ulcer area, left panel) and histological damage scores (right panel) at 48 hours. PMA (50 μg) was injected into the subserosal layer of the anterior wall of the glandular stomach, and vehicle (20% ethanol in saline) was injected as a control. The ulcer area is expressed as the area of mucosal lesions (mm^2) and histological damage was evaluated in paraffin sections stained with hematoxylin and eosin and given a score on a 0–4 scale. * $P < 0.05$ vs. control. # $P < 0.05$ vs. PMA alone. Values are means \pm SEM in six animals.

Figure 2. Time course of changes in C18 and C24 ceramide in the gastric mucosal walls after 50 μg PMA injection. After the extraction of major lipids, samples were resolved in thin-layer chromatography using petroleum ether and diethyl ether (7:3) as the first solvent and chloroform and methanol (95:5) as the second solvent. The HPTLC plates were sprayed with primulin reagent until they became wet and then air dried thoroughly. The lipids were visualized under UV light at 365 nm and analyzed with a densitometer. (A) Representative pictures of thin-layer chromatography. C18 and C24: standard of C18 and C24 ceramide (1 mg/ml). (B) Densitometric analysis of changes in C18 (left panel) and C24 (right panel) ceramide contents. * $P < 0.05$ vs. 0 h (no treatment). Values are means \pm SEM in six animals.

Figure 3. Inhibitory effect of FB1 and NF- κB inhibitors on ceramide activation in the gastric mucosal walls 3 hours after 50 μg PMA injection. (A) Representative pictures of thin-layer chromatography. C18 and C24: standard of C18 and C24 ceramide (1 mg/ml).

JPET/2002/45195

FB1: fumonisin B1 at the dose of 10 μ M. PDTC: NF- κ B inhibitor, pyrrolidine dithiocarbamate (100 mM). Decoy of NF- κ B (15 mM). These inhibitors were locally administered to the gastric walls in combination with PMA. (B) Densitometric analysis of changes in C18 (left panel) and C24 (right panel) ceramide contents. As a control, vehicle (20% ethanol in saline) was injected. * P <0.05 vs. control. # P <0.05 vs. PMA alone. Values are means \pm SEM in six animals.

Figure 4. Effect of FB1 on activation of transcription factors NF- κ B (A) and AP-1 (B) in the PMA-injected gastric mucosa as determined by electrophoretic mobility shift assay (EMSA) using FITC-labeled consensus oligonucleotides. PMA treatment induced a significant increase in NF- κ B and AP-1 binding activity as compared with vehicle treatment (control) at 3 hours as indicated by arrows. FB1 was locally injected into the stomach in combination with PMA. This shows a representative picture from four experiments with similar results. Activation of NF- κ B (C) and AP-1 (D) was estimated by densitometric scanning and expressed as a percent of the control. * P <0.05 vs. control (vehicle treatment). # P <0.05 vs. PMA alone. Values are means \pm SEM in four animals.

Figure 5. Effect of caspase inhibitors on PMA-induced macroscopically assessed mucosal lesions (ulcer area, left panel) and histological damage scores (right panel). The inhibitors of caspase 3/7 (10, 100 nM) and caspase 1/4 (10, 100 nM) were locally administered to the gastric walls with PMA and the area of mucosal lesions was assessed as an ulcer area at 48 hours. * P <0.05 vs. control (vehicle treatment). # P <0.05 vs. PMA alone. Values are means \pm SEM in six animals.

Figure 6. Representative photographs of the microscopic findings of apoptotic cells in the rat gastric mucosa after subserosal injection of PMA (50 μ g). Apoptotic cells were determined by immunohistochemistry using a polyclonal antibody against ss-DNA. Upper panels shows pictures taken with a x200 objective and lower panels shows their

JPET/2002/45195

higher magnification (x400). (A) Vehicle (20% ethanol in saline) treatment as control. (B) 0.5 hours after PMA treatment. (C) 3 hours after PMA treatment. (D) 6 hours after PMA treatment. (E) 24 hours after PMA treatment. (F) 24 hours after PMA treatment with FB1 (10 μ M) treatment.

Figure 7. Time course of changes in apoptotic cell number in the gastric mucosa after 50 μ g PMA injection and the inhibitory effect of FB1 and NF- κ B inhibitors. The number of apoptotic cells was determined by immunohistochemistry on more than five sections per animal using a polyclonal antibody against ss-DNA and expressed as a positive cell number per microscopic field average (x400). 0h: no treatment. 0.5h–48h: number of hours after PMA treatment. Control: 24 hours after vehicle treatment (20% ethanol alone). +FB1: 24 hours after PMA with FB1 (10 μ M) treatment. +PDTC: 24 hours after PMA with pyrrolidine dithiocarbamate (100 mM). +decoy of NF- κ B: 24 hours after PMA with decoy of NF- κ B (15 mM). +caspase 3/7 inhibitor: 24 hours after PMA with caspase 3/7 inhibitor (100 nM). +caspase 1/4 inhibitor: 24 hours after PMA with caspase 1/4 inhibitor (100 nM). * P <0.05 vs. time 0 h. # P <0.05 vs. PMA alone (24 h). Values are means \pm SEM in six animals.

Figure 8. Schematic diagram illustrating the pathways involved in PMA-induced ulcer formation and the steps experimentally performed in this study.

FOOTNOTES

This study was supported in part by Grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Science and Culture of Japan and by grants from Keio University, School of Medicine, and from National Defense Medical College.

Address for reprints request: Hiromasa Ishii, M.D.

Professor of Internal Medicine, School of Medicine, Keio University

35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan


Tel: 81-3-3353-1211, ext 62260, Fax: 81-3-3356-9654, E-mail: hishii@sc.itc.keio.ac.jp

Keita Uehara,¹⁾ Soichiro Miura,²⁾ Tetsu Takeuchi,¹⁾ Takao Taki,³⁾ Manabu Nakashita,¹⁾ Masayuki Adachi,¹⁾ Toshiaki Inamura,¹⁾ Toshiko Ogawa,¹⁾ Yasutada Akiba,¹⁾ Hidekazu Suzuki,¹⁾ Hiroshi Nagata¹⁾ and Hiromasa Ishii¹⁾

Department of Internal Medicine¹⁾, School of Medicine, Keio University, Tokyo and Second Department of Internal Medicine, National Defense Medical College²⁾, Saitama and Molecular Medical Science Institute Otsuka Pharmaceutical³⁾ Co., Ltd, Japan.

Fig. 1

A



PMA alone

**PMA
+ fumonisin B1
(0.1 μ M)**

**PMA
+ fumonisin B1
(10 μ M)**

Fig. 1

B

(mm²)





Fig. 2




Fig. 3






Fig. 4

A **NF- κ B**



control PMA + FB1 + FB1 + FB1 + FB1
alone 0.01 μ M 0.1 μ M 1 μ M 10 μ M

B **AP-1**



control PMA + FB1 + FB1 + FB1 + FB1
alone 0.01 μ M 0.1 μ M 1 μ M 10 μ M

Fig. 4




Fig. 5







Fig. 6

A



B



C







Fig. 6

D



E



F






Fig. 7

Fig. 8

