

JPET/2003/063065

## **Histamine H<sub>1</sub>- and H<sub>2</sub>-Receptor Gene and Protein Levels Are Differentially Expressed in the Hearts of Rodents and Humans**

**NAOYUKI MATSUDA, SUBRINA JESMIN, YOSHIKA TAKAHASHI,  
EIICHIRO HATTA, MASANOBU KOBAYASHI, KAZUTO MATSUYAMA,  
NOZOMI KAWAKAMI, ICHIRO SAKUMA, SATOSHI GANDO, HIYOYUKI  
FUKUI, YUICHI HATTORI, ROBERTO LEVI**

*Departments of Pharmacology (Y.H.), Anesthesiology & Critical Care Medicine (N.M.,  
Y.T., S.G.), Cardiovascular Medicine (S.J., I.S.) and Cardiovascular Surgery (E.H.),  
Hokkaido University School of Medicine, Sapporo, Japan; Division of Cancer Biology,  
Institute for Genetic Medicine, Hokkaido University, Sapporo, Japan (M.K.); Department  
of Pharmacology, Faculty of Pharmaceutical Sciences, University of Tokushima,  
Tokushima, Japan (K.M., N.K., H.F.); and Department of Pharmacology, Weill Medical  
College of Cornell University, New York, New York (R.L.)*

Running title: H<sub>1</sub>- and H<sub>2</sub>-Receptor Expression in the Heart

Corresponding author: Yuichi Hattori, M.D.  
Department of Pharmacology  
Hokkaido University  
School of Medicine  
Sapporo 060-8638, Japan  
Tel: +81-11-706-6920  
Fax: +81-11-706-7378  
E-mail: [yhattori@med.hokudai.ac.jp](mailto:yhattori@med.hokudai.ac.jp)

Pages: 31

Tables: 0

Figures: 12

References: 40

The number of words of Abstract: 244

The number of words of Introduction: 655

The number of words of Discussion: 1,414

**ABBREVIATIONS:** AV, atrioventricular; CHO, Chinese hamster ovary; PAGE, polyacrylamide gel electrophoresis; PCR, polymerase chain reaction; PBS, phosphate-buffered saline; SA, sinoatrial

## ABSTRACT

Histamine is highly concentrated in the heart of animals and humans. Excessive release in pathophysiological conditions, such as immediate hypersensitivity and septic shock, causes cardiac dysfunction and arrhythmias. Previous pharmacological studies revealed that H<sub>1</sub>- and H<sub>2</sub>-receptors mediate these effects. Yet, an accurate estimate of the distribution and molecular characteristics of cardiac histamine receptors is missing. Recently, the genes encoding H<sub>1</sub>- and H<sub>2</sub>-receptors have been cloned, and the amino acid sequence and protein structure have been elucidated. Accordingly, we analyzed gene and protein expression levels of H<sub>1</sub>- and H<sub>2</sub>-receptors in atria and ventricles of guinea pig, rabbit, rat and human hearts. With immunocytochemical techniques we examined the regional expression of H<sub>1</sub>- and H<sub>2</sub>-receptor proteins in the sinoatrial and atrioventricular nodes and surrounding myocardium of the guinea-pig heart. Northern and Western blot studies revealed that cardiac histamine H<sub>1</sub>- and H<sub>2</sub>-receptors are variably distributed among different mammalian species and different regions of the heart, while H<sub>2</sub>-receptors are abundantly expressed in human atrial and ventricular myocardium. These findings agree with those of previous pharmacological studies, clearly demonstrating that the responses of the heart to histamine depend on the expression level of H<sub>1</sub>- and H<sub>2</sub>-receptors. The highly abundant expression of H<sub>2</sub>-receptors in the human heart substantiates histamine arrhythmogenicity in various disease states. The new knowledge of a differential distribution of histamine receptor subtypes in the human heart will foster a better understanding of histamine roles in cardiovascular pathophysiology and may contribute to new therapeutic approaches to histamine-induced cardiac dysfunctions.

The actions of histamine as a cardiac stimulant have been appreciated for more than 90 years since the seminal work of Dale and Laidlaw (1910). The primary and direct actions of histamine in the heart are characterized by an increase in sinus rate and ventricular automaticity, a decrease in AV conduction velocity, and an increase in force of contraction (Levi et al., 1991; Hattori, 1999). Since histamine is present in high concentrations in cardiac tissues in most animal species, including humans (Bristow et al., 1982; Wolff and Levi et al., 1986), its release from cardiac stores and its subsequent actions on the heart may be of importance in pathological conditions associated with histamine release (Levi, 1988). In this context, we recently reported that superinduction of histamine receptor gene products and histamine overproduction contribute to the hemodynamic derangement characteristic of septic shock (Matsuda et al., 2002).

The negative dromotropic and positive chronotropic effects of histamine invariably result from the activation of H<sub>1</sub>- and H<sub>2</sub>-receptors, respectively (Levi et al., 1991; Hattori, 1999). In contrast, there is a marked species difference in the subtype of histamine receptors mediating the positive inotropic effect. Moreover, different receptor subtypes mediate the inotropic effect in different parts of the heart within the same species. In the guinea-pig heart, only H<sub>1</sub>-receptors are responsible for the positive inotropic effect of histamine in left atrium, whereas H<sub>2</sub>-receptors predominantly mediate its positive inotropic effect in ventricle (Steinberg and Holland, 1975; Verma and McNeill, 1977; Hattori et al., 1994). In contrast, in the rabbit heart, H<sub>2</sub>-receptors mediate the positive inotropic effect of histamine in left atrium whereas H<sub>1</sub>-receptors are predominantly involved in the ventricle (Hattori et al., 1988, 1990, 1994). On the other hand, the positive inotropic effect of histamine in the human heart, whether in the atria or in the ventricle, is invariably blocked by cimetidine, and not by pyrilamine, indicating an exclusive H<sub>2</sub>-receptor involvement (Eckel et al., 1982; Du et al., 1993). In yet another

mode, a positive inotropic response in the rat occurs only with very high concentrations of histamine, and is blocked by propranolol or reserpine pretreatment, suggesting the involvement of adrenergic mechanisms (Laher and McNeill, 1980).

Whether such species differences result from a different distribution of H<sub>1</sub>- and H<sub>2</sub>-receptors remains an open question. To date, the distribution of histamine receptors in different mammalian tissues has been studied with selective radioligands for each receptor subtype. Yet, these studies have been hampered by high nonspecific binding and/or significant binding to secondary non-histamine-receptor sites (Chang et al., 1979; Hill and Young, 1980; Rising and Norris, 1985; Foreman et al., 1985; Liu et al., 1992). Thus, many of the radioligands used to label H<sub>1</sub>- and H<sub>2</sub>-receptors appear to be of limited value for an accurate estimate of the distribution and molecular characteristics of cardiac histamine receptors (Hattori et al., 1991, 1994).

In recent years, the genes encoding each of the H<sub>1</sub>- and H<sub>2</sub>-receptor subclasses have been cloned (Yamashita et al., 1991a; Fujimoto et al., 1993; Horio et al., 1993; Gantz et al., 1991b; Traiffort et al., 1995), and the amino acid sequence and structure of their proteins have been elucidated. Thus, the application of molecular biology techniques to the study of histamine receptor subtypes has enabled us to assess by Northern and Western blotting the expression of their mRNAs and proteins in the heart. In the present study, we analyzed the gene and protein expression levels of H<sub>1</sub>- and H<sub>2</sub>-receptors in atria and ventricles from guinea pig, rabbit, rat and human hearts. Furthermore, we used immunocytochemical techniques to examine the regional expression of H<sub>1</sub>- and H<sub>2</sub>-receptor proteins in the sinus node, AV conduction system, and surrounding myocardium of the guinea pig heart.

We report evidence from Northern and Western blot studies that cardiac histamine H<sub>1</sub>- and H<sub>2</sub>-receptors are variably distributed not only among different

mammalian species, but also in different regions of the heart, and that H<sub>2</sub>-receptors are abundantly expressed in human atrial and ventricular myocardium.

## Materials and Methods

### Animal and Human Tissue

Male Hartley guinea pigs (250-350 g), New Zealand White rabbits (2.2-2.5 kg) and Wistar rats (180-220 g) were used in these experiments. The animals were killed by exsanguination under anesthesia with an overdose of pentobarbital (guinea pigs and rabbits) or gaseous diethyl ether (rats), and the hearts and brains were harvested. All animal procedures were reviewed and approved by the Hokkaido University School of Medicine Animal Care and Use Committee.

Right atrial tissue was obtained from patients undergoing aorta-coronary bypass surgery; these patients had no heart failure, or only a moderate form of it, as measured by echocardiography. Also, portions of left ventricle were excised from patients with moderate heart failure during surgical resection of left ventricular aneurysms. The apex of the left temporal lobe of the brain was taken from patients undergoing internal decompression following subarachnoid hemorrhage with severe cerebral edema. Written informed consent was obtained for use of organs in research from patients undergoing cardiac surgery, or in the case of neurosurgery, from their relatives.

### Northern Blot Analysis

Total RNA was extracted from tissues by the guanidinium thiocyanate-phenol-chloroform method with Isogen (Nippon Gene, Toyama, Japan) used routinely in our laboratory (Matsuda et al., 1999). RNA purity was determined by the ratio of optical density measured at 260 and 280 nm ( $OD_{260}/OD_{280}$ ), and RNA quantity was estimated at  $OD_{260}$ .

RNA (30  $\mu$ g per lane) was subjected to electrophoresis on

agarose/formaldehyde gels and then transferred to a Hybond-N<sup>+</sup> nylon membrane (Amersham, Little Chalfont, Buckinghamshire, UK). The membrane was pre-hybridized in pre-warmed rate-enhanced hybridization buffer (Rapid-hyb buffer; Amersham) at 65°C for 60 min. The amino acid sequence of the human histamine H<sub>1</sub>-receptor displays an 81% and 75% homology with the rat and guinea pig receptors, respectively. The comparison of the amino acid sequence of the human histamine H<sub>2</sub>-receptor with that of the rat and guinea pig H<sub>2</sub>-receptors revealed an overall identity of 85% and 86%, respectively. The molecular cloning of the rabbit H<sub>1</sub>- and H<sub>2</sub>-receptors has not been yet reported, but their amino sequences would be predicted to show high homology with those of other rodents. We thus used guinea pig H<sub>1</sub>-receptor cDNA (Horio et al., 1993) and the rat H<sub>1</sub>-receptor cDNA (Fujimoto et al., 1993), both of which were kindly provided by Prof. E. Senba (Department of Anatomy, Wakayama Medical College, Wakayama, Japan). The human H<sub>2</sub>-receptor cDNA (accession no. NM\_022304) was isolated from gel after electrophoretic separation of the products which were amplified by the PCR using two oligodeoxynucleotide primers (sense: 5'-CTCTACCGCATGCAAGATCA-3', and antisense: 5'-CCCCAGGTGGATAGACAGAA-3') (Matsuda et al., 2002). The cDNA probes were radio-labeled using a random primer labeling system (Rediprime; Amersham) in the presence of [<sup>32</sup>P]dCTP (6000 Ci/mmol; Amersham). The blot was washed in 2 x standard saline citrate/0.1% SDS at room temperature for 10 min and in 1 x standard saline citrate/0.1% SDS twice at 65°C for 5 min. The histamine H<sub>1</sub>- or H<sub>2</sub>-receptor mRNA was quantified by counting the radioactivity using a bioimaging analyzer (Fujix BAS 2000; Fuji Photo Film, Tokyo, Japan), as previously described (Matsuda et al., 1999). Ethidium bromide staining was used as a control to verify gel loading, and the expression of H<sub>1</sub>- or H<sub>2</sub>-receptor mRNA was normalized as the ratio of the H<sub>1</sub>- or H<sub>2</sub>-receptor mRNA over 28 S ribosomal RNA.



### Antibody Specificity to Histamine Receptors

The coding regions of the human H<sub>1</sub>-receptor gene and of the human H<sub>2</sub>-receptor gene were subcloned into a pBluescript SK(+) vector (La Jolla, CA). Then, nucleotide sequences that encoded 10-amino acid peptides (EQKLISEEDL) of the human *c-myc* epitope were inserted between the amino-terminal initiator methionine and the second amino acid of the human H<sub>1</sub>- or H<sub>2</sub>-receptor by PCR. PCR was carried out with the following primers using pBluescript vectors containing the human H<sub>1</sub>- or H<sub>2</sub>-receptor gene as template. Primary pairs (forward: 5'-GCGGCCGCATGGAGC-AAAAGCTCATCAGTGAGGAAGACCTAAGCCTCCC CAATTCCTGCCTC-3', and reverse: 5'-GCGGCGGCTAACCTCGCTTATACGTCTTAAGA-3') were used for the H<sub>1</sub>-receptor. Primary pairs (forward: 5'-GCGGCCGCATGGAGCAAAGCTCATCA-GTGAGGAAGACCTAGCACCCAA TGGAACAGCCTCT-3', and reverse: 5'-GCG-GCGGCTAATGGACAGACACCGAGGGACCC-3') were used for the H<sub>2</sub>-receptor. The resulting PCR products were digested and ligated into the pBluescript vectors. The entire coding region, including the epitope tag, was sequenced and transferred into a mammalian expression vector, pdKCR-dhfrs. CHO cells deficient in dihydrofolate reductase were transfected with the plasmid constructs using the calcium phosphate precipitation method (Fujimoto et al., 1993). Cells were cultured in a-minimum essential medium without ribonucleosides and deoxyribonucleosides (Life Technologies, Rockville, MD) supplemented with 10% fetal calf serum. After 2 weeks of incubation, individual colonies were transferred to new plates.

Viewed by immunostaining, CHO cells were confirmed as being successfully

transfected with the plasmid-containing the H<sub>1</sub>- or H<sub>2</sub>-receptor by the expression of *c-myc*. Transfected CHO cells expressing the H<sub>1</sub>- and H<sub>2</sub>-receptor proteins were stained with anti-human H<sub>1</sub>- and H<sub>2</sub>-receptor rabbit polyclonal antibody (Chemicon International, Temecula, CA) employed in this study. However, control cells that were subjected to the transfection procedure with no plasmid present showed no reactivity with these antibodies. The specificity of the histamine receptor antibodies was also certified in transfected cells by Western blot (Fig. 1). Therefore, these data confirmed that anti-human H<sub>1</sub>- and H<sub>2</sub>-receptor antibodies were specific to the H<sub>1</sub>- and H<sub>2</sub>-receptor proteins, respectively.

### **Western Blot Analysis**

Samples of tissue homogenate (20 µg) were run on SDS-PAGE (14% polyacrylamide gel), and electro-transferred to a polyvinylidene difluoride filter membrane. To reduce non-specific binding, the membrane was pre-incubated for 60 min at room temperature in PBS (137 mM NaCl, 2.7 mM KCl, 8.1 mM Na<sub>2</sub>HPO<sub>4</sub>, and 1.5 mM KH<sub>2</sub>PO<sub>4</sub>) containing 1% bovine serum albumin. The membranes were then incubated overnight 4°C with anti-histamine receptor antibody at 1:500 dilution in PBS (10 µg/ml). After extensive washing with in PBS containing 0.05% Tween 20, the membrane was incubated with horseradish peroxidase-conjugated anti-rabbit antibody (1:6000 dilution; Bio-Rad Laboratories, Hercules, CA) for 60 min at room temperature. The blots were washed twice in PBS-Tween buffer and subsequently visualized with an enhanced chemiluminescence detection system (Amersham), exposed to X-ray film, and analyzed by NIH Image software produced by Wayne Rasband (National Institutes of Health, Bethesda, MD). To check for protein loading/transfer variations, all blots were stained with Ponceau red (washable, before incubation with antibodies) and with Coomassie

brilliant blue (permanent, after the enhanced chemiluminescence detection system). Intensity of total protein bands per lane was evaluated by densitometry. Loading/transfer variation between samples was negligible.

### **Immunohistochemistry**

For immunohistochemical determination of histamine receptors, tissue specimens were fixed in 10% buffered-formalin solution, dehydrated, and then embedded in paraffin. The preparations were cut in 5- $\mu$ m sections, de-paraffinized, and treated for 10 min with citrate buffer (10 mM citric acid; pH 6.0) in a microwave oven (750 W) before immuno-staining. Endogenous peroxidase activity was quenched by incubation in 3% hydrogen peroxide for 7 min. After incubation with primary antibodies overnight at 4°C, the sections were washed with PBS and exposed to DAKO EnVision, Peroxidase, Rabbit (DAKO Corporation, Carpinteria, CA) for 30 min at room temperature. Slides were rinsed in PBS, incubated with DAKO Liquid DAB Large Volume Substrate-Chromogen System (DAKO Corporation), rinsed gently with distilled water, and counterstained. Cover slips were mounted with Malinol (Muto Pure Chemicals, Tokyo, Japan). The specificity of immuno-reactivity was confirmed by negative controls in which non-immune IgG was used instead of the primary antibodies.

### **Immunofluorescent Staining**

After overnight incubation with each primary antibody as mentioned above, the sections were exposed to the fluorescent secondary antibody, Cy3-conjugated AffiniPure anti-rabbit IgG or fluorescein-conjugated AffiniPure goat anti-rabbit IgG (Jackson Immuno Research Laboratories, Westgrove, PA), for 2 h according to the manufacturer's instructions. Samples processed without primary antibodies served as negative controls.

Coverslips were mounted with Immunon (Thermo Shandon, Pittsburgh, PA). Immuno-fluorescent images were observed under a Laser Scanning Confocal Imaging System (MRC-1024, Bio-Rad, Hemel Hempstead, UK).

### **Data Analysis**

All values are expressed as mean  $\pm$  S.E.M. of  $n$  observations, where  $n$  represents the number of animals or patients. The results were examined by Student's  $t$  test or one-way ANOVA with repeated measurements followed by Bonferroni's multiple comparison test when appropriate.

## Results

### Histamine H<sub>1</sub>- and H<sub>2</sub>-Receptor Gene Expression

Typical Northern blots for detection of the H<sub>1</sub>-receptor mRNA in cerebral and myocardial tissues from guinea pigs, rabbits, rats and human are shown in Fig. 2. The hybridization with the H<sub>1</sub>-receptor probes revealed two mRNAs for the H<sub>1</sub>-receptor at 3.3 and 3.9 kb. In the guinea pig, the H<sub>1</sub>-receptor transcripts were abundantly expressed in atrial tissue (Figs. 2A and 3A). Thus, the H<sub>1</sub>-receptor mRNA levels in atrium were 86-94% of the cerebral level. Guinea pig ventricular tissue also exhibited a significant presence of the H<sub>1</sub>-receptor transcripts, but the transcript level (48-68% of the cerebral level) was evidently less than that in atrial tissue. In the rabbit, the H<sub>1</sub>-receptor transcripts were barely detectable in atrial tissue (Figs. 2B and 3B). However, the presence of the H<sub>1</sub>-receptor transcripts was evident in ventricle. Its H<sub>1</sub>-receptor mRNA level was 31-43% of the cerebral level. In the rat, abundant expression of the H<sub>1</sub>-receptor mRNA was detected in both atrial and ventricular tissues (75-80% and 72-73% of the cerebral level, respectively) (Figs. 2C and 3C). In the human atrium there was very little expression of H<sub>1</sub>-receptor mRNA (Figs 2D and 3D). Furthermore, the expression level of H<sub>1</sub>-receptor mRNA was markedly lower in the ventricle when compared with the cerebral level (12-26%).

Northern blot analysis using the H<sub>2</sub>-receptor probe revealed a major mRNA species of 3.2 kb in guinea pig and rabbit tissues (Fig. 4, A and B). In rat and human tissues, however, 7 kb H<sub>2</sub>-receptor mRNA was also detected (Fig. 4, C and D). In the guinea pig, H<sub>2</sub>-receptor mRNA expression was much higher in ventricle than in atrium (Figs. 4A and 5A). Its expression level was 26% in atrium and 64% of the cerebral level in ventricle. In contrast, in the rabbit, H<sub>2</sub>-receptor mRNA expression was higher in the atrium than in the ventricle (Figs. 4B and 5B). The levels of the H<sub>2</sub>-receptor mRNA were

66% and 30% of the cerebral level in atrium and ventricle, respectively. In the rat, the H<sub>2</sub>-receptor transcripts were less abundantly expressed in both atrial and ventricular tissues (Figs. 4C and 5C). When compared with the H<sub>2</sub>-receptor transcript level in cerebral tissue, the levels in atrial and ventricular tissues were 17% and 23%, respectively. In the human, a considerable expression level of the H<sub>2</sub>-receptor transcripts was detected in myocardial tissues (Figs. 4D and 5D). Thus, atrial and ventricular tissues showed 68-94% and 43-71% of the H<sub>2</sub>-receptor transcript levels in cerebral tissue, respectively.

### **Histamine Receptor Protein Expression**

Immunological detection of the histamine H<sub>1</sub>-receptor was performed using a anti-human H<sub>1</sub>-receptor polyclonal antibody, which recognized a 57-kDa band in mammalian cerebral and myocardial tissues (Fig. 6). In the guinea pig, the band was evidently lighter in ventricle than in atrium and cerebrum (Fig. 6A). Densitometric quantification of the signal showed that the H<sub>1</sub>-receptor protein level was 86% in atrium and 65% in ventricle when compared with the cerebral level (Fig. 7A). In the rabbit, the atrial band was very faint, and the ventricular band was modest but clearly evident (Fig. 6B). Quantitative analysis revealed that the atrial and ventricular levels of the H<sub>1</sub>-receptor protein were 10% and 38% of the cerebral level, respectively (Fig. 7B). In the rat, the protein bands were intensely labeled in both atrial and ventricular tissues (Fig. 6C). Thus, the relative amount of immunodetectable H<sub>1</sub>-receptor, as determined by densitometric scanning, was more than 70% of the cerebral level in these tissues (Fig. 7C). In the human, the H<sub>1</sub>-receptor protein was much less abundant in myocardial tissues when compared with cerebral tissue (Fig. 6D). Quantitative analysis of immunoblots showed that the amounts of atrial and ventricular H<sub>1</sub>-receptor protein were only 4% and 15% of the cerebral level, respectively (Fig. 7D).

Immunoblot analysis using anti-human H<sub>2</sub>-receptor polyclonal antibody showed a single protein band with a molecular mass of 69 kDa in mammalian cerebral and myocardial tissues (Fig. 8). In the guinea pig, the band was markedly less in atrium than in ventricle and cerebrum (Fig. 8A). Quantification of the H<sub>2</sub>-receptor protein indicated that the protein levels in atrial and ventricular tissues were 21% and 64% of the cerebral level, respectively (Fig. 9A). In the rabbit, the atrial band was apparently darker than the ventricular band, although it was not so marked in comparison with the cerebral band (Fig. 8B). On scanning these bands, the atrial and ventricular levels were 67% and 30% of the cerebral level, respectively (Fig. 9B). In the rat, the H<sub>2</sub>-receptor protein signal was marginally visible in myocardial tissues (Fig. 8C). Densitometric analysis revealed that the H<sub>2</sub>-receptor protein levels in atrial and ventricular tissues were 13% and 15% of the cerebral level, respectively (Fig. 9C). In the human, immunodetectable H<sub>2</sub>-receptor was found at high levels in both atrial and ventricular tissues (Fig. 8D). Compared with the cerebrum, the atrial and ventricular expression levels of the H<sub>2</sub>-receptor protein were 79% and 70%, respectively (Fig. 9D).

As can be seen in Fig. 10, immunofluorescence studies confirmed the results obtained from Western blotting. Thus, immunofluorescence staining for H<sub>1</sub>-receptor protein in the guinea pig heart showed that its expression was high in the atrial tissue and was low in the ventricular tissue. About 83% of cells in the atrium and 50% of cells in the ventricle showed positive immunoreactivity for the H<sub>1</sub>-receptor. In contrast, positive staining for H<sub>2</sub>-receptor protein was strong in the ventricle, but weak in the atrium: 90% of ventricular cells and only 45% of atrial cells showed positive staining for H<sub>2</sub>-receptors.

## **Identification of Histamine Receptors on the Specialized Conductive Tissue of the Heart**

Immunohistochemical studies showed abundant expression of both H<sub>1</sub>- and H<sub>2</sub>-receptor proteins in the SA node in the guinea pig heart (Fig. 11). Furthermore, the AV node appeared to equally express H<sub>1</sub>- and H<sub>2</sub>-receptor proteins (Fig. 12).



## Discussion

Using Northern blotting, immunoblotting, and immunohistochemical techniques we uncovered the presence and distribution pattern of histamine H<sub>1</sub>- and H<sub>2</sub>-receptor mRNA transcripts and proteins in the heart of various mammals, including humans. On Western blots, we detected H<sub>1</sub>- and H<sub>2</sub>-receptor proteins in every mammalian cardiac and brain tissue studied. H<sub>1</sub>- and H<sub>2</sub>-receptor proteins migrated as two single bands at ~57 and 69 kDa, respectively.

Photoaffinity binding studies using [<sup>125</sup>I]iodoazidophenpyramine and subsequent SDS-PAGE analysis had previously indicated that the H<sub>1</sub>-receptor protein has a molecular weight of 56 kDa under reducing conditions in rat, guinea-pig and mouse brain (Ruat and Schwartz, 1989). Similarly, studies in bovine adrenal medulla membranes with the photoaffinity ligand [<sup>3</sup>H]azidobenzamide had found labeled peptides in the size range of 53 to 58 kDa (Yamashita et al., 1991b). Moreover, when the bovine adrenal medulla H<sub>1</sub>-receptor had been expression-cloned in *Xenopus* oocytes, it was deduced to correspond to a 491 amino acid protein with a calculated molecular weight of 56 kDa (Yamashita et al., 1991a). Thus, the molecular weight for the H<sub>1</sub>-receptor estimated in this study is comparable to those noted in the previous reports.

On the other hand, the molecular weight of 69 kDa obtained for the H<sub>2</sub>-receptor is considerably larger than the previously calculated molecular weights (40.2 to 40.5 kDa) for the cloned H<sub>2</sub>-receptors (Ganz et al., 1991a; Ruat et al., 1991; Traiffort et al., 1995; Kobayashi et al., 1996). This difference could be explained by glycosylation of the native H<sub>2</sub>-receptor. Indeed, cloned H<sub>2</sub>-receptor protein carries N-glycosylation sites in the N-terminus region (Ganz et al., 1991a, b; Ruat et al., 1991; Traiffort et al., 1995).

We found that the expression levels of H<sub>1</sub>- and H<sub>2</sub>-receptor proteins in cardiac tissues from each mammalian species correlated with their mRNA expression levels, as

determined by Northern blot analysis. This correlation implies that the expression of H<sub>1</sub>- and H<sub>2</sub>-receptors in the mammalian heart is regulated in a transcriptional manner. Northern blot analysis of cardiac and cerebral tissues from the four mammals, including human, revealed two hybridizing forms of mRNA for the H<sub>1</sub>-receptor, 3.3-kb and 3.9-kb transcripts. These transcripts may correspond to the use of distinct start sites in the 5' region of the gene or distinct polyadenylation sites or, alternatively, to products of distinct genes (Traiffort et al., 1994). Northern blot analysis of H<sub>2</sub>-receptor mRNA identified a single transcript of 3.2 kb in guinea pig and rabbit cardiac and cerebral tissues, whereas the second higher molecular weight species of 7 kb was clearly evidenced in rat and human tissues. While hybridization in most previous studies indicated the presence of a single H<sub>2</sub>-receptor transcript of 4.6-6 kb (Ruat et al. 1991; Traiffort et al., 1995; Karlstedt et al., 2001), it is interesting to note that, although not addressed by the authors, an earlier study had shown the presence of several H<sub>2</sub>-receptor transcripts on Northern blots from various canine tissues (Gantz et al., 1991b).

In guinea pigs, we found H<sub>1</sub>-receptors to be expressed in the heart at a relatively high level, mainly in the atrium and less in the ventricle, compared with their abundant expression in the brain. In contrast, H<sub>2</sub>-receptors were abundant in the ventricle and slightly expressed in the atria. Conversely, in rabbits, H<sub>2</sub>-receptors appeared to be the predominant histamine-receptor subtype in the atrium, with much less H<sub>1</sub>-receptor expression; whereas the ventricle expressed an abundance of H<sub>1</sub>-receptors and much less H<sub>2</sub>-receptors.

Indeed, previous pharmacological studies had indicated a marked difference between guinea-pig and rabbit hearts in terms of which receptor subtypes mediate the positive inotropic effect of histamine in left atrium and ventricle. While the positive inotropic effect of histamine in the left atrium of the rabbit is exclusively mediated by

H<sub>2</sub>-receptors (Hattori et al., 1988), in the guinea pig, the positive inotropic effect in the left atrium is mediated entirely by H<sub>1</sub>-receptors (Steinberg and Holland, 1995; Verma and McNeill, 1977). On the other hand, in guinea pig ventricular muscle, the positive inotropic effect of histamine is mediated predominantly by H<sub>2</sub>-receptors (Hattori et al., 1994), whereas H<sub>1</sub>-receptors almost exclusively mediate the inotropic effect in the rabbit ventricle (Hattori et al., 1990, 1994). Thus, the particular distribution of H<sub>1</sub>- and H<sub>2</sub>-receptors in guinea pig and rabbit myocardial tissues that we report here, well explains previous pharmacological findings.

In the rat heart, H<sub>2</sub>-receptors were present at a rather low level both in the atrium and ventricle; in contrast, the expression of H<sub>1</sub>-receptors was significant in both tissues. Yet, in the rat heart, histamine is known to cause a weak positive inotropic effect and only at very high concentrations. Notably, this inotropic effect is attenuated by  $\beta$ -blockers rather than by H<sub>1</sub>-receptor antagonists (Laher and McNeill, 1980; Levi et al., 1991). Although this suggest that the weak positive inotropic response in the rat heart may result from catecholamine release, it is also conceivable that in the rat heart H<sub>1</sub>-receptors are weakly coupled to their transduction pathways.

In human atrial and ventricular tissue, H<sub>1</sub>-receptor expression was less abundant, whereas H<sub>2</sub>-receptors were richly expressed. This is the first study showing that H<sub>2</sub>-receptors are present as the predominant histamine receptor subtype in the human heart at both mRNA and protein levels, using Northern and Western blot analysis. The present data are also in good agreement with the previous functional reports that histamine produces a positive inotropic effect in human atrial and ventricular muscles likely due to the exclusive activation of H<sub>2</sub>-receptors (Eckel et al., 1982; Levi et al., 1991; Du et al., 1993).

Our immunohistochemical studies clearly show the presence of both H<sub>1</sub>- and

H<sub>2</sub>-receptors in the SA and AV nodes of the guinea-pig heart. Indeed, pharmacological evidence for the presence of H<sub>2</sub>-receptors in the SA node was already strong. In fact, a sinus rate increase in response to histamine had been demonstrated in isolated hearts from guinea pigs, rabbits and cats, and in isolated right atria from guinea pigs, rabbits and monkeys (Levi et al., 1991; Hattori, 1999). In these preparations, the histamine-induced increase in sinus rate was selectively and competitively antagonized by H<sub>2</sub>-receptor antagonists such as cimetidine, indicating that H<sub>2</sub>-receptors which are present in the SA node primarily mediate the positive chronotropic effect of histamine. On the other hand, it is not clear at this time what role, if any, is played by the H<sub>1</sub>-receptors present in the SA node.

In the guinea pig heart, histamine is known to impair AV conduction (Levi, 1972; Levi and Kuye, 1974). This negative dromotropic effect is effectively abolished by H<sub>1</sub>-receptor antagonists and is mimicked by selective H<sub>1</sub>-receptor agonists (Levi et al., 1975), suggesting that H<sub>1</sub>-receptors mediate the histamine-induced slowing of AV conduction. However, histamine may also enhance automaticity in the AV node. Electrophysiological experiments using AV node preparations from the rabbit and guinea pig have indicated that H<sub>2</sub>-receptors promoting an increase in automaticity are likely to exist in the AV node (Borchard and Hafner, 1986; Sanchez-Chapula and Elizalde, 1987). Therefore, our immunohistochemical data demonstrating the presence of both H<sub>1</sub>- and H<sub>2</sub>-receptors in the AV conduction system are in good agreement with the results of previous pharmacological studies.

In conclusion, we provide here for the first time evidence from Northern and Western blot studies that histamine H<sub>1</sub>- and H<sub>2</sub>-receptors are variably distributed not only among different mammalian species, but also in different regions of the heart. Notably, these novel molecular findings agree with previous pharmacological results obtained in

whole animal and isolated tissue experiments. With the exception of the rat heart, where pharmacological responses are apparently dissociated from the abundant expression of H<sub>1</sub>-receptors, and where adrenergic mechanism may also play a role, we find that in guinea pig, rabbit and human heart the expression level of H<sub>1</sub>- and H<sub>2</sub>-receptors clearly determines the functional response to histamine. The fact that a highly abundant expression of H<sub>2</sub>-receptors was found in human atrial and ventricular myocardium, substantiates our previous evidence of histamine arrhythmogenicity in the human heart in various disease states (Levi et al., 1981; Wolff and Levi, 1986; Levi, 1988). Moreover, we recently reported that endotoxin-induced sepsis results in a dramatic increase in the expression of H<sub>1</sub>- and H<sub>2</sub>-receptor and histidine decarboxylase genes in the cardiovascular system, causing a marked increase not only in the production of histamine but also in its effects (Matsuda et al., 2002). These changes may well contribute to the cardiovascular deterioration associated with sepsis in humans. The new knowledge of a differential distribution of histamine receptor subtypes in the human heart will foster a better understanding of histamine roles in cardiovascular pathophysiology and possibly contribute to new therapeutic approaches to histamine-induced cardiac dysfunctions.

## References

Borchard U, and Hafner D (1986) Electrophysiological characterization of histamine receptor subtypes in mammalian heart preparations. *Naunyn-Schmiedeberg's Arch Pharmacol* **334**: 294-302.

Bristow MR, Ginsburg R, and Harrison DC (1982) Histamine and the human heart: the other receptor system. *Am J Cardiol* **49**: 249-251.

Chang RSL, Tran VT, and Snyder SH (1979) Characteristics of histamine H<sub>1</sub>-receptors in peripheral tissues labeled with [<sup>3</sup>H]mepyramine. *J Pharmacol Exp Ther* **209**: 437-442.

Dale HH, and Laidlaw PP (1910) The physiological action of beta-imidazolylethylamine. *J Physiol* **41**: 318-344.

Du XY, Schoemaker RG, Bax WA, Bos E, and Saxena PR (1993) Effects of histamine on porcine isolated myocardium: Differentiation from effects on human tissue. *J Cardiovasc Pharmacol* **22**: 468-473.

Eckel L, Gristwood RW, Nawrath H, Owen DAA, and Satter P (1982) Inotropic and electrophysiological effects of histamine on human ventricular heart muscle. *J Physiol* **330**: 111-123.

Foreman JC, Norris DB, Rising TJ, and Webber SE (1985) The binding of [<sup>3</sup>H]-tiotidine to homogenates of guinea-pig lung parenchyma. *Br J Pharmacol* **86**: 475-482.

Fujimoto K, Horio Y, Sugama K, Ito S, Liu YQ, and Fukui H (1993) Genomic cloning of the rat histamine H<sub>1</sub> receptor. *Biochem Biophys Res Commun* **190**: 294-301.

Gantz I, Munzert G, Tashiro T, Schäffer M, Wang L, and Yamada T (1991a) Molecular cloning of the human histamine H<sub>2</sub> receptors. *Biochem Biophys Res Commun* **178**: 1386-1392.

Gantz I, Schäffer M, DelValle J, Logsdon C, Campbell V, Uhler M, and Yamada T (1991b) Molecular cloning of a gene encoding the histamine H<sub>2</sub> receptor. *Proc Natl Acad Sci USA* **88**: 429-433.

Hattori Y (1999) Cardiac histamine receptors: Their pharmacological consequences and signal transduction pathways. *Methods Find Exp Clin Pharmacol* **21**: 123-131.

Hattori Y, Endou M, Gando S, and Kanno M (1991) Identification and characterization of histamine H<sub>1</sub>- and H<sub>2</sub>-receptors in guinea-pig left atrial membranes by [<sup>3</sup>H]-mepyramine and [<sup>3</sup>H]-tiotidine binding. *Br J Pharmacol* **103**: 1573-1579.

Hattori Y, Gando S, Nagashima N, and Kanno M (1994) Histamine receptors mediating a positive inotropic effect in guinea pig and rabbit ventricular myocardium: Distribution of the receptors and their possible intracellular coupling processes. *Jpn J Pharmacol* **65**: 327-336.

Hattori Y, Nakaya H, Endou M, and Kanno M (1990) Inotropic, electrophysiological and

biochemical responses to histamine in rabbit papillary muscles: Evidence for coexistence of H<sub>1</sub>- and H<sub>2</sub>-receptors. *J Pharmacol Exp Ther* **253**: 250-256.

Hattori Y, Sakuma I, and Kanno M (1988) Differential effects of histamine mediated by histamine H<sub>1</sub>- and H<sub>2</sub>-receptors on contractility, spontaneous rate and cyclic nucleotides in the rabbit heart. *Eur J Pharmacol* **153**: 221-230.

Hill SJ, and Young JM (1980) Histamine H<sub>1</sub>-receptors in the brain of the guinea-pig and the rat: differences in ligand binding properties and regional distribution. *Br J Pharmacol* **68**: 687-696.

Horio Y, Mori Y, Higuchi I, Fujimoto K, Ito S, and Fukui H (1993) Molecular cloning of the guinea-pig histamine H<sub>1</sub> receptor gene. *J Biochem* **114**: 408-414.

Karlstedt K, Senkas A, Åhman M, and Panula P (2001) Regional expression of the histamine H<sub>2</sub> receptor in adult and developing rat brain. *Neuroscience* **102**: 201-208.

Kobayashi T, Inove I, Jenkins NA, Gilbert DJ, Copeland NG, and Watanabe T (1996) Cloning RNA expression and chromosomal location of a mouse histamine H<sub>2</sub>-receptor gene. *Genomics* **37**: 390-394.

Laher IE, and McNeill JH (1980) Effects of histamine on rat isolated atria. *Can J Physiol Pharmacol* **58**: 1256-1261.

Levi R (1972) Effects of exogenous and immunologically released histamine on the



isolated heart: A quantitative comparison. *J Pharmacol Exp Ther* **182**: 227-238.

Levi R (1988) Cardiac anaphylaxis: models, mediators, mechanisms and clinical considerations. In *Human Inflammatory Disease, Clinical Immunology* (Marone G, Lichtenstein LM, Condorelli M and Fauci AS eds) pp 93-105, B. C. Decker Inc., Toronto, Canada.

Levi R, Capurro N, and Lee C-H (1975) Pharmacological characterization of cardiac histamine receptors: Sensitivity to H<sub>1</sub>- and H<sub>2</sub>-receptor agonists and antagonists. *Eur J Pharmacol* **30**: 328-335.

Levi R, and Kuye JO (1974) Pharmacological characterization of cardiac histamine receptors: Sensitivity to H<sub>1</sub>-receptor antagonists. *Eur J Pharmacol* **27**: 330-338.

Levi R, Malm JR, Bowman FO and Rosen MR (1981) The Arrhythmogenic Actions of Histamine on Human Atrial Fibers. *Circ Res* **49**: 545-550.

Levi R, Rubin LE, and Gross SS (1991) Histamine in cardiovascular function and dysfunction: Recent developments. In *Handbook of Experimental Pharmacology, Vol. 97 Histamine and Histamine Antagonists* (Uvnäs B ed) pp 347-383, Springer-Verlag, Berlin.

Liu YQ, Horio Y, Mizuguchi H, Fujimoto K, Imamura I, Abe Y, and Fukui H (1992) Re-examination of <sup>3</sup>H-mepyramine binding assay for histamine H<sub>1</sub>-receptor using quinine. *Biochem Biophys Res Commun* **189**: 378-384.

Matsuda N, Hattori Y, Gando S, Akaishi Y, Kemmotsu O, and Kanno M (1999) Diabetes-induced down-regulation of  $\beta_1$ -adrenoceptor mRNA expression in rat heart. *Biochem Pharmacol* **58**: 881-885.

Matsuda N, Hattori Y, Sakuraya F, Kobayashi M, Zhang X-H, Kemmotsu O, and Gando S (2002) Hemodynamic significance of histamine synthesis and histamine H<sub>1</sub>- and H<sub>2</sub>-receptor gene expression during endotoxemia. *Naunyn-Schmiedeberg's Arch Pharmacol* **366**: 513-521.

Rising TJ, and Norris DB (1985) Histamine H<sub>2</sub>-receptor radioligand binding studies. In *Frontiers in Histamine Research, Advances in the Biosciences* (Ganellin CR, and Schwartz JC ed) pp 61-67, Pergamon, New York.

Ruat M, and Schwartz JC (1989) Photoaffinity labeling and electrophoretic identification of the H<sub>1</sub>-receptor: comparison of several brain regions and animal species. *J Neurochem* **53**: 335-339.

Ruat M, Traiffort E, Arrang JM, Leurs R, and Schwartz JC (1991) Cloning and tissue expression of a rat histamine H<sub>2</sub>-receptor gene. *Biochem Biophys Res Commun* **179**: 1470-1478.

Sanchez-Chapula J, and Elizalde A (1987) Characterization of the effects of histamine on the transmembrane electrical activity of guinea-pig and rabbit SA- and AV-node cells. *Naunyn-Schmiedeberg's Arch Pharmacol* **336**: 218-223.

Steinberg MI, and Holland DR (1975) Separate receptors mediating the positive inotropic and chronotropic effect of histamine in guinea-pig atria. *Eur J Pharmacol* **34**: 95-104.

Traiffort E, Leurs R, Arrang JM, Tardivel-Lacombe J, Diaz J, Schwartz J-C, and Ruat M (1994) Guinea pig histamine H<sub>1</sub> receptor. I. Gene cloning, characterization, and tissue expression revealed by in situ hybridization. *J Neurochem* **62**: 507-518.

Traiffort E, Vizuete ML, Tadvellacombe J, Souile E, Schwartz JC, and Ruat M (1995) The guinea-pig histamine H<sub>2</sub> receptor-gene cloning, tissue expression and chromosomal localization of its human counterpart. *Biochem Biophys Res Commun* **211**: 570-577.

Verma SC, and McNeill JH (1977) Cardiac histamine receptors: Differences between left and right atria and ventricle. *J Pharmacol Exp Ther* **200**: 352-362.

Wolff AA, and Levi R (1986) Histamine and cardiac arrhythmias. *Circ Res* **58**: 1-16.

Yamashita M, Fukui H, Sugama K, Horio Y, Ito S, Mizoguchi H, and Wada H (1991a) Expression cloning of a cDNA encoding the bovine histamine H<sub>1</sub> receptor. *Proc Natl Acad Sci USA* **88**: 11515-11519.

Yamashita M, Ito S, Sugama K, Fukui H, Smith B, Nakanishi K, and Wada H (1991b) Biochemical characterization of histamine H<sub>1</sub>-receptors in bovine adrenal medulla. *Biochem Biophys Res Commun* **177**: 1233-1239.

## Figure Legends

**Fig. 1.** Demonstration of antibody specificity to histamine receptors. CHO cells were transfected with a plasmid containing the human *c-myc* epitope and human H<sub>1</sub>-receptor. Note that the bands with the same size (60 kDa) were detected when Western blot was performed using anti-*c-myc* antibody and anti-H<sub>1</sub>-receptor antibody. Control cells (cells transfected with no plasmid present) exhibited no bands regardless of which antibody was employed. The same results were obtained in cells transfected with the H<sub>2</sub>-receptor instead of the H<sub>1</sub>-receptor.

**Fig. 2.** H<sub>1</sub>-receptor mRNA expression in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D) (representative Northern blots). H<sub>1</sub>-receptor mRNA was detected as two major bands of 3.3 and 3.9 kb using a <sup>32</sup>P-labeled cDNA fragment of the guinea pig (0.50 kb) or rat (0.35 kb) H<sub>1</sub>-receptor as a probe. The locations of ribosomal RNA (28 S and 18 S) are indicated. Ethidium bromide staining was used as a control for gel loading (lower part of each panel).

**Fig. 3.** Quantification of steady-state levels of H<sub>1</sub>-receptor mRNA in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). H<sub>1</sub>-receptor mRNA was normalized as the ratio of its mRNA level over the 18 S level and was expressed as percentage of the value obtained in each cerebral tissue. Values are means  $\pm$  S.E.M. ( $n = 5$ ). For the human, the average of two separate experiments is given.  $P < 0.05$  indicates a significant difference between atrial and ventricular tissues.

**Fig. 4.** H<sub>2</sub>-receptor mRNA expression in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D) (representative Northern blots). H<sub>2</sub>-receptor mRNA was detected as a major single band of 3.2 kb (for guinea pig and rabbit) or two bands of 3.2 and 7 kb (for rat and human) by using a <sup>32</sup>P-labeled cDNA fragment (0.46 kb) of the human H<sub>2</sub>-receptor as a probe. The locations of ribosomal RNA (28 S and 18 S) are indicated. Ethidium bromide staining was used as a control for gel loading (lower part of each panel).

**Fig. 5.** Quantification of the steady-state levels of H<sub>2</sub>-receptor mRNA in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). H<sub>2</sub>-receptor mRNA was normalized as the ratio of its mRNA over the 18 S level and was expressed as percentage of the value obtained in each cerebral tissue. Values are means  $\pm$  S.E.M. ( $n = 5$ ). For the human, the average of two separate experiments is given.  $P < 0.05$  indicates a significant difference between atrial and ventricular tissues.

**Fig. 6.** Representative Western blots for the H<sub>1</sub>-receptor protein in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). The antibody recognized a single protein of 57 kDa referred to as H<sub>1</sub>-receptor.

**Fig. 7.** Summary of immunoblot analysis of the H<sub>1</sub>-receptor protein in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). Densitometric results are expressed as percentage of the 57-kDa band

obtained with cerebrum in each experiment. Values are means  $\pm$  S.E.M. ( $n = 5$ ).  $P < 0.05$  indicates a significant difference between atrial and ventricular tissues.

**Fig. 8.** Representative Western blots for the H<sub>2</sub>-receptor protein in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). The antibody recognized a single protein of 69 kDa referred to as H<sub>2</sub>-receptor.

**Fig. 9.** Summary of immunoblot analysis of the H<sub>2</sub>-receptor protein in cerebral and myocardial (atrium and ventricle) tissues from the guinea pig (A), rabbit (B), rat (C) and human (D). Densitometric results are expressed as percentage of the 69-kDa band obtained with cerebrum in each experiment.  $P < 0.05$  indicates a significant difference between atrial and ventricular tissues.

**Fig. 10.** Immunofluorescent findings for H<sub>1</sub>-receptors (*red*) and H<sub>2</sub>-receptors (*green*) in the guinea pig brain and heart. A, H<sub>1</sub>-receptor immunoreactivity in the cerebellum used as a positive control. B, Immunofluorescence labeling for H<sub>1</sub>-receptors was relatively weak in the ventricle. C, Intense H<sub>1</sub>-receptor staining was detected in the atrium. D, Secondary antibody used for H<sub>1</sub>-receptor labeling showed no immunoreactivity with the atrium. E, H<sub>2</sub>-receptor immunoreactivity in the cerebrum used as a positive control. F, Intense H<sub>2</sub>-receptor staining was found in the ventricle. G, Weak H<sub>2</sub>-receptor immunoreactivity was seen in the atrium. H, Secondary antibody used for H<sub>2</sub>-receptor labeling showed no reactivity with the ventricle. Magnification, x400.

**Fig. 11.** Immunohistological localization of H<sub>1</sub>- and H<sub>2</sub>-receptors in the SA node of the

guinea pig heart. Arrows indicate the SA node area. Positive staining (*brown*) for expression of H<sub>1</sub>-receptors (central panels) and H<sub>2</sub>-receptors (right panels) was demonstrated in the SA node. Left panels show hematoxylin and eosin (HE) stained sections adjacent to those in central and right panels. Magnification, x80 for upper panels, x400 for lower panels.

**Fig. 12.** Immunohistological localization of H<sub>1</sub>- and H<sub>2</sub>-receptors in the AV node of the guinea pig heart. Arrows indicate the AV node area. Positive staining (*brown*) for expression of H<sub>1</sub>-receptors (central panels) and H<sub>2</sub>-receptors (right panels) was demonstrated in the AV node. Left panels show hematoxylin and eosin (HE) stained sections adjacent to those in central and right panels. Magnification, x80 for upper panels, x200 for lower panels.






Figure 1




### A Guinea Pig



### B Rabbit



### C Rat



### D Human




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9






Figure 10




Figure 11




Figure 12