

**Role of $\alpha 7$ Nicotinic Acetylcholine Receptors in the Pressor Response to
Intracerebroventricular Injection of Choline: Blockade by Amyloid Peptide A β 1-42**

Xinyu D. Li and Jerry J. Buccafusco

Department of Pharmacology and Toxicology, Alzheimer's Research Center, Medical College
of Georgia, and the Veterans Administration Medical Center, Augusta, GA 30912, U.S.A.

JPET #63321

Running Title: Amyloid peptide, nicotinic receptors, choline, blood pressure, heart rate, intracerebroventricular.

Address correspondence to:

Jerry J. Buccafusco, Ph.D., Director

Alzheimer's Research Center

Medical College of Georgia

Augusta, Georgia 30912-2300

U.S.A.

Ph: (706) 721-6355 FAX: (706) 721-9861

email: jbuccafu@mcg.edu

No. Text Pages	32
Tables	1
Figures	6
References	53
Abstract Words	282
Introduction Words	711
Discussion Words	1466

Abbreviations:

amyloid beta peptide (1-42)	A β 1-42
dihydro- β -erythroidin	DH β E
intracerebroventricular	icv
methyllaconitine	MLA
nicotinic acetylcholine receptor	α 7nAChR

JPET #63321

Abstract

Systemic blood pressure and cardiac function have long been known to be under the control of central autonomic and hormonal pathways that, in part, utilize cholinergic neural systems. Recently choline, a precursor and product of acetylcholine metabolism, has been shown to serve as a selective endogenous agonist for the $\alpha 7$ subtype of the nicotinic acetylcholine receptor ($\alpha 7$ nAChR). This receptor subtype mediates several responses to nicotine in animals, most notably, neuroprotection and enhanced cognition. The purpose of this study was to determine whether the cardiovascular changes induced by central injection of choline in rats also were mediated by $\alpha 7$ nAChRs. Moreover, we sought to determine whether these cardiovascular changes to choline could be blocked by central pretreatment with A β 1-42, a neurotoxic component of cerebral amyloid that is known to bind with high affinity to $\alpha 7$ nAChRs. Central, intracerebroventricular (icv) injection of choline (50, 100 or 150 μ g) produced dose-dependent (10-15 min duration) pressor response of up to about 20 mmHg. The most consistent change in heart rate included a brief increase (up to 40 beats/min) that lasted 2-3 min, followed by a prolonged decrease averaging 50 beats/min that lasted up to 30 min. Pretreatment (icv) with the selective $\alpha 7$ nAChR antagonists α -bungarotoxin and methyllycaconitine (MLA) significantly inhibited the pressor and heart rate responses to subsequent injection of choline. Pretreatment with the non- $\alpha 7$ -preferring antagonist dihydro- β -erythroidin (DH β E) was not effective. These findings suggested that the cardiovascular response to icv injection of choline was mediated at least in part through $\alpha 7$ nAChRs. Pretreatment (30 min) with low doses (1 – 100 pmoles) of amyloid peptide A β 1-42 (but not with A β 40-1) administered by the icv route significantly inhibited the choline-induced blood pressure increase as well as the choline-induced decrease in heart rate.

JPET #63321

One of the most prominent pathological changes associated with the Alzheimer's disease (AD) is the loss of cholinergic innervation of the cortex and hippocampus. This neuronal loss is accompanied by a decrease in the brain density of the $\alpha 7$ subtype of the nicotinic acetylcholine receptor ($\alpha 7$ nAChR). The $\alpha 7$ nAChR, is a unique member of the nicotinic receptor family which in recent years has been shown to enjoy significant physiological importance. In particular, this receptor subtype is known to mediate several responses to nicotine administration in animals, including neuroprotection and cognitive enhancement. For example, $\alpha 7$ nAChR preferring agonists, such as anabaseine and GTS-21, were shown to improve working memory in rats (Meyer et al., 1994; Arendash et al., 1995; Briggs et al., 1997). Another $\alpha 7$ nAChR agonist, ARR 17779, was shown to reverse the working memory deficits induced by fimbria-fornix lesions (Levin et al., 1999). However, other than procedures that estimate aspects of behavior, no objective measure of central $\alpha 7$ nAChR activation *in vivo* has been described.

The peripheral circulation and cardiac function, to varying degrees and under specific conditions, are continuously regulated through the output of central autonomic and hormonal pathways. Although many important questions remain unanswered, a great level of understanding has been achieved concerning the central control of blood pressure. In this complicated and interconnected regulatory system, central cholinergic neural pathways have been shown to play important roles; and in unanesthetized rats, cholinergic agonists, both muscarinic (see, Brezenoff, 1982; Buccafusco, 1992) and nicotinic (Brezenoff and Giuliano, 1982; Buccafusco and Yang, 1993; Khan et al., 1994), have been shown to induce increases in arterial blood pressure. Changes in heart rate often are unpredictable, depending upon the particular agonist, and upon ongoing arousal and activity (Magri and Buccafusco, 1988). Some years ago, Caputi

JPET #63321

and Brezenoff (1980) demonstrated that central intracerebroventricular (icv) injection of choline induced a short lasting, but significant increase in arterial pressure and a much longer lasting bradycardic response in unanesthetized rats. They failed to inhibit the pressor response through pretreatment of the animals by icv injection with either the muscarinic antagonist atropine or the nicotinic antagonist mecamylamine. Neither drug alone nor their combination modified the pressor response. However, neither of these compounds is subtype selective within their respective class, and atropine induces acetylcholine release. In light of the recently discovered ability of choline to serve as a selective agonist at $\alpha 7$ nAChRs (Papke et al., 1996; Alkondon et al., 1999; Pereira et al., 2002), it is reasonable to consider that this receptor subtype mediates the choline-induced pressor response. With reasonably subtype selective nAChR antagonists now available this possibility can readily be examined. Thus, the first portion of this study was directed at determining whether the cardiovascular response to icv administration of choline was mediated predominantly through activation of $\alpha 7$ nAChRs.

In the Alzheimer's disease (AD) brain large deposits of amyloid plaques are present that consist chiefly of amyloid β peptides. In fact, recently, amyloid β peptides containing a crucial binding sequence, including the neurotoxic A β 1-42, have been shown to bind with high affinity to $\alpha 7$ nAChRs (Wang et al., 1999; Wang et al., 2000). Moreover, A β and $\alpha 7$ nAChRs appear to co-localize within the AD brain. A β peptides have been shown to inhibit the cellular responses to nAChR agonists (Pettit, et al., 2000; Liu, et al., 2001) and to prevent the cytoprotective action of nicotine *in vitro* (Li and Buccafusco, 2003). There also exists evidence to suggest that amyloid deposition in AD-related transgenic mouse strains alters the expression of nicotinic receptors *in vivo* (Bednar, et al., 2002). Whereas A β peptides have been directly introduced into the

JPET #63321

cerebrospinal fluid of rodents by prolonged icv administration to elevate brain amyloid and to produce neurotoxicity, regimens specific for pharmacological interactions between the peptide and $\alpha 7$ nAChRs *in vivo* have not been determined. Therefore, the second portion of this study was designed to assess the ability of A β peptides to inhibit the magnitude and expression of the acute pressor response to choline. In addition to the practical application of the study, these results may provide further *in vivo* evidence for the ability of A β peptides to target $\alpha 7$ nAChRs, a subtype which is highly expressed on neural cells known to be selectively vulnerable to the toxic consequences of AD (Davies and Feisullin, 1981; Sugaya, et al., 1990; Hellstrom-Lindahl, et al., 1999).

JPET #63321

Material and Methods

Materials

Choline iodide, methyllycaconitine citrate (MLA), atropine sulfate, amyloid β peptide 1-42, amyloid β peptide 40-1, and α -bungarotoxin were obtained from Sigma Chemicals, St. Louis, MO. Dihydro- β -erythridine (DH β E) was donated by Merck & Co., Inc, Rahway, NJ. The doses used in this study for each nicotinic receptor antagonist were derived from previous studies in which they were shown to exert nicotinic receptor blocking actions in vivo with relative selectivity (Buccafusco and Yang, 1993; Jonnala et al., 2002). The β amyloid peptides were dissolved in sterile distilled water at 100 μ M, and they were stored as aliquots at -20°C. They were incubated at 37°C for 24 hr before use to allow for aggregation.

Experimental Animals:

Male Wistar rats (Harlan Sprague-Dawley, Indianapolis, IN), weighing 280 – 380 g were housed in an environmentally controlled room on a 12-hr/12-hr day/night cycle and they were maintained on standard rat chow and tap water (unlimited). All animal protocols were previously approved by the Medical College of Georgia Committee on Animal Use for Research and Education.

Surgical Procedures:

Rats were anesthetized initially with methohexital (sodium), 65 mg/kg, i.p. and supplemented as needed. The subject's head was positioned in a stereotaxic frame with the skull in a horizontal position. A midline incision was made to expose the skull. Using stereotaxic landmarks a burr hole was placed over the left lateral cerebral ventricle. An additional burr hole was placed to the

JPET #63321

right of the first hole and a stainless steel anchoring screw was inserted. A sterile stainless steel or cannula guide was attached to a micromanipulator on the stereotaxic frame. Each guide (Plastics One, Roanoke, VA) had a stainless steel shaft equivalent to 22 gauge tubing which was inserted into a wider threaded region made of Teflon at the top of the guide. The threaded region permits the introduction of a screw cap and dummy cannula to plug the guide when not in use. The cannula guide (11 mm long) was lowered to a depth of 4.0 mm below the skull through the burr hole over the ventricle. The tip of the guide was placed over, but without penetrating, the ventricular space. While in position, the guide and screw were covered with acrylic cement. After complete hardening of the cement, the manipulator was retracted and the guide plugged and capped. The skin was pulled around (but not over) the guide and sutured closed.

One week later rats again were anesthetized and a surgical incision was made in the midline of the abdomen to expose the lower portion of the abdominal aorta. A small nick was made in the lower portion of the aorta and a telemetry catheter (implant model TA11PA-C40, Data Sciences International, St. Paul, MN) was introduced and advanced until the tip rested below the origin of the renal arteries. The tip of the catheter contains a non-thrombogenic gel to prevent clot formation. The catheter end was fixed in place in the aorta by using Vetbond tissue adhesive (3M Animal Care Products, St. Paul, MN). The transmitter end of the catheter was placed subcutaneously over the abdomen and the wound was sutured closed.

Measurement of mean arterial pressure and heart rate:

Blood pressure and heart rate were measured using the DSI PhysioTel[®] telemetry system (Data Sciences International, St. Paul, MN), which monitored the animals while they moved freely

JPET #63321

within their cages. Each implanted transducer/transceiver transmitted pre-calibrated digitized blood pressure data via radio frequency signals to a nearby receiver. The data was sampled continuously, and by using the Dataquest[®] data acquisition system (Data Sciences International, St. Paul, MN) which also calculated heart rate continuously from the pressure waveforms. Mean arterial pressure (MAP) and heart rate values were averaged and presented graphically in 30 sec bins.

Microinjections:

For icv injections, a 50 μ l Hamilton micro-syringe (Reno, NV) was connected to a 28 gauge internal cannula (Plastics One, Inc., Roanoke, VA), which was inserted into the cranial 22 gauge cannula guide so that the tip rested within the lateral ventricle. All test compounds were dissolved in 10 μ l sterile normal saline (vehicle), and they were delivered by hand over a period of 15-20 sec. At the completion of experiments each injection site was verified by injecting 10 μ l India ink. The pretreatment drugs (or vehicle) were administered 10 or 30 min (as indicated) prior to the choline (or vehicle) injection. Rats were allowed to recover for at least one day after each experiment (i.e., one choline microinjection). Each rat was used in no more than 5 experiments.

Statistical analysis:

The data depicted in the text and figures is presented as mean \pm S.E.M. The existence of a significant difference between or among experimental groups was determined by analysis of variance with repeated measures. A modified *t* test with Bonferroni's correction for multiple comparisons using the error mean-square term from the analysis of variance was used for

JPET #63321

analysis between groups of data. The criterion for statistical significance was $P < 0.05$ for all comparisons.

JPET #63321

Results

Because all of the data derived from this study are presented as the absolute change from resting (baseline) MAP and heart rate, Table 1 provides the baseline means for each of the experimental paradigms. There were no significant differences among the means for MAP [(F(10,65)=1.25, P=0.28)]. There also were no significant differences among means for resting heart rate [(F(10,65)=0.50, P=0.89)]. Thus, any change noted below in the magnitude or duration of the choline induced cardiovascular responses produced by pretreatment drugs could not be attributed to alterations in baseline (pre-choline) MAP or heart rate.

Microinjection of 10 μ l of vehicle (sterile saline) on average produced only a brief (<3 min) increase in mean arterial pressure (MAP) of up to 3 mmHg. Microinjection of solutions of choline (50, 100, or 150 μ g) within the same volume produced immediate and dose-dependent increases in MAP ranging from 10 to 20 mmHg (Fig. 1). There was a significant between dose component of the ANOVA [F(2,10)=5.58, P=0.024]. The responses peaked 3-4 min after injection and returned to pre-injection levels within 10-15 min. In some animals, MAP declined below preinjection values towards the end of the observation period. Simultaneously, heart rate initially increased (to about 40 beats/min) and it peaked about the same time as the increase in MAP (Fig. 2). Thereafter, heart rate declined to below pre-injection values. The late decrease in heart rate may reflect a decrease in cardiac output driving the late fall in MAP observed in some animals. Although the heart rate changes were not statistically dose-dependent [F(2,10)=1.67, P=0.24], the profile of response to choline was quite similar to that reported 13 years ago by Caputi and Brezenoff (1980) except for the brief initial increase in heart rate which they did not report observing. It is possible that the increase in heart rate may have reflected

JPET #63321

animal activity just after the injection as it was not present in all animals. As observed by Caputi and Brezenoff (1980), and in this study, the duration of the decrease in heart rate usually far exceeded (at least up to 30 min) the duration of the increase in MAP. Also, soon after choline administration, an increase in locomotor activity was often observed, with the effect quickly diminishing; and the rats actually remained quiet, possibly sedate for the remainder of the observation period. Although the behavioral effects were not quantified, they were not apparent after icv injection of vehicle.

To determine whether $\alpha 7$ nAChRs contribute to the cardiovascular response produced by choline, rats were pretreated by icv injection either with methyllycaconitine (MLA) or α -bungarotoxin, both $\alpha 7$ nAChR-specific antagonists. In the first series, administration of 10 μ g MLA (10 min prior to 150 μ g of choline) resulted in a complete abolition of the pressor response to choline. Likewise, the biphasic heart rate change after choline was essentially normalized by MLA pretreatment (Fig. 3). In contrast, pretreatment with 0.3 μ g atropine failed to significantly modify the pressor response to choline. This icv dose of atropine was shown previously to completely block the pressor response to central microinjection of muscarinic agonists (Caputi and Brezenoff, 1980). Atropine pretreatment partly, but significantly, reversed the changes in heart rate produced by choline. The administration of 10 μ g α -bungarotoxin (30 min prior to 150 μ g of choline), like MLA, significantly inhibited the magnitude of the choline-induced pressor response. These data are presented in Figure 4. The longer pretreatment time for α -bungarotoxin in this series was in consideration of the drug's slow binding kinetics. Toxin pretreatment significantly inhibited the choline-induced initial increase in heart rate, but it had no effect on the subsequent bradycardic response (Fig 4, inset). To further examine the specificity

JPET #63321

of the choline response, the β 2-subtype preferring antagonist DH β E was used as the pretreatment agent. Unlike MLA and α -bungarotoxin, pretreatment with 6 μ g of DH β E failed to significantly inhibit the pressor response to subsequent injection of 150 μ g of choline (Fig. 4). The heart rate changes induced by choline also were not affected by DH β E (Fig. 4, inset).

With the data suggesting that the initial pressor response to icv microinjection of choline was mediated predominantly by α 7nAChRs, we next sought to determine whether pretreatment with the amyloid peptide (also known to block α 7nAChRs) would effectively inhibit the choline response. A β 1-42 (1, 10, or 100 pmol) was administered 30 min prior to choline. The peptide itself elicited no change in resting MAP or heart rate (Table 1). The two higher doses of A β 1-42 (10 pmol and 100 pmol) significantly inhibited the increases in MAP produced by icv injection of 150 μ g of choline (Fig. 5). Under the same conditions the reverse peptide A β 40-1 (100 pmol) failed to significantly affect the pressor response to choline (Fig. 6, upper panel). As with the receptor antagonists, the higher dose of the A β 1-42 (100 pmol) normalized the biphasic heart rate response to icv injection of choline, while the same dose of the reverse peptide did not (Fig 6, lower panel).

JPET #63321

Discussion

$\alpha 7$ nAChRs have been suggested to participate in important higher brain functions such as working memory (Addy et al., 2003), in cell survival and neuroprotection (Li et al., 2000; Shimohama et al., 2001; Jonnala and Buccafusco, 2001), and to serve as a potential drug target in disease states such as schizophrenia (Marutle et al., 2001; Araki et al., 2002), epilepsy (Gil et al., 2002), Alzheimer's disease (Meyer et al., 1997; Briggs et al., 1997; Kem, 2000), and Parkinson's disease (Burghaus et al., 2003). As new $\alpha 7$ nAChR compounds become available it would be advantageous to have an efficient, objective, and experimentally simple model for drug efficacy. In this study, the most technically challenging procedure was the installation of the aortic blood pressure telemetry catheter. However, similar results may be obtained by using a standard catheter inserted into any major artery. In fact, for our very first dose finding experiments we directly catheterized the iliac artery and exteriorized the catheter to a water-tight swivel for the measurement of MAP in unanesthetized animals as we have described previously (Buccafusco and Yang, 1993). The use of the telemetry catheter was somewhat more advantageous as there are no exterior catheters to maintain, and the preparation has a longer useful lifetime.

The most reproducible response produced by icv microinjection of choline was the pressor response that occurred during the first 10-15 min after injection. Except for the magnitude of the pressor response (which was greater than that to choline), the cardiovascular changes to icv administered nicotine as we described them several years ago (Buccafusco and Yang, 1993) were quite similar to those produced by icv administration of choline. The maximal pressor response to nicotine (averaging about 30 mmHg) was achieved with 50 μ g (0.11 μ moles) of the drug, as

JPET #63321

compared with 150 μg (0.65 μmoles) of choline. This 6 fold difference in potency was somewhat surprising in view of the 50 fold difference in binding potencies (vs. [^{125}I] α -bungarotoxin) between the two drugs for the $\alpha 7\text{nAChR}$ (Jonnala et al., 2003). However, Alkondon and colleagues (1999; 2000), in studying $\alpha 7$ -mediated currents recorded from CA1 interneurons in rat hippocampus, reported that the EC_{50} values for choline and nicotine were, respectively, 2270 mM and 158 mM, representing only a 14 fold difference in potencies. Likewise there was no more than a 10 fold difference in potency between choline and nicotine in their ability to increase intracellular Ca^{2++} in cultured porcine superior cervical ganglion cells (Si and Lee, 2002). We had noted similar mismatches between binding affinity and pharmacological response earlier in studying nicotine and choline as cytoprotective agents in vitro (Jonnala et al., 2003), so it is not likely that the differences in drug effectiveness could be attributed to differences in their brain distribution or elimination. Also, choline appears to be slightly less efficacious than nicotine though it has been characterized as a full $\alpha 7\text{nAChR}$ agonist (Pereira et al., 2002). Again, we had noted this potential discrepancy in our earlier cytoprotection study.

Although the effect of icv choline on heart rate was more variable than the effect on MAP, the responses appeared to be susceptible to the same antagonists as was the initial increase in MAP. MLA pretreatment almost completely normalized the biphasic change in heart rate to choline; whereas α -bungarotoxin pretreatment appeared to inhibit only the initial tachycardic phase. The differential effect may be due to differences in drug distribution between the two antagonists, and it suggests that the two phases of the heart rate response may be mediated in different brain regions. The ability of icv-injected atropine to inhibit the heart rate changes to choline suggests

JPET #63321

that both muscarinic and nicotinic receptors play a role within the pathway mediating the bradycardic response to choline. The lack of effectiveness of atropine on the increase in blood pressure to choline indicates that the pressor pathway may not include muscarinic receptors.

This model not only provides a useful means for assessing the effectiveness of $\alpha 7$ nAChR agonists and antagonists, but the results suggest that this subtype of nicotinic receptors plays some role in central cardiovascular regulation. For example, we noted earlier that central nicotinic receptors partly mediate the pressor responses to icv administration of certain neuro-active peptides such as bradykinin, angiotensin II, and substance P (Buccafusco and Serra, 1985; Trimarchi et al., 1986). More recently we reported that in rat models genetically manipulated to maintain an elevated blood pressure, there is a link between decreased expression of brain $\alpha 7$ nAChRs and working memory deficits (Gattu et al., 1997; Terry et al., 2001). In the present study the pressor response to icv-administered choline was blocked by pretreatment with the $\alpha 7$ subtype selective antagonists, MLA and α -bungarotoxin. The lack of effect of pretreatment with DH β E, the $\beta 2$ subtype-preferring antagonist, supported the concept that the initial pressor response to choline was mediated by the $\alpha 7$ subtype. Of course the participation of other non- $\alpha 7$ containing subtypes cannot be completely ruled out at this time; but the data do fit the known specificity of choline for $\alpha 7$ nAChRs (Papke et al., 1996; Pereira et al., 2002; Jonnala et al., 2003). Additional support may be derived from the finding that an analog of choline, CDP-choline, also increased blood pressure and decreased heart rate after icv microinjection in rats (Savci et al., 2002). The pressor effect was significantly reduced following pretreatment with the non-selective nicotinic antagonist, mecamylamine.

JPET #63321

Having characterized the specificity of the receptors mediating the pressor response to choline, it was then possible to determine whether icv administration of A β 1-42 would inhibit the subsequent pressor response to icv-injected choline. Cerebroventricular infusion, or repeated icv injection of A β 1-42 in rats has been used to assess the various pharmacological responses induced by the peptide, particularly regarding its potential amnestic actions (Maurice et al., 1996; Yamada et al., 1998; Yamaguchi et al., 2001; Nakamura et al., 2001; Olariu et al., 2002). Moreover, the related peptide A β 25-35 (15 nmol/injection) has been reported to impair short-term working memory even after a single acute injection (Stepanichev et al., 2003). The dose used in the memory experiment was 150 fold greater than the most effective dose we used to block the pressor response to choline.

The mechanism by which A β 1-42 interferes with the performance of certain memory-related tasks is not clear, but the peptide has this feature in common with the nicotinic receptor antagonist mecamylamine (Elrod and Buccafusco, 1991; Hiramatsu et al., 1998; Levin and Simon, 1998). Future studies will be required to determine whether the amnestic effects induced following central microinjection of A β peptides are related to their ability to bind to α 7nAChRs. Nevertheless based on the data presented here, it is clear that A β 1-42 does produce a functional blockade of α 7nAChRs at low pharmacologic doses. The specificity of the A β 1-42 inhibition of the choline pressor response was supported by the failure of A β 40-1 peptide (which contains the critical α 7nAChR binding sequence in reverse order) to block the choline response. Although A β 1-42 significantly inhibited the pressor response to choline, the maximal degree of inhibition amounted to about a 60% decrease relative to the control response. Clearly the nicotinic antagonists were more effective, nearly eliminating the choline-induced pressor response. This

JPET #63321

difference might reflect differences in the nature of the binding kinetics between A β 1-42 and the antagonists. Alternatively, A β 1-42 might not be distributed throughout the brain as readily as MLA and α -bungarotoxin. Indeed, the potential site of action for choline in eliciting its pressor response is not known. Stimulation of specific hypothalamic sites with cholinergic drugs has been reported to generate a pressor response mediated via nicotinic receptors (Schaeppi, 1967; Day and Roach, 1977). A hypothalamic site of action would explain the rapid onset of the pressor response to choline as the drug would readily redistribute to periventricular hypothalamic regions. To support a hypothalamic site of action, the pressor response to icv injection of CDP-choline was partly mediated via the release of vasopressin into the circulation (Savci et al., 2002). Also, α -bungarotoxin binding sites are relatively highly expressed in periventricular hypothalamic regions encompassing the dorsomedial, ventromedial, posterior hypothalamic nuclei, the dorsal premammillary nucleus and the arcuate nucleus (Gattu et al., 1997; Tohyama and Takatsuji, 1998).

In summary, these data support the concept that the pressor response to icv microinjection of choline may be used as an unbiased measure of α 7nAChR stimulation *in vivo*. They also support the ability of acute injection of A β 1-42 to produce a rapid functional blockade of α 7nAChRs at pharmacologically (and likely physiologically in the case of AD) relevant doses. The use of telemetric catheters for monitoring cardiovascular status permits free movement of the animals and could allow for the assessment of various parameters of locomotor activity and other behaviors. Finally, some consideration should be given to the possibility that amyloid A β - α 7nAChR interactions may play a role in the dysautonomia exhibited by some individuals with AD (Giubilei et al., 1998).

JPET #63321

Acknowledgements

The authors are grateful to Laura C. Shuster, who provided excellent technique support.

JPET #63321

References

Addy NA, Nakijama A, Levin ED (2003) Nicotinic mechanisms of memory: effects of acute local DH β E and MLA infusions in the basolateral amygdala. *Brain Res Cogn Brain Res* **16**:51-57.

Alkondon M, Pereira EFR, Eisenberg HM, Albuquerque EX (1999) Choline and selective antagonists identify two subtypes of nicotinic acetylcholine receptors that modulate GABA release from CA1 interneurons in rat hippocampal slices. *J Neurosci* **19**:2693-2705.

Alkondon M, Pereira EFR, Almeida LEF, Randall W, Albuquerque EX (2000) Nicotine at concentrations found in cigarette smokers activates and desensitizes acetylcholine receptors in CA1 interneurons of rat hippocampus. *Neuropharmacology* **39**:2726-2739.

Araki H, Suemaru K, Gomita Y (2002) Neuronal nicotinic receptor and psychiatric disorders: functional and behavioral effects of nicotine. *Jpn J Pharmacol* **88**:133-138.

Bednar I, Paterson D, Marutle A, Pham TM, Svedberg M, Hellstrom-Lindahl E, Mousavi M, Court J, Morris C, Perry E, Mohammed A, Zhang X, Nordberg A (2002) Selective nicotinic receptor consequences in APP(SWE) transgenic mice. *Mol Cell Neurosci* **20**:354-365.

Briggs CA, Anderson DJ, Brioni JD, Buccafusco JJ, Buckley MJ, Campbell JE, Decker MW, Donnelly-Roberts D, Elliott RL, Holladay MW, Hui Y-H, Jackson WJ, Kim DJB, Marsh KC,

JPET #63321

O'Neill A, Prendergast M A, Ryther KB, Sullivan JP, Arneric SP (1997) Functional characterization of the novel neuronal nicotinic acetylcholine receptor ligand GTS-21. In vitro and in vivo. *Pharmacol Biochem Behav* **57**:231-241.

Buccafusco JJ (1992) Neuropharmacologic and behavioral actions of clonidine: Interactions with central neurotransmitters. *Int Rev Neurobiol* **33**:55-107.

Buccafusco JJ, Serra (1985) Role of cholinergic neurons in the cardiovascular responses evoked by central injection of bradykinin or angiotensin II in conscious rats. *Eur J Pharmacol* **113**:43-51.

Buccafusco JJ, Yang XH (1993) Mechanism of the hypertensive response to central injection of nicotine in conscious rats. *Brain Res Bull* **32**:35-41.

Brezenoff HE, Giuliano R (1982) Cardiovascular control by cholinergic mechanisms in the central nervous system. *Annu Rev Pharmacol Toxicol* **22**:341-381.

Burghaus L, Schutz U, Krempel U, Lindstrom J, Schroder H (2003) Loss of nicotinic acetylcholine receptor subunits alpha4 and alpha7 in the cerebral cortex of Parkinson patients. *Parkinsonism Relat Disord* **9**:243-246.

Caputi AP, Brezenoff HE (1980) Cardiovascular effects produced by choline injected into the lateral cerebral ventricle of the unanesthetized rat. *Life Sci* **26**:1029-1036.

JPET #63321

Davies P, Feisullin S (1981) Postmortem stability of α -bungarotoxin binding sites in mouse and human brain. *Brain Res* **216**:449-454.

Day MD, Roach AG (1978) Cardioascular effects of carbachol and other cholinomimetics administered into the cerebral ventricles of conscious cats. *Clin Exp Pharmacol Physiol* **4**:431-442.

Elrod K, Buccafusco JJ (1991) Correlation of the amnestic effects of nicotinic antagonists with inhibition of regional brain acetylcholine synthesis in rats. *J Pharmacol Exp Ther* **258**:403-409.

Gattu M, Pauly JR, Boss KL, Summers JB, Buccafusco JJ (1997) Cognitive impairment in spontaneously hypertensive rats: role of central nicotinic receptors. I. *Brain Res* **771**:89-103.

Gil Z, Sack RA, Kedmi M, Harmelin A, Orr-Urtreger A (2002) Increased sensitivity to nicotine-induced seizures in mice heterozygous for the L250T mutation in the alpha7 nicotinic acetylcholine receptor. *Neuroreport* **13**:191-196.

Giubilei F, Strano S, Imbimbo BP, Tisei P, Calcagnini G, Lino S, Frontoni M, Santini M, Fieschi C. (1998) Cardiac autonomic dysfunction in patients with Alzheimer disease: possible pathogenetic mechanisms. *Alzheimer Disease & Associated Disorders* **12**:356-361.

Hellstrom-Lindahl E, Mousavi M, Zhang X, Ravid R, Norberg A (1999) Regional distribution of

JPET #63321

nicotinic receptor subunit mRNAs in human brain: Comparison between Alzheimer and normal brain. *Mol Brain Res* **66**:94-103.

Hiramatsu M, Murasawa H, Nabeshima T, Kameyama T (1998) Effects of U-50,488H on scopolamine-, mecamylamine- and dizocilpine-induced learning and memory impairment in rats. *J Pharmacol Exp Ther* **284**:858-867.

Jonnala RR, Buccafusco JJ (2001) Relationship between the increased cell surface alpha7 nicotinic receptor expression and neuroprotection induced by several nicotinic receptor agonists. *J Neurosci Res* **66**:565-572.

Jonnala RR, Graham JH III, Terry AV Jr, Beach JW, Young JA, Buccafusco JJ (2003) Relative levels of cytoprotection produced by analogs of choline and the role of $\alpha 7$ -nicotinic acetylcholine receptors. *Synapse* **47**:262-269.

Jonnala RR, Terry AV Jr, Buccafusco JJ (2002) Nicotine increases the expression of high affinity nerve growth factor receptors in both in vitro and in vivo. *Life Sci* **15**:543-554.

Kem WR (2000) The brain alpha7 nicotinic receptor may be an important therapeutic target for the treatment of Alzheimer's disease: studies with DMXBA (GTS-21). *Behav Brain Res* **113**:169-181.

Khan IM, Taylor P, Yakash TL (1994) Stimulatory pathways and sites of action of intrathecally

JPET #63321

administered nicotinic agents. *J Pharmacol Exp Ther* **271**:1550-1557.

Levin ED, Bettegowda C, Blosser J, Gordon J (1999) AR-R17779, and alpha7 nicotinic agonist, improves learning and memory in rats. *Behav Pharmacol* **10**:675-680.

Levin ED, Simon BB (1998) Nicotinic acetylcholine involvement in cognitive function in animals. *Psychopharmacology (Berl)* **138**:217-230.

Li XD, Buccafusco JJ (2003) Effect of beta-amyloid peptide 1-42 on the cytoprotective action mediated by $\alpha 7$ nicotinic acetylcholine receptors in growth factor-deprived differentiated PC-12 cells. *J Pharmacol Exp Ther* **307**:670-675.

Li Y, King MA, Meyer EM (2000) Alpha7 nicotinic receptor-mediated protection against ethanol-induced oxidative stress and cytotoxicity in PC12 cells. *Brain Res* **861**:165-167.

Liu QS, Kawai J, Berg DK (2001) β -amyloid peptide blocks the response of $\alpha 7$ -containing nicotinic receptors on hippocampal neurons. *Proc Natl Acad Sci USA* **98**:4734-4739.

Magri' V, Buccafusco JJ (1988) Hypertension following intrathecal injection of cholinergic agonists in conscious rats: Role of endogenous acetylcholine. *J Auton Nerv Sys* **25**:69-77.

Marutle A, Zhang X, Court J, Piggott M, Johnson M, Perry R, Perry E, Nordberg A 2001.

Laminar distribution of nicotinic receptor subtypes in cortical regions in schizophrenia. *J Chem*

JPET #63321

Neuroanat **22**:115-126.

Maurice T, Lockhart BP, Privat A (1996) Amnesia induced in mice by centrally administered beta-amyloid peptides involves cholinergic dysfunction. *Brain Res* **706**:181-193.

Meyer EM, Tay ET, Papke RL, Meyers C, Huang GL, de Fiebre CM (1997) 3-[2,4-Dimethoxybenzylidene]anabaseine (DMXB) selectively activates rat $\alpha 7$ receptors and improves memory-related behaviors in a mecamylamine-sensitive manner. *Brain Res* **768**:49-56.

Nakamura S, Murayama N, Noshita T, Annoura H, Ohno T (2001) Progressive brain dysfunction following intracerebroventricular infusion of beta(1-42)-amyloid peptide. *Brain Res* **912**:128-36.

Olariu A, Yamada K, Mamiya T, Hefco V, Nabeshima T (2002) Memory impairment induced by chronic intracerebroventricular infusion of beta-amyloid (1-40) involves downregulation of protein kinase C. *Brain Res* **957**:278-286.

Papke R, Bencheriff M, Lippiello P (1996) An evaluation of neuronal nicotinic acetylcholine receptor activation by quaternary nitrogen compounds indicates that choline is selective for the $\alpha 7$ subtype. *Neurosci Lett* **213**:201-204.

Pereira EFR, Hilmas C, Santos MD, Alkondon M, Maelicke A, Albuquerque EX (2002) Unconventional ligands and modulators of nicotinic receptors. *J Neurobiol* **53**:479-500.

JPET #63321

Pettit DL, Shao Z, Yakel JL (2001) β -amyloid₁₋₄₂ peptide directly modulates nicotinic receptors in the rat hippocampal slice. *J Neurosci* **21**:RC120 1-5.

Savci V, Cavun S, Goktalay G, Ulus IH (2002) Cardiovascular effects of intracerebroventricularly injected CDP-choline in normotensive and hypotensive animals: the involvement of cholinergic system. *Naunyn Schmiedebergs Arch Pharmacol* **365**:388-398.

Schaeppi U (1967) Effects of nicotine administration to the cat's lower brain stem upon electroencephalogram and autonomic system. *Ann NY Acad Sci* **142**:40-48.

Shimohama S, Kihara T (2001) Nicotinic receptor-mediated protection against beta-amyloid neurotoxicity. *Biol Psychiatry* **49**:233-239.

Si ML, Lee TJ (2002) Alpha7-nicotinic acetylcholine receptors on cerebral perivascular sympathetic nerves mediate choline-induced nitrenergic neurogenic vasodilation. *Circ Res* **91**:62-69.

Stepanichev MY, Moiseeva YV, Lazareva NA, Onufriev MV, Gulyaeva NV (2003) Single intracerebroventricular administration of amyloid-beta (25-35) peptide induces impairment in short-term rather than long-term memory in rats. *Brain Res Bull* **61**:197-205.

Sugaya K, Giacobini E, Chiappinelli VA (1990) Nicotinic Acetylcholine receptor subtypes in human frontal cortex: Changes in Alzheimer's disease. *J Neurosci Res* **27**:349-359.

JPET #63321

Terry AV Jr, Hernandez CM, Buccafusco JJ (2001) Dahl salt-sensitive and salt-resistant rats: examination of learning and memory performance, blood pressure, and the expression of central nicotinic acetylcholine receptors. *Neurosci* **103**:351-363.

Tohyama M, Takatsuji K (1998) Atlas of Neuroactive Substances and Their Receptors in the Rat. *Oxford University Press, NY*.

Trimarchi GR, Glisson WC, Thompson WM, VanLingen J, Buccafusco, JJ (1986) Cholinergic neurons and the cardiovascular response produced by central injection of substance P in the normotensive rat. *Life Sci* **39**:1579-1588.

Yamada K, Tanaka T, Senzaki K, Kameyama T, Nabeshima T (1998) Propentofylline improves learning and memory deficits in rats induced by beta-amyloid protein-(1-40). *Eur J Pharmacol* **349**:15-22.

Yamaguchi Y, Kawashima S (2001) Effects of amyloid-beta-(25-35) on passive avoidance, radial-arm maze learning and choline acetyltransferase activity in the rat. *Eur J Pharmacol* **412**:265-272.

Wang HY, Lee DHS, D'Andrea MR, Peterson PA, Shank RP (1999) β -amyloid₁₋₄₂ binds to α 7 nicotinic acetylcholine receptor with high affinity. *J Biol Chem* **275**:5626-5632.

JPET #63321

Wang HY, Lee DHS, Davis CB, Shank RP (2000) Amyloid peptide A β_{1-42} binds selectively and with picomolar affinity to $\alpha 7$ nicotinic acetylcholine receptors. *J Neurochem* **75**:1155-1161.

JPET #63321

Footnotes

This study was supported by the Office of Research and Development, Medical Research Service, Department of Veterans Affairs.

JPET #63321

Figure Legends

Figure 1.

The time course for the change in mean arterial pressure (MAP) after icv injection of choline (50, 100, or 150 μ g) in freely-moving rats. Each point and vertical bar represents the mean \pm SEM derived from 4-6 experiments.

Figure 2.

The time course for the changes in heart rate after (HR) icv injection of choline (50, 100, or 150 μ g) in freely-moving rats. Each point and vertical bar represents the mean \pm SEM derived from 4-6 experiments.

Figure 3.

Effect of icv pretreatment with 10 μ l of saline (vehicle), or with 10 μ g of MLA, or with 0.3 μ g of atropine on the change in MAP to icv injection of 150 μ g of choline. Saline MLA, or atropine was injected 10 min prior to choline. Compared to saline, MLA significantly inhibited the pressor effect induced by choline [F(1,497)=80.1, P<0.0001]. Atropine pretreatment was without significant effect on the pressor response to choline [F(1,497)=0.72, P=0.40]. **Inset:** Inset: Accompanying HR changes to choline. Both MLA pretreatment [F(1,497)=55.5, P<0.0001] and atropine pretreatment [F(1,495)=9.6, P=0.002] inhibited the changes in HR to choline

Figure 4.

JPET #63321

Effect of icv pretreatment with 10 μ l of saline (control) or 10 μ g of α -bungarotoxin (BTX), or 6 μ g or dihydro- β -erythroidin (DH β E) on the change in MAP to icv injection of 150 μ g of choline. Saline or the antagonists were injected 30 min prior to choline. When compared to the saline pretreatment group, α -bungarotoxin significantly inhibited the increase in MAP produced by 150 μ g of choline [F(1,465)=19.4, P<0.0001]. Under the same conditions, 6 μ g of DH β E failed to significantly alter choline induced pressor response [F(1,350)=0.67, P=0.41]. There also was a significant difference between the BTX and the DH β E treated groups [F(1,350)=51.6, P<0.0001]. Inset: Accompanying HR changes to choline. As with the effects on MAP, significant differences were found between the BTX vs. Saline pretreatment groups [F(1,469)=34.2, P<0.0001] and between the BTX vs. DH β E pretreatment groups [F(1,350)=4.9, P=0.028]. DH β E pretreatment failed to significantly inhibit the HR changes produced by choline [F(1,350)=2.2, =0.14].

Figure 5.

Dose effect of A β 1-42 (1, 10, or 100 pmol, icv) on the choline (150 μ g, icv)-induced changes in MAP. Amyloid peptides were administrated 30 min before choline. A β 1-42 (100 pmol) significantly inhibited the increase in mean arterial pressure (MAP) produced by icv injection of choline [F(1,645)=208.3, P<0.0001]. The 10 pmol dose of A β 1-42 also significantly inhibited the increase in MAP produced by choline relative both to the saline control [F(1,492)=96.0, P<0.0001]; and with respect to the 1 pmole dose [F(1,492)=9.6, P=0.002]. The 1 pmol dose of A β 1-42 was not significantly effective in blocking the choline response (P>0.05). The first time point (0 min) indicates the time of choline administration.

JPET #63321

Figure 6.

Comparison of the abilities of icv pretreatment with A β 1-42 and the reverse sequence peptide A β 40-1 to block the choline induced changes in MAP and heart rate. The peptides were administered 30 min before choline. **Upper panel:** The pressor response to choline (150 μ g, icv) was significantly greater following pretreatment with A β 40-1 as compared with A β 1-42 [F(1,497)=33.3, P<0.0001]; and A β 40-1 pretreatment failed to significantly inhibit the pressor response to choline as compared to the data derived from vehicle pretreated rats [F(1,405)=0.065, P=0.8]. **Lower panel:** The changes in heart rate to choline were significantly greater following pretreatment with A β 1-42 as compared with saline [F(1,487)=46.7, P<0.0001]; and A β 40-1 pretreatment failed to significantly inhibit the changes in heart rate to choline as compared to the data derived from vehicle pretreated rats [F(1,485)=0.58, P=0.4].

JPET #63321

Table 1

Baseline values for Mean Arterial Pressure and Heart Rate for each study protocol.

	MAP (mmHg)	Heart Rate (beats/min)	N
Choline Dose Response (after saline pretreatment)			
50 µg	94±7.5	393±13.8	6
100 µg	108±5.8	384±16.1	6
150 µg	100±5.4	398±15.8	6
Treatments prior to Saline			
Saline	101±3.5	386±14.8	6
Treatments prior to Choline (150µg)			
MLA	99±7.4	359±18.3	6
BTX	105±8.7	392±15.8	6
Atropine	89±6.4	397±7.7	6
Aβ1-42 (100pmol)	96±4.2	385 ± 10.7	11
Aβ1-42 (10pmol)	98±5.4	394±13.6	6
Aβ1-42 (1pmol)	106±4.6	390±12.1	6
Aβ40-1 (100pmol)	90±3.4	385±8.3	6


Measurements were made during the 3-5 min preceding choline administration. MAP (mean

JPET #63321

arterial pressure); MLA (methyllycaconitine); BTX (α -bungarotoxin). Each value represents the mean \pm SEM. * P=0.005 as compared with saline (no injection) mean.


JPET #63321

Figure 1




JPET #63321

Figure 2




JPET #63321

Figure 3




JPET #63321

Figure 4



JPET #63321

Figure 5



JPET #63321

Figure 6

