

JPET #75283

**Sulodexide (KRX-101) Attenuates Myocardial Ischemia/Reperfusion  
Injury and the Deposition of C-Reactive Protein in Areas  
of Infarction without Affecting Hemostasis**

D. Adam Lauver, Erin A. Booth, Andrew J. White,

Enrique Poradosu and Benedict R. Lucchesi

University of Michigan Medical School, Department of Pharmacology, Ann Arbor,

Michigan; DAL, EAB, AJW and BRL

Keryx Biopharmaceuticals, New York, New York; EP

JPET #75283

Running Title:

Sulodexide Attenuates Myocardial Injury

Correspondence:

Dr. Benedict R. Lucchesi

Department of Pharmacology,

University of Michigan Medical School,

1301 MSRB III,

Ann Arbor, Michigan.

E-mail: [benluc@umich.edu](mailto:benluc@umich.edu)

Number of text pages: 24

Number of figures: 6

Number of references: 29

Number of words in the *Abstract*: 240

Number of words in the *Introduction*: 510

Number of words in the *Discussion*: 1,069

Nonstandard abbreviations: MAC, Membrane attack complex; TTC, 2,3,5-triphenyltetrazolium chloride.

Recommended Section Assignment: *Cardiovascular*

## Abstract

Several glycosaminoglycans (GAGs) have been demonstrated to protect the ischemic heart against reperfusion injury, in part, by modulating activation of the complement cascade. The present study assessed the cardioprotective effects of sulodexide (KRX-101), a mixture of GAGs composed of 80% low molecular weight heparin and 20% dermatan sulfate. KRX-101 differs from other GAGs (e.g. heparin) in that it has limited anticoagulant efficacy and can be administered orally. The experimental protocol was designed to determine if KRX-101 could protect the ischemic myocardium. Anesthetized, New Zealand white rabbits underwent 30 minutes of coronary artery occlusion. Intravenous doses of KRX-101 (0.5 mg/kg,  $n = 10$ ) or drug diluent ( $n = 10$ ) were administered at the end of regional ischemia and at each hour of reperfusion. Infarct size, as a percentage of the area at risk, was calculated for both groups. Myocardial infarct size was  $31.3 \pm 4.1\%$  in the vehicle and  $17.3 \pm 3.2\%$  in the KRX-101 treated animals ( $p < 0.05$  vs. vehicle). Activated partial thromboplastin times determined at baseline (pre-ischemia) and at each hour of reperfusion ( $n = 4$ ) were not significantly different between vehicle and KRX-101 treated groups ( $p = ns$ ). Myocardial injury was further assessed by measuring serum levels of cardiac-specific troponin I. KRX-101 administration significantly reduced ( $p < 0.05$ ) the serum concentration of troponin I during reperfusion. The results suggest that KRX-101 may be an effective adjunctive agent in myocardial revascularization procedures, without the risk of increased bleeding.

JPET #75283

The restoration of blood flow to a previously ischemic region is associated with a complex series of events that lead to tissue injury greater than what is attributed to the original ischemic insult, an event referred to as reperfusion injury. The complement system is a component of the innate immune system consisting of a group of proteins found circulating in the blood. The components of the complement system act together to recognize and destroy foreign pathogens. Activation of the complement system, however, can also have adverse actions on host tissues. In the setting of myocardial reperfusion injury, the complement system represents an integral mechanism through which the ischemic tissue undergoes injury leading to cell death and necrosis (Kilgore et al., 1994; Park and Lucchesi, 1999). Previous studies provide evidence for the role of complement by demonstrating the deposition of the terminal complement complex, or the Membrane Attack Complex (MAC), in irreversibly injured myocardial tissue (Schafer et al., 1986). It is hypothesized, therefore, that pharmacological inhibition of complement activation may be beneficial in reducing tissue injury associated with ischemia and reperfusion.

Glycosaminoglycans (GAGs) are a group of compounds reported to be of benefit to the ischemic myocardium by preserving contractile function and reducing tissue injury (Friedrichs et al., 1994; Black et al., 1995; Gralinski et al., 1996; Kilgore et al., 1998; Tanhehco et al., 1999). In addition, selective GAGs are known to possess anti-complement activity in addition to their classical roles as anticoagulants. In the following study, we focus on sulodexide (KRX-101), a purified GAG obtained from porcine mucosa composed of 80% low molecular weight heparin and 20% dermatan sulfate. KRX-101 differs from heparin in that it has a reduced risk of bleeding and is readily

JPET #75283

absorbed upon oral administration. Although GAGs such as heparin are better recognized for their anticoagulant effect, less is known about their anti-inflammatory actions. Identification of more efficacious GAGs, especially those with a limited capacity to alter normal hemostasis, are of potential interest to provide adjunctive therapies to be used in procedures aimed at reducing injury resulting from reperfusion.

An important aspect of complement-mediated myocardial injury that has been the focus of recent research is the role C-reactive protein (CRP) in the activation of the complement system. More sensitive assays for the measurement of CRP have revealed that increased CRP values, even within the range previously considered normal, may be predictive of future coronary events. Once thought of as only a nonspecific indicator of systemic inflammation, recent epidemiological research indicates that CRP might be directly involved in the pathogenesis of ischemic diseases through the activation the complement system (Beranek, 1997; Du Clos, 2000; Agrawal et al., 2001). Previous studies have demonstrated that plasma CRP concentration is directly related to infarct size following ischemia/reperfusion (Barrett et al., 2002; Hirschfield and Pepys, 2003).

In the current study, the cardioprotective effects of KRX-101 were evaluated in male New Zealand white rabbits subjected to regional myocardial ischemia/reperfusion. The protocol was designed to test the ability of KRX-101 to reduce the extent of myocardial injury associated with regional ischemia and reperfusion.

## Materials and Methods

**Guidelines for Animal Research.** The procedures used in this study were in agreement with the guidelines of the University of Michigan Committee on the Use and Care of Animals. The University of Michigan Unit for Laboratory Animal Medicine provides veterinary care. The University of Michigan is accredited by the American Association of Accreditation of Laboratory Animal Health Care, and the animal care use program conforms to the standards in *The Guide for the Care and Use of Laboratory Animals*, publication number NIH 86-23.

**Surgical Preparation.** Male New Zealand White rabbits (2.2-2.6 kg) were anesthetized with a combination of xylazine (3.0 mg/kg) and ketamine (35 mg/kg) administered intramuscularly, followed by an intravenous injection of sodium pentobarbital (15 mg/kg). After insertion of an endotracheal tube, the animals were placed on positive pressure ventilation with room air. The right jugular vein was isolated and cannulated for drug administration. The right carotid artery was isolated and instrumented with a Millar™ catheter micro-tip pressure transducer (Millar Instruments Inc., Houston, TX) positioned immediately above the aortic valve to monitor aortic blood pressure. The lead II electrocardiogram was monitored throughout the experiment. A left thoracotomy and pericardiotomy were performed, followed by identification of the left anterior descending coronary artery. A silk suture (3-0; Genzyme Corporation, Fall River, MA) was passed under the artery and around a short length of polyethylene tubing. Simultaneous downward displacement of the polyethylene tubing while applying upward traction on the suture resulted in occlusion of the coronary artery and cessation of regional myocardial blood flow. Coronary artery occlusion was maintained for 30 min after

JPET #75283

which time reperfusion was initiated by withdrawing the polyethylene tubing. Regional myocardial ischemia was verified by the presence of a zone of cyanosis in the area of distribution of the occluded vessel and by changes in the electrocardiogram consistent with the presence of transmural regional myocardial ischemia (ST-segment elevation).

**Experimental Protocol.** The animals were allowed to stabilize for 15 min before beginning the protocol that involved two experimental treatment groups. The respective groups received intravenous doses of KRX-101 (0.5 mg/kg, n = 7) or saline (drug diluent, n = 7) administered at the end of regional ischemia and at each hour after the start of reperfusion, excluding the end of the fourth and final hour.

**Determination of Infarct Size.** At the completion of the 4 hr reperfusion period, the hearts were removed, the aorta was cannulated, and the coronary vascular bed was perfused on a Langendorff apparatus with Krebs-Henseleit buffer at a constant flow of 30 to 32 ml/min. The hearts were perfused with buffer for 10 min to clear the vascular compartment of plasma and blood cellular elements. Fifty milliliters of a 1% solution of triphenyltetrazolium chloride (TTC) in phosphate buffer (pH 7.4, 37°C) was perfused through the heart. TTC demarcates the noninfarcted myocardium within the area at risk with a brick red color, indicating the presence of a formazan precipitate resulting from reduction of TTC by dehydrogenases present in viable myocardial tissue. Irreversibly injured tissue, lacking cytosolic dehydrogenases, is unable to form the formazan precipitate and appears pale yellow. Upon completion of the TTC infusion, the left anterior descending coronary artery was ligated at the site identical to that ligated during the induction of regional myocardial ischemia. The perfusion pump was stopped, and 3 ml of a 0.25% solution of Evans Blue was injected slowly through a side-arm port

## JPET #75283

connected to the aortic cannula. The dye was passed through the heart for 10 seconds to ensure its uniform tissue distribution. The presence of Evans Blue was used to demarcate the left ventricular tissue that was not subjected to regional ischemia, as opposed to the risk region. The heart was removed from the perfusion apparatus and cut into transverse sections at right angles to the vertical axis. The right ventricle, apex, and atrial tissue were discarded. Both surfaces of each tissue section were traced onto clear acetate sheets. The images were then digitized using a flatbed scanner. The areas of the normal left ventricle non-risk region, area at risk, and infarct region were determined by calculating the number of pixels occupying each area using Adobe PhotoShop software (Adobe Systems, Seattle, WA). Total area at risk is expressed as the percentage of the left ventricle. Infarct size is expressed as the percentage of the area at risk.

**Hematological Measurements.** Determinations of the whole blood aPTT and *ex vivo* platelet aggregation were made at baseline and repeated after the end of regional ischemia and at each hour of reperfusion. Blood (5 ml) was withdrawn from the right jugular vein cannula into a plastic syringe containing 3.7% sodium citrate as the anticoagulant [1:10 citrate to blood (v/v)] for aPTT and *ex vivo* platelet aggregation determinations. Two milliliters of blood were used in the determination of activated partial thromboplastin time (aPTT) using a Hemochron analyzer (Technidyne, Edison, NJ) with the reagents supplied by the manufacturer. Platelet-rich plasma (PRP), the supernatant present after centrifugation of the remaining 3 ml of whole blood at 660 rpm for 10 min, was diluted with phosphate buffered saline to achieve a platelet count of 200,000 cells/ml. Platelet-poor plasma (PPP) was prepared by centrifuging the remaining blood at 12,000 g for 10 min and discarding the bottom cellular layer. *Ex vivo* platelet



## JPET #75283

aggregation was assessed by established spectrophotometric methods with the use of a four-channel aggregometer (BioData PAP-4; BioData Corp., Horsham, PA) by recording the increase in light transmission through a stirred suspension of PRP maintained at 37°C. Aggregation was induced with arachidonic acid (AA, 0.65 mM) or  $\gamma$ -thrombin (25 nM). A subaggregatory concentration of epinephrine (550 nM) was used to prime the platelets before addition of the agonists to induce platelet aggregation. Values for platelet aggregation are expressed as percentage of light transmission standardized to PRP and PPP samples yielding 0 and 100% light transmission, respectively.

**Biochemical Markers of Myocardial Damage.** Plasma concentrations of cardiac-specific troponin I (cTnI) were determined by enzyme-linked immunosorbent assays (Developed in conjunction with Dr. Chris Chadwick, Life Diagnostic, Inc.). Briefly, serum was prepared from whole blood drawn at baseline, after ischemia and each hour of reperfusion. Samples were frozen immediately in liquid nitrogen and store at -80°C. On the day of the assay, samples were thawed over ice and diluted appropriately with sample diluent supplied with each assay kit. Protein concentrations were determined using the OD of each sample as compared to a standard curve.

**Immunofluorescent detection of the membrane attack complex and C-reactive protein.** Tissue samples used for infarct size determination were fixed in 10% buffered formalin immediately after the completion of the experimental protocol. The tissue samples were embedded in paraffin blocks and cut into sections of 2  $\mu$ m in thickness, which were mounted on glass slides. Two consecutive sections (mirror images) from a single heart slice were mounted on each slide. The slides were deparaffinized in three washes of xylene and rehydrated in an ethanol gradient. To remove any residual formalin

JPET #75283

and paraffin, antigen unmasking was performed using a commercially available antigen unmasking solution (Vector Laboratories, Burlingame, CA) and a pressure cooker (Fagor America, Inc., Lyndhurst, NJ). Briefly, slides were placed in a boiling solution of the unmasking agent and water. The pressure cooker was sealed and slides were incubated for 1 minute once the cooker reached maximal pressure (approximately 15 pounds per square inch). The slides were cooled immediately in a tap water bath and blocked with 3% immunohistochemical grade bovine serum albumin (BSA) for 30 minutes. Primary antibodies were incubated at room temperature in a humidity chamber for 45 minutes. One section per slide was incubated with a chicken anti-rabbit CRP antibody (5 µg/ml final concentration, Immunology Consultants Laboratory, Inc.). The other section was incubated with a chicken anti-rabbit MAC antibody (1:2500 final dilution, developed in conjunction with Lampire Biological Laboratories, Pipersville, PA). After three BSA washes, each section was incubated with a biotinylated secondary antibody (Goat anti-chicken, 1.5 µg/ml final, Vector Laboratories) for 30 minutes. Following 3 additional BSA washes, the slides were incubated with Fluorescein and Texas Red (CRP and MAC sections, respectively) labeled streptavidin (Fluorescent Streptavidin Kit, Vector Laboratories) to visualize the proteins. The fluorescent streptavidin reagents were allowed to incubate for 10 minutes. ProLong Gold antifade (Molecular Probes, Eugene, Oregon) and cover slips were used to preserve the sections. For comparison, digital images were captured using a digital camera (Sony DKC5000, Sony Corporation of America, New York) connected to a Leica fluorescent stereoscope (Leica MZ FLIII) and the accompanying software (Leica Microsystems, Germany). Images were analyzed using IP Lab (Scanalytics, Inc., Fairfax, VA) software to determine mean fluorescence

JPET #75283

intensity per heart section. The sections were normalized to the amount of background on each slide. The mean intensities for three hearts in each treatment group were averaged and compared.

**Statistical Analysis.** Results are expressed as mean values  $\pm$  S.E.M. Parameters between the two groups were compared using the Student's *t* test for unpaired comparisons. *P* values of  $< 0.05$  and  $< 0.01$  are regarded as significant and denoted by an asterisk and double asterisk, respectively.

## Results

**Cardiac contractile parameters.** Hemodynamic variables were obtained to determine the effects of KRX-101 in mediating alterations in arterial blood pressure and heart rate. The rate-pressure product (RPP), defined as mean arterial blood pressure multiplied by the heart rate divided by 100, was used as an indicator of myocardial oxygen consumption. As depicted in Figure 1, the RPP decreased slightly in each group (KRX-101 and vehicle) after equilibration and then remained stable throughout the duration of the protocol. The Lead II electrocardiogram did not detect any changes upon administration of KRX-101 or vehicle. Sustained ST-segment elevation was observed in all rabbits during regional myocardial ischemia, however these changes returned toward baseline upon reperfusion. No deaths, from either cardiac arrhythmias or pump failure were noted in any of the groups.

**Effect of KRX-101 on myocardial infarct size.** Each treatment group consisted of ten animals in which either KRX-101 or the control vehicle was administered at the end of regional ischemia and at each hour of reperfusion, excluding the end of the fourth and final hour. The size of the area at risk expressed as a percentage of the total left ventricle was similar in each of the groups. Rabbits treated with KRX-101 exhibited significantly smaller infarcts expressed as a percentage of the area at risk compared with rabbits treated with vehicle (Figure 2).

**Blood Coagulation Parameters.** The aPTT determinations were similar in both vehicle and KRX-101 treatment groups at baseline, post ischemia and each hour of reperfusion (Figure 3). Percentage of platelet aggregation response did not change through out the course of the experiment in the control group. Whereas platelet aggregation in responses

JPET #75283

to AA remained unchanged, aggregation induced by  $\gamma$ -thrombin was reduced significantly in PRP prepared from blood obtained from rabbits treated with KRX-101 (Figure 4).

**Plasma concentrations of biochemical markers of cardiac damage.** Serum concentrations of cardiac-specific troponin I (cTnI) were similar at baseline and immediately after reperfusion. KRX-101 treated rabbits exhibited significantly lower values cTnI at 1, 2, 3 and 4 hours after the onset of reperfusion as compared to vehicle controls (Figure 5).

**Immunofluorescence.** Left ventricular tissue sections used for immunofluorescence were taken from hearts that had been subjected to 30 minutes of regional ischemia, followed by 4 hours of reperfusion. Hearts from animals treated with the drug vehicle demonstrated bright fluorescence with both anti-CRP and anti-MAC antibodies, indicating the deposition of both proteins in the area of infarction. Conversely, hearts treated with KRX-101 exhibited reduced fluorescence and therefore a reduction in the deposition of CRP and MAC (Figure 6a-d). The mean intensity of fluorescence for hearts treated with KRX-101 was significantly ( $p < 0.05$ ) lower than vehicle for both CRP and MAC stained sections (Figure 6e).

## Discussion

It is understood that heparin and other GAGs have therapeutic uses beyond their traditional role as anticoagulants (Rajtar et al., 1993). In this study we demonstrated that the intravenous administration of KRX-101 results in a decrease in myocardial infarct size in a model of *in vivo* regional ischemia/reperfusion. The results are in accordance with previous studies in which it was found that other glycosaminoglycans have the ability to reduce infarct size *in vivo* (Friedrichs et al., 1994; Black et al., 1995; Gralinski et al., 1996; Kilgore et al., 1998; Tanhehco et al., 1999).

The chemical composition of KRX-101 is defined as 80% low molecular weight (7000 D) heparin fraction and 20% dermatan sulfate. Low molecular weight heparin contains the same dimeric components as unfractionated heparin, but has a lower degree of sulfation and shorter polysaccharide chain length. Dermatan sulfate is a polysaccharide made up of many various disaccharide units with a mean molecular weight of 25,000 D. As a result of the presence of both fractions, KRX-101 potentiates the antiprotease activities of both antithrombin III and heparin cofactor II simultaneously. Although structurally similar, KRX-101 has major differences from unfractionated heparin including prolonged half-life, reduced effect on global coagulation and oral bioavailability (Callas et al., 1993; Buchanan et al., 1994). Low molecular weight heparin has previously been showed to reduce infarct size following ischemia and reperfusion (Gralinski et al., 1997; Libersan et al., 1998), however dermatan sulfate has yet to be individually investigated.

Due to their ability to inhibit the complement cascade, it is hypothesized that GAGs may prevent the adverse events associated with complement activation associated

JPET #75283

with ischemia/reperfusion. Activation of the complement cascade leads to the assembly of the MAC on cell membranes. Deposition of a sufficient number of MAC molecules on a target cell results in the disruption of the cell membrane and ultimately cell lysis.

Intravenous administration of 0.5 mg/kg KRX-101 immediately after an ischemic period and at hourly intervals during reperfusion was associated with a significant decrease in myocardial infarct size expressed as a percentage of the area at risk when compared with vehicle treated control rabbits. The AAR in hearts from both treated and control groups was not statistically different, thereby demonstrating that a similar amount of myocardial tissue was subjected to regional ischemia, allowing a valid comparison of infarct size between the two groups when expressed as a percentage of the risk region. Hemodynamic parameters including the rate-pressure product were similar between groups, suggesting that KRX-101 is not acting to protect the myocardium by decreasing the myocardial demand for oxygen. These data indicate that differences in the myocardial oxygen demand did not contribute to protecting the myocardium after ischemia/reperfusion.

Analysis of platelet reactivity, as determined by *ex vivo* platelet aggregation, demonstrated a decrease in  $\gamma$ -thrombin-induced platelet aggregation in KRX-101 treated animals as compared to vehicle treated animals. These data agree with previously reported findings (Rajtar et al., 1993; Cerletti et al., 1994) which indicated that KRX-101 inhibits thrombin-induced platelet activation. Interestingly, evaluation of the effects of KRX-101 on coagulation, using the activated partial thromboplastin time (aPTT) demonstrated that at the dose used in our study, there was little or no change in hemostasis.

JPET #75283

As another method of quantifying cardiac injury after ischemia/reperfusion, we measured the serum concentration of a biochemical marker of tissue injury. Cardiac-specific troponin I (cTnI) is a component of the contractile machinery within myocytes. Upon cell lysis, the protein is released into the blood and can be measured using a specific immunoassay. As would be predicted based on infarct size data, it was found that KRX-101 significantly reduced the concentration of this marker during reperfusion.

Along with other GAGs, the precise mechanism by which KRX-101 achieves myocardial protection after ischemia/reperfusion has yet to be determined. Since KRX-101 affects several aspects of reperfusion injury, various hypotheses may be drawn concerning its role in cardioprotection. Previous studies have indicated that GAGs are effective inhibitors of the complement system. We therefore sought to investigate the likelihood that KRX-101 acts to protect the myocardium through inhibition of the complement cascade.

C-reactive protein (CRP) is an acute phase protein that has been demonstrated to be a highly sensitive, but nonspecific, marker of inflammation. Not only are plasma levels of CRP elevated in inflammatory diseases, increased CRP concentrations are associated with increased mortality due to cardiovascular events (de Beer et al., 1982; Yeh et al., 2001). Thus, CRP may be an indicator of myocardial injury, as well as being involved in the pathogenesis of irreversible myocardial injury (de Beer et al., 1982; Ridker et al., 1998; Westhuyzen and Healy, 2000; Yeh et al., 2001). The proposed mechanism of CRP involvement is through local activation of the complement system (Volanakis, 1982). Systemic administration of CRP was found to increase the extent myocardial necrosis, through complement-dependent mechanisms, in experimental acute



JPET #75283

myocardial infarction (Griselli et al., 1999). The endogenous production of CRP in response to a remote inflammatory response likewise results in more extensive myocardial injury after ischemia/reperfusion (Barrett et al., 2002). CRP has been shown to activate the classical complement pathway providing a possible mechanism linking CRP to mortality due to myocardial infarction (Kaplan and Volanakis, 1974; Wolbink et al., 1996; Nijmeijer et al., 2003). Using an immunofluorescent method to determine the presence of tissue bound CRP and MAC; we were able to show that KRX-101 significantly reduced the deposition of both CRP and MAC, which were found localized within the area of infarction.

The results of this study demonstrate that KRX-101 effectively reduces infarct size after reperfusion of the ischemic myocardium. Although the exact mechanism by which KRX-101 protects the myocardium has yet to be determined, we provide evidence that KRX-101 acts to inhibit complement activation, possibly through the inhibition of CRP, resulting in attenuation of myocardial ischemia/reperfusion injury. The presence of the MAC in infarcted myocardium, together with the ability of inhibitors of complement to protect the ischemic myocardium, underscore the importance of pharmacologic agents capable of inhibiting the complement system in the modulation of reperfusion injury. This study also provides evidence that there is a clear separation between the anticoagulant and anti-complement effects of glycosaminoglycans. We show that a dose of KRX-101 that was previously found to have little or no effect on coagulation (Harenberg, 1998) retains the ability to protect the ischemic myocardium. Therefore, KRX-101 may be a useful agent in the reduction of ischemia/reperfusion injury with a reduced risk of adverse effects on hemostasis.

## References

- Agrawal A, Shrive AK, Greenhough TJ and Volanakis JE (2001) Topology and structure of the C1q-binding site on C-reactive protein. *J Immunol* **166**:3998-4004.
- Barrett TD, Hennan JK, Marks RM and Lucchesi BR (2002) C-reactive-protein-associated increase in myocardial infarct size after ischemia/reperfusion. *J Pharmacol Exp Ther* **303**:1007-1013.
- Beranek JT (1997) C-reactive protein and complement in myocardial infarction and postinfarction heart failure. *Eur Heart J* **18**:1834-1836.
- Black SC, Gralinski MR, Friedrichs GS, Kilgore KS, Driscoll EM and Lucchesi BR (1995) Cardioprotective effects of heparin or N-acetylheparin in an in vivo model of myocardial ischaemic and reperfusion injury. *Cardiovasc Res* **29**:629-636.
- Buchanan MR, Liao P, Smith LJ and Oforu FA (1994) Prevention of thrombus formation and growth by antithrombin III and heparin cofactor II-dependent thrombin inhibitors: importance of heparin cofactor II. *Thromb Res* **74**:463-475.
- Callas DD, Hoppensteadt DA, Jeske W, Iqbal O, Bacher P, Ahsan A and Fareed J (1993) Comparative pharmacologic profile of a glycosaminoglycan mixture, Sulodexide, and a chemically modified heparin derivative, Suleparotide. *Semin Thromb Hemost* **19 Suppl 1**:49-57.
- Cerletti C, Rajtar G, Marchi E and de Gaetano G (1994) Interaction between glycosaminoglycans, platelets, and leukocytes. *Semin Thromb Hemost* **20**:245-253.

- de Beer FC, Hind CR, Fox KM, Allan RM, Maseri A and Pepys MB (1982)  
Measurement of serum C-reactive protein concentration in myocardial ischaemia  
and infarction. *Br Heart J* **47**:239-243.
- Du Clos TW (2000) Function of C-reactive protein. *Ann Med* **32**:274-278.
- Friedrichs GS, Kilgore KS, Manley PJ, Gralinski MR and Lucchesi BR (1994) Effects of  
heparin and N-acetyl heparin on ischemia/reperfusion-induced alterations in  
myocardial function in the rabbit isolated heart. *Circ Res* **75**:701-710.
- Gralinski MR, Driscoll EM, Friedrichs GS, DeNardis MR and Lucchesi BR (1996)  
Reduction of Myocardial Necrosis After Glycosaminoglycan Administration:  
Effects of a Single Intravenous Administration of Heparin or N-Acetylheparin 2  
Hours Before Regional Ischemia and Reperfusion. *J Cardiovasc Pharmacol Ther*  
**1**:219-228.
- Gralinski MR, Park JL, Ozeck MA, Wiater BC and Lucchesi BR (1997) LU 51198, a  
highly sulfated, low-molecular-weight heparin derivative, prevents complement-  
mediated myocardial injury in the perfused rabbit heart. *J Pharmacol Exp Ther*  
**282**:554-560.
- Griselli M, Herbert J, Hutchinson WL, Taylor KM, Sohail M, Krausz T and Pepys MB  
(1999) C-reactive protein and complement are important mediators of tissue  
damage in acute myocardial infarction. *J Exp Med* **190**:1733-1740.
- Harenberg J (1998) Review of pharmacodynamics, pharmacokinetics, and therapeutic  
properties of sulodexide. *Med Res Rev* **18**:1-20.
- Hirschfield GM and Pepys MB (2003) C-reactive protein and cardiovascular disease:  
new insights from an old molecule. *Qjm* **96**:793-807.

- Kaplan MH and Volanakis JE (1974) Interaction of C-reactive protein complexes with the complement system. I. Consumption of human complement associated with the reaction of C-reactive protein with pneumococcal C-polysaccharide and with the choline phosphatides, lecithin and sphingomyelin. *J Immunol* **112**:2135-2147.
- Kilgore KS, Friedrichs GS, Homeister JW and Lucchesi BR (1994) The complement system in myocardial ischaemia/reperfusion injury. *Cardiovasc Res* **28**:437-444.
- Kilgore KS, Naylor KB, Tanhehco EJ, Park JL, Booth EA, Washington RA and Lucchesi BR (1998) The semisynthetic polysaccharide pentosan polysulfate prevents complement-mediated myocardial injury in the rabbit perfused heart. *J Pharmacol Exp Ther* **285**:987-994.
- Libersan D, Khalil A, Dagenais P, Quan E, Delorme F, Uzan A and Latour JG (1998) The low molecular weight heparin, enoxaparin, limits infarct size at reperfusion in the dog. *Cardiovasc Res* **37**:656-666.
- Nijmeijer R, Lagrand WK, Lubbers YT, Visser CA, Meijer CJ, Niessen HW and Hack CE (2003) C-reactive protein activates complement in infarcted human myocardium. *Am J Pathol* **163**:269-275.
- Park JL and Lucchesi BR (1999) Mechanisms of myocardial reperfusion injury. *Ann Thorac Surg* **68**:1905-1912.
- Rajtar G, Marchi E, de Gaetano G and Cerletti C (1993) Effects of glycosaminoglycans on platelet and leucocyte function: role of N-sulfation. *Biochem Pharmacol* **46**:958-960.

JPET #75283

- Ridker PM, Buring JE, Shih J, Matias M and Hennekens CH (1998) Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. *Circulation* **98**:731-733.
- Schafer H, Mathey D, Hugo F and Bhakdi S (1986) Deposition of the terminal C5b-9 complement complex in infarcted areas of human myocardium. *J Immunol* **137**:1945-1949.
- Tanhehco EJ, Kilgore KS, Naylor KB, Park JL, Booth EA and Lucchesi BR (1999) Reduction of myocardial infarct size after ischemia and reperfusion by the glycosaminoglycan pentosan polysulfate. *J Cardiovasc Pharmacol* **34**:153-161.
- Volanakis JE (1982) Complement activation by C-reactive protein complexes. *Ann N Y Acad Sci* **389**:235-250.
- Westhuyzen J and Healy H (2000) Review: Biology and relevance of C-reactive protein in cardiovascular and renal disease. *Ann Clin Lab Sci* **30**:133-143.
- Wolbink GJ, Brouwer MC, Buysmann S, ten Berge IJ and Hack CE (1996) CRP-mediated activation of complement in vivo: assessment by measuring circulating complement-C-reactive protein complexes. *J Immunol* **157**:473-479.
- Yeh ET, Anderson HV, Pasceri V and Willerson JT (2001) C-reactive protein: linking inflammation to cardiovascular complications. *Circulation* **104**:974-975.

JPET #75283

**Footnote:**

Supported by the Cardiovascular Research Fund, University of Michigan Medical School.

JPET #75283

Figure 1. Rate-pressure product for KRX-101 (closed squares) and vehicle (open circles) treated animals. Rate-pressure product was calculated with the formula beats per minute x millimeters of Hg/100. No significant differences were noted between groups at any time point or within the same group throughout the reperfusion period. Arrow indicates the onset of reperfusion.

Figure 2. Effects of KRX-101 on myocardial infarct size after 30 min of left anterior descending coronary artery occlusion and 4 hours of reperfusion compared with vehicle. KRX-101 was administered at doses of 0.5 mg/kg at the onset and each hour of reperfusion excluding the fourth and final hour. The areas at risk were similar between groups, which indicates that the degree of the insult was similar. Infarct size after ischemia/ reperfusion is expressed as a percentage of the area at risk. The infarct region is decreased significantly in the group treated with KRX-101 compared with vehicle. Values are presented as mean  $\pm$  SEM; vehicle group, n = 10 (white bars); KRX-101 group, n = 10 (black bars); \*  $p < 0.05$  versus baseline.

Figure 3. The effect of KRX-101 administration on whole blood activated partial thromboplastin time (aPTT). No significant differences were noted between groups at any time point. Values are presented as mean  $\pm$  SEM; vehicle group, n = 4 (white bars); KRX-101 group, n = 4 (black bars).

Figure 4. Effect of KRX-101 administration on serum levels of a biochemical marker of cardiac damage. Administration of KRX-101 significantly reduced serum concentrations

JPET #75283


of cardiac specific troponin I (cTnI) at one, two, three and four hours after reperfusion compared to vehicle control. Values are presented as mean  $\pm$  SEM; vehicle group, n = 6 (white bars); KRX-101 group, n = 6 (black bars); \* p < 0.05 versus vehicle control.

Figure 5. The effect of KRX-101 administration on  $\gamma$ -thrombin-induced platelet reactivity (A), and arachidonic acid induced platelet aggregation (B). Administration of KRX-101 significantly reduced  $\gamma$ -thrombin induced platelet aggregation (white bars, n = 4) beginning at one hour and continuing throughout the experimental protocol compared to vehicle administration (black bars, n=4). Arachidonic acid induced platelet aggregation (B) did not differ between groups. Values are presented as mean  $\pm$  SEM; \*\* p < 0.01 versus vehicle control.


Figure 6. Representative fluorescent images of a heart from a control animal (A and C) and an animal treated with KRX-101 (B and D) after 30 min of ischemia and 4 hr of reperfusion. In the vehicle treated animal, staining for CRP (A) and MAC (C) are present in areas of infarction. In KRX-101 treated hearts, little or no staining for CRP (B) or MAC (D) can be observed in areas of infarction. E, Graph illustrating the comparison of mean fluorescence intensity per heart section. Values are presented as mean  $\pm$  SEM; vehicle group, n = 3 (white bars); KRX-101 group, n = 3 (black bars); \* p < 0.05 versus vehicle control.



# Figure 1



## Figure 2



### Figure 3






Figure 4



## Figure 5

A.



B.




Figure 6

