

Pharmacological and Electrophysiological Characterization of Store-operated Currents and Capacitative Ca^{2+} Entry in Vascular Smooth Muscle Cells

Liubov I. Brueggemann, Daniel R. Markun, Kyle K. Henderson,

Leanne L. Cribbs, and Kenneth L. Byron

Loyola University Chicago

Departments of Pharmacology (LIB, DRK, KKH, KLB) and Medicine (LLC)

Maywood, IL 60153

Running title: I_{SOC} & CCE in vascular smooth muscle cells.

Address for correspondence:

Kenneth L. Byron, Ph.D.

Loyola University Medical Center

Dept. of Pharmacology & Experimental Therapeutics

2160 South First Avenue

Maywood, IL 60153

Tel.: 708-327-2819. Fax.: 708-216-6596. E-mail: kbyron@lumc.edu

The number of text pages: 41 pages

Number of tables: 0

Number of figures: 9

Number of references: 49

Number of words in the Abstract: 244

Number of words in the Introduction: 637

Number of words in the Discussion: 2134

Nonstandard abbreviations used in the paper:

$[Ca^{2+}]_i$, cytosolic free Ca^{2+} concentration

2-APB, 2-aminoethoxydiphenylborane

AS-TRPC1-GFP, adenoviral vector for expression of antisense TRPC1 DNA and co-expression of
Green Fluorescent Protein

BAPTA, 1,2-Bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid

BSA, bovine serum albumin

CCE, capacitative Ca^{2+} entry

CPA, cyclopiazonic acid

InsP_3 , inositol 1,4,5-trisphosphate

I_{SOC} , store-operated current

LOE 908, (R,S)-(3,4-dihydro 6,7-dimethoxy-isoquinoline-1-yl)-2-phenyl- N,N-di-[2-(2,3,4-trimethoxyphenyl)ethyl]-acetamide

MASMC, mesenteric artery smooth muscle cells

MOI, multiplicity of infection

NCCE, non-capacitative Ca^{2+} entry

OAG, 1-oleoyl-2-acetyl-sn-glycerol

PCR, polymerase chain reaction

PKC, protein kinase C

S.E., standard error of the mean

siRNA, small interfering RNA

SOCE, store-operated Ca^{2+} entry

SR, sarcoplasmic reticulum

TRP, Transient Receptor Potential non-selective cation channel

VSM vascular smooth muscle

Recommended section assignment: Cardiovascular

ABSTRACT

Capacitative Ca^{2+} entry (CCE) in vascular smooth muscle cells contributes to vasoconstrictor and mitogenic effects of vasoactive hormones. In A7r5 rat aortic smooth muscle cells, measurements of cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) have demonstrated that depletion of intracellular Ca^{2+} stores activates CCE. However, there is disagreement in published studies regarding the regulation of this mechanism by the vasoconstrictor hormone, $[\text{Arg}^8]$ -vasopressin (AVP). We have employed electrophysiological methods to characterize the membrane currents activated by store depletion (I_{SOC} , store-operated current). Because of different recording conditions, it has not been previously determined whether I_{SOC} corresponds to CCE measured using fura-2. Nor has the channel protein responsible for CCE been identified. In the present study, the pharmacological characteristics of I_{SOC} , including its sensitivity to blockade by 2-aminoethoxydiphenylborane, diethylstilbestrol, or micromolar Gd^{3+} , were found to parallel the effects of these drugs on thapsigargin- or AVP-activated CCE measured under identical external ionic conditions using fura-2. Thapsigargin-stimulated I_{SOC} was also measured in freshly isolated rat mesenteric artery smooth muscle cells (MASMC). Members of the TRP family of non-selective cation channels, TRPC1, TRPC4, and TRPC6 were detected by RT-PCR and Western blot in both A7r5 cells and MASMC. TRPC1 expression was reduced in a stable A7r5 cell line expressing a small interfering RNA (siRNA) or by infection of A7r5 cells with an adenovirus expressing a TRPC1 antisense nucleotide sequence. Thapsigargin-stimulated I_{SOC} was reduced in both the TRPC1 siRNA- and TRPC1 antisense-expressing cells, suggesting that the TRPC1 channel contributes to the I_{SOC} /CCE pathway.

INTRODUCTION

Vasoconstrictor hormones cause contraction of vascular smooth muscle (VSM) cells by increasing cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$), which in turn activates the cells' contractile apparatus. A well-characterized signal transduction pathway leading to elevation of $[\text{Ca}^{2+}]_i$ involves inositol trisphosphate- (InsP_3)- mediated release of Ca^{2+} from the sarcoplasmic reticulum (SR) and Ca^{2+} entry via non-selective calcium-permeable cation channels. This pathway may also contribute to the mitogenic actions of vasoactive hormones (Wang *et al.*, 2001).

The A7r5 rat aortic smooth muscle cell line is among the most extensively characterized vascular smooth muscle preparations in terms of agonist-stimulated Ca^{2+} signaling pathways. Using the Ca^{2+} -sensitive fluorescent indicator, fura-2, the vasoconstrictor hormone $[\text{Arg}^8]$ -vasopressin (AVP) was found to activate both capacitative (also known as store-operated) and non-capacitative Ca^{2+} entry (CCE and NCCE, respectively) in A7r5 cells (Byron and Taylor, 1995; Broad et al., 1999). Several subsequent studies (Moneer and Taylor, 2002; Moneer et al., 2003; Moneer et al., 2005) have utilized fura-2 fluorescence measurements to examine the regulation of these Ca^{2+} entry pathways by AVP, with some disagreement among the published results (Brueggemann et al., 2005). It remains unclear whether CCE contributes to a sustained elevation of $[\text{Ca}^{2+}]_i$ while AVP is present.

CCE and NCCE in A7r5 cells probably involve different Ca^{2+} -permeable channels because they appear to have different permeabilities to Sr^{2+} and Mn^{2+} (Byron and Taylor, 1995) and are differentially sensitive to pharmacological blockers, such as 2-aminoethoxydiphenylborane (2-APB), LOE 908 and Gd^{3+} (Broad et al., 1999; Iwamuro et al., 1999; Moneer and Taylor, 2002). However, the earlier fura-2 fluorescence studies did not characterize the electrophysiological properties of the CCE pathway, and the channel proteins that are the presumed targets of the pharmacological blockers remain unidentified.

A number of earlier electrophysiological studies (Nakajima et al., 1996; Iwasawa et al., 1997; Iwamuro et al., 1999; Jung et al., 2002) failed to detect store-operated currents in A7r5 cells despite evidence from fura-2 fluorescence studies that Ca^{2+} influx is stimulated following depletion of Ca^{2+} stores. In a recent study (Brueggemann *et al.*, 2005), we reported conditions under which store-operated currents can be detected in A7r5 cells, but the resulting current differs in its apparent cation selectivity and voltage-dependence from I_{SOC} recorded from other vascular smooth muscle preparations. Conditions used for recording isolated currents are generally much different than the ionic conditions used to detect Ca^{2+} entry using fura-2. These conditions may affect the behavior of the channels and their responses to pharmacological agents. Thus there is a paucity of information related to the electrophysiological or molecular nature of the CCE pathway in vascular smooth muscle cells and disagreements among different groups as to the mechanisms regulating its activity.

Molecular candidates for the channels involved in capacitative Ca^{2+} entry pathways may be emerging from the discovery in the mid-1990s of the *Drosophila* Transient Receptor Potential (TRP) channels and 7 canonical mammalian TRP homologues (TRPC1-TRPC7; reviewed by Montell (2005)). The primary structure of these proteins indicates that they have six membrane-spanning domains reminiscent of the voltage-gated K^{+} channel family. In analogy to K^{+} channels, the TRP channels are believed to form as tetramers of individual TRP homologue protein subunits. The TRPC proteins are ubiquitously expressed in mammalian tissues, with many cell types, including VSM, expressing multiple TRPC homologues. When expressed heterologously they generally form functional channels that exhibit electrophysiological properties of non-selective cation channels.

In the present study we use electrophysiological techniques to record store-operated currents (I_{SOC}) in A7r5 cells and mesenteric artery smooth muscle cells (MASMC). The pharmacological characteristics of I_{SOC} are compared with CCE recorded with fura-2 under similar ionic conditions. The findings suggest that CCE is active during the sustained phase of AVP Ca^{2+} signaling. We have also

identified several TRPC homologues expressed in A7r5 cells and MASMC. Evidence is presented to support a role of TRPC1 as a component of the store-operated Ca^{2+} entry pathway.

METHODS

A7r5 cell culture

A7r5 cells were cultured as described previously (Byron and Taylor, 1993). Cells were subcultured onto rectangular (9x22-mm, no.1½) glass coverslips or plastic tissue culture dishes (Corning, Acton, MA, USA). For fura-2 fluorescence measurements, confluent cell monolayers grown on glass coverslips were used 2-5 days after plating. For electrophysiological recordings, cells grown on plastic tissue culture dishes were trypsinized (0.25% trypsin, 1% EDTA) and replated on 12 mm round glass coverslips. Cells were allowed to adhere for approximately 15 min, then transferred to external solution. Cells were then used for patch clamp recording within 30 min or maintained at 4°C for up to 2 hrs prior to recording.

Mesenteric artery smooth muscle cell (MASMC) isolation

Male Sprague-Dawley rats weighing approximately 300 g were anesthetized with isoflurane (4% by inhalation) and a section of the mesenteric arcade from the small intestine was surgically excised. The rats were then humanely killed by thoracotomy and cardiac extirpation. All experiments were conducted in accordance with the policies of the Public Health Service and the Animal Welfare Act. 10-15 segments (6-10 mm in length) of mesenteric artery were quickly dissected free of connective tissue in ice-cold solution containing (in mM): NaCl 145; KCl 4.7; NaH₂PO₄ 1.2; MgSO₄ 1.17; Pyruvic Acid 2; EDTA 0.02; MOPS 3; CaCl₂ 0.1; D-Glucose 5; with 10 mg/ml bovine serum albumin (BSA). MASMC were isolated essentially as described by (Plane et al., 2005). Arteries were placed in an isolation solution: (in mM) NaCl 120; NaHCO₃ 25; KCl 4.2; MgCl₂ 1.2; KH₂PO₄ 0.6; D-Glucose 11; pH 7.4 when bubbled with 5%CO₂, 21% O₂ (at room temperature) with BSA (1 mg/ml) and 0.1 mM CaCl₂. The arteries were cut into 4-5 mm segments, which were then digested in isolation solution containing papain (1.0 mg/ml), dithiothreitol (1 mg/ml) and BSA (1 mg/ml) for 10 min at 37 °C. The artery segments were then transferred to isolation solution containing collagenase (0.3 mg/ml

collagenase type H and 0.7 mg/ml collagenase type F), 0.01 mM CaCl_2 , dithiothreitol (1 mg/ml), and BSA (1 mg/ml) for 10 min at 37 °C. The tissue was then washed 10 times in ice-cold isolation solution without collagenase, and triturated gently with polished Pasteur pipettes to separate the individual cells. Freshly isolated cells with characteristic elongated morphology of vascular myocytes were added directly to the chamber and allowed to adhere for 10-15 min for patch clamp electrophysiology or collected for preparation of RNA for RT-PCR (see below).

Whole cell voltage clamp

The whole cell ruptured and perforated (200 $\mu\text{g/ml}$ Amphotericin B) patch configurations were used to measure membrane currents under voltage-clamp conditions in single A7r5 cells or MASMC. All experiments were performed at room temperature with continuous perfusion of bath solution. Resistances of patch pipettes were 1.5-2.5 $\text{M}\Omega$ after filling with internal solution. Series resistance was 2-4 $\text{M}\Omega$ in ruptured path experiments and was not compensated.

Solutions for recording store-operated current (I_{SOC}) contained (in mM): 100 Na aspartate (or 100 NMDG aspartate as indicated in figure legends), 20 Ca aspartate, 1 MgCl_2 , 10 HEPES, pH 7.3 (external); 90 Cs aspartate, 20 CsCl, 4.6 MgCl_2 , 5 HEPES, 10 BAPTA- Cs_4 , pH 7.2 (internal). For divalent cation selectivity experiments, external solutions contained (in mM): 100 NMDG aspartate, 20 CaCl_2 , 10 HEPES, pH 7.3, or 20 CaCl_2 was replaced with equimolar concentrations of SrCl_2 , BaCl_2 or MnCl_2 .

Osmolality was adjusted to 265-270 mOsm/l for A7r5 cells and 290 mOsm/l for MASMC with D-glucose. 10 μM verapamil and 100 μM spermine chloride were added to all bath solutions to block L-type Ca^{2+} channels and Mg^{2+} -inhibitable currents, respectively, as previously described (Brueggemann et al., 2005). Currents were recorded with a 100 ms voltage ramp protocol (from +85 mV to -115 mV) from -15 mV holding potential every 10 sec; at least 1 min (6 time points) of stable current recording was used for current density measurements. Liquid junction potentials were

calculated using Junction Potential Calculator provided by PCLAMP8 software and subtracted off-line. Experiments in whole cell perforated patch configuration were started with series resistance (R_s) below 30 M Ω ; cells with an abrupt decrease in R_s were discarded. Recordings of I_{SOC} in ruptured patch configuration were started 30 sec after break-in. Voltage-clamp command potentials were generated using an Axopatch 200B amplifier under control of PCLAMP8 software. Whole-cell currents were digitized at 10 kHz, filtered at 5 kHz and analyzed off-line. Currents were normalized to membrane capacitance. Data are presented as mean \pm standard error of the mean (S.E.).

A well accepted characteristic of store-operated currents is that they are inhibited by lanthanides such as Gd^{3+} . CCE in A7r5 cells is also reportedly inhibited by low concentrations of Gd^{3+} ($<5 \mu M$; Broad et al., 1999; Moneer and Taylor, 2002; Moneer et al., 2003; Moneer et al., 2005). As described previously (Brueggemann et al., 2005), the Gd^{3+} -sensitivity of I_{SOC} in A7r5 cells was evaluated by addition of 100 μM $GdCl_3$ to the external solution. We estimate that the final concentration of free Gd^{3+} is in the 1 to 5 μM range (calculated with WINMAXC software using Gd-aspartate binding constants kindly provided by Chris Patton, Stanford University). For some experiments, 7 mM $GdCl_3$ was added to provide a free $[Gd^{3+}]$ of about 100 μM (also calculated using WINMAXC software).

Thapsigargin was used in several experiments to passively empty intracellular Ca^{2+} stores. Its effectiveness in this regard has been demonstrated previously using fura-2 fluorescence techniques (Byron and Taylor, 1995). In control experiments, we have confirmed that sustained thapsigargin-activated store-operated currents persist after its removal from the bathing medium and are not different if thapsigargin is continuously present.

$[Ca^{2+}]_i$ measurements with fura-2

Essentially as described previously (Fan and Byron, 2000), A7r5 cells were grown to confluence on glass coverslips. Coverslips were washed twice with control medium (135 mM NaCl,

5.9 mM KCl, 1.5 mM CaCl_2 , 1.2 mM MgCl_2 , 11.5 mM glucose, 11.6 mM HEPES, pH 7.3) and then incubated in the same medium with 5 μM fura-2-AM, 0.1% bovine serum albumin, and 0.02% Pluronic F127 detergent for 90 min at room temperature (22-25 $^\circ\text{C}$) in the dark. The cells were then washed twice and incubated in the dark in control medium (or pretreated with drugs) for 0.5-5 h prior to the start of the experiment. All experiments were performed at room temperature. Fura-2 fluorescence (excitation alternating between 340 and 380 nm, emitted fluorescence detected at 509 nm) was measured in cell populations with a Perkin-Elmer LS50B fluorescence spectrophotometer or a BioTek fluorescence plate reader (used only for 2-APB, Gd^{3+} , and diethylstilbestrol (DES) concentration-response determinations). For the latter, A7r5 cells were plated on 96-well plates, grown to confluence, and loaded with fura-2 as described above. Immediately prior to each recording, the control medium was replaced with 100 μl of nominally Ca^{2+} -free control medium (CaCl_2 was omitted) \pm varying concentrations of 2-APB, Gd^{3+} , or DES. Two injectors were used, first to inject thapsigargin (\pm 2-APB, Gd^{3+} , or DES in a volume of 100 μl of Ca^{2+} -free medium at time = 60s) and then CaCl_2 (plus thapsigargin \pm 2-APB, Gd^{3+} , or DES; 100 μl of a 4.5 mM CaCl_2 solution to restore external $[\text{Ca}^{2+}]$ to 1.5 mM at time 13 min); fura-2 fluorescence was recorded at 5s intervals. Each concentration of 2-APB, Gd^{3+} , or DES was tested in triplicate in 3-4 separate experiments. All $[\text{Ca}^{2+}]_i$ recordings were made in the continuous presence of 10 μM verapamil to eliminate any contributions of voltage-gated L-type Ca^{2+} channels.

RT-PCR detection and quantification of TRPC homologue expression

Total RNA was prepared from cultured A7r5 cells or freshly isolated mesenteric artery smooth muscle cells (MASMC) by the guanidinium method (Chomczynski and Sacchi, 1987). Reverse transcription was performed using 1 μg RNA and MMLV-RT according to the manufacturer's protocol (Life Technologies Inc., Gaithersburg, MD). One tenth of the reverse transcription reaction product was used for PCR using Platinum PCR SuperMix (Invitrogen, Carlsbad, CA) and 10 pmol of each

primer according to the manufacturer's protocol. Primer pairs for TRPC1-TRPC7 were as published by McDaniel et al. (2001). Negative controls were minus reverse transcriptase, and positive controls were total rat brain RNA (or in the case of TRPC2, total rat testes RNA). For MASMCMC PCR, individual freshly isolated arterial myocytes (~500) were selected based on their morphology using a micropipette, and placed directly into guanidinium buffer for RNA isolation (Chomczynski and Sacchi, 1987). RNA was divided into eight separate portions then used in RT-PCR reactions using SuperScript One-Step RT-PCR System (Invitrogen, Carlsbad, CA) and primer pairs for TRPC1 through TRPC7. One tenth of the PCR reaction was run on 0.8% agarose/TBE gels. For real time RT-PCR, RNA was prepared from cultured A7r5 cells using RNeasy Minikit (Qiagen, Valencia, CA), and RT performed using i-Script cDNA Synthesis Kit (BioRad, Hercules, CA). cRNA was subjected to real time PCR using Platinum Quantitative PCR SuperMix-UDG w/ROX (Invitrogen, Carlsbad, CA) and TaqMan® Gene Expression Assays and 18s RNA endogenous controls with a 7300 Real-Time PCR System (Applied Biosystems, Foster City, CA). Results were analyzed using RQ Relative Quantitation software (Applied Biosystems, Foster City, CA).

Western blot analysis of TRPC proteins

Confluent cultures of A7r5 cells or MASMCMC were lysed in a solution containing 1% sodium deoxycholate, 0.1% SDS, 1% Triton X-100, 100 mM NaF, 10 mM sodium pyrophosphate, 1 mM EGTA, 1.5 mM MgCl₂, 10% glycerol, 150 mM NaCl, 10 µg/ml leupeptin, 10 µg/ml aprotinin, 1 mM Na₃VO₄, 50 mM HEPES, pH 7.4. Following centrifugation to remove particulate material, protein concentration in supernatants was determined using a bicinchoninic acid protein assay kit (Pierce) and equal amounts of protein (100 – 400 µg) were separated by SDS-PAGE on a 7.5% acrylamide gel. The proteins were then transferred to nitrocellulose and blotted for TRPC1, TRPC4, and/or TRPC6 using commercially available antibodies from Alomone Labs (1:500 dilution). In additional experiments (results not shown), duplicate samples blotted using antibodies preadsorbed with immunogenic peptide

served as negative controls to confirm correct identification of the bands of interest. Chemiluminescent detection of proteins and densitometric analysis of blots was performed as previously described (Byron and Lucchesi, 2002).

TRPC1 siRNA Construct

A vector containing a neomycin-resistance gene and expressing rat TRPC1 siRNA was generously provided by Dr. M.L. Villereal (University of Chicago). This construct was previously demonstrated to selectively reduce TRPC1 expression in H19 -7 hippocampal neuronal cells (Wu et al., 2004) and HEK-293 cells (TRPC3, TRPC4, TRPC6, and TRPC7 levels were not affected; Zagranichnaya et al., 2005). A7r5 cells were transfected with the rat siTRPC1 construct and grown in selection medium containing G418 (500 µg/ml) to establish a stable cell line.

Transfection

Proliferating A7r5 cells were grown in 150-mm cell culture dishes to 50% confluence and transfected the next day with 20 µg plasmid DNA using Lipofectamine transfection reagent (Invitrogen, Inc.) according to standard transfection protocol. 48 h after transfection, 500 µg/ml G418 was added. Parallel cultures of non-transfected A7r5 cells were also treated with G418 to verify its lethality in the absence of the selection marker; these cultures died within five days. Surviving transfected cells were maintained in the presence of G418 for several weeks.

TRPC1 antisense adenovirus

We constructed an adenoviral vector for expression of antisense TRPC1 DNA and co-expression of Green Fluorescent Protein (AS-TRPC1-GFP). A 1694-bp portion of the human TrpC1 cDNA (generously provided by Dr. I. Ambudkar, NIH) was inserted in the antisense direction into the adenovirus GFP-tagged shuttle vector pAdTrack-CMV (generously provided by Dr. B. Vogelstein, Johns Hopkins University). Recombinant virus was generated using the AdEasy system essentially as

described by He et al. (1998), using electrocompetent BJ5183-AD-1 cells (Stratagene, La Jolla, CA). A7r5 cells were infected with AS-TRPC1-GFP at a multiplicity of 30:1, and cells were analyzed 4 to 6 days later by electrophysiology and RT-PCR as described above.

Immunohistochemical detection of TRPC1

A7r5 cells were subcultured on 12 mm round glass coverslips and infected with AS-TRPC1-GFP at a multiplicity of infection of 100. After 5 days, the cells were fixed with 2% paraformaldehyde in phosphate buffered saline (PBS: 138 mM NaCl, 2.7 mM KCl, 8.1 mM Na₂HPO₄, 1.2 mM KH₂PO₄, pH 7.4) for 15 min, washed twice with PBS and permeabilized with 0.5% Triton X-100 (Sigma) in PBS for 15 min. After washing with 0.1% Triton X-100 in PBS, coverslips were blocked with Image-iTTM FX Signal Enhancer (Molecular Probes Inc., Eugene, OR) according to the manufacturer's instructions for 30 min. An additional blocking step with 10% goat serum for 2 hr was used for staining with anti-TRPC1 (H-105) rabbit polyclonal IgG antibody (Santa Cruz Biotechnology, Inc.). Coverslips were washed 3 times with 0.1% Triton X-100 in PBS and incubated with anti-TRPC1 antibodies either generously provided by Dr. M.L. Villereal (University of Chicago) or commercially available (Santa Cruz Biotechnology, Inc.) at a 1:100 dilution over night at +4°C. Coverslips incubated without primary antibody were used to control for non-specific background staining. After three washing steps with 0.1% Triton X-100 in PBS for 15 min each, coverslips were incubated for 2 hr at room temperature in the dark with Alexa Fluor 594 goat anti-rabbit secondary antibody (1:400 dilution in 0.1% Triton X-100 plus 10% goat serum in PBS). Three additional washing steps for 15 min with 0.1% Triton X-100 in PBS were performed before mounting coverslips on glass slides using SlowFade Light Antifade Kit (Molecular Probes Inc., Eugene, OR). Cell images were acquired using C Imaging System (Compix Inc.) with an Olympus 1X71 inverted epifluorescence microscope (10x fluorescent objective) and Simple PCI software (Vers.5.3.1.). Two images of each field were captured, at 490 nm

excitation wavelength for GFP fluorescence, and at 595 nm excitation wavelength for Alexa Fluor 594 fluorescence, respectively.

Statistics

Data are expressed as means \pm standard error of the mean (S.E.) Statistical comparisons were performed using Sigma Stat 3.1 (Systat Software, Inc.). Differences between two groups were evaluated with Student's paired or unpaired *t* test as appropriate. Concentration-response relationships were analyzed by One Way Repeated Measures Analysis of Variance and Holm-Sidak method for comparisons with controls. Differences were considered to be significant when $p < 0.05$.

Materials

Rabbit polyclonal anti-TRPC1, TRPC4, and TRPC6 antibodies were from Alomone Laboratories (Jerusalem, Israel); PCR primers were from Integrated DNA Technologies, Inc. (Coralville, IA); cyclopiazonic acid and 1-oleoyl-2-acetyl-sn-glycerol were from Biomol Research Laboratories, Inc. (Plymouth Meeting, PA); collagenase types H and F, diethylstilbestrol, gadolinium chloride, flufenamic acid, [Arg⁸]-vasopressin, verapamil, and thapsigargin were from Sigma-Aldrich (St. Louis, MO); Fura-2-AM, Alexa Fluor 594 anti-rabbit IgG, SlowFade Light Antifade Kit, and Pluronic F127 were from Molecular Probes, Inc. (Eugene, OR). Papain was from Worthington Biochemical Corporation (Lakewood, NJ). Anti-TRPC1 rabbit polyclonal IgG was from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA).

RESULTS:

In a previous study (Brueggemann et al., 2005), we identified patch clamp recording conditions for detection of store-operated currents (I_{SOC}) in A7r5 cells. Under these recording conditions (perforated patch configuration in 0 Na^+ , 20 mM Ca^{2+} external solution), I_{SOC} is activated rapidly in response to emptying of intracellular Ca^{2+} stores with 100 nM AVP, reaching a peak within 1.5 min followed by partial inactivation (Fig. 1A). The current declines slowly following washout of AVP. As described previously (Brueggemann et al., 2005), I_{SOC} has an inwardly rectifying current-voltage (I - V) relationship and reversal potential near +40 mV (Fig. 1B). Under the same recording conditions, I_{SOC} was also detected in freshly isolated mesenteric artery smooth muscle cells (MASMC, Fig. 1 C,D). Small inwardly rectifying currents were detected in 5 of 7 cells exposed to thapsigargin (1 μM); mean current amplitude at -115 mV was estimated as -0.37 ± 0.14 pA/pF ($n=4$).

We reported previously (Brueggemann et al., 2005) that I_{SOC} developed in A7r5 cells as a result of passive emptying of Ca^{2+} stores with 1 μM thapsigargin (irreversible inhibitor of the sarcoplasmic reticulum Ca^{2+} ATPase) or by dialyzing cells with 10 mM BAPTA in the ruptured patch configuration. A number of pharmacological agents have been used to characterize store-operated currents or Ca^{2+} entry pathways in a variety of cell types (reviewed by Parekh and Putney, 2005). We previously demonstrated (Brueggemann et al., 2005) that 100 μM GdCl_3 in aspartate-based external solution (about 5 μM free Gd^{3+}) completely inhibited thapsigargin-activated I_{SOC} . Here we show that 5 μM free Gd^{3+} inhibits I_{SOC} in A7r5 cells reversibly. A significant recovery, to $88 \pm 7\%$ of its original level, was recorded after washout of Gd^{3+} ($n=6$, $p<0.001$, paired Student's t test (Fig. 2A)). 2-APB also significantly inhibited Gd -sensitive I_{SOC} (by $95 \pm 2\%$, $n=4$, $p<0.001$, paired Student's t test) and this effect partially reversed after washout (current returned to $56 \pm 17\%$ of its original level, $n=4$). At a concentration of 10 μM , inhibition by 2-APB was reproducibly slower than by Gd^{3+} (Fig.2B). We also found that diethylstilbestrol (DES), a drug recently reported to inhibit capacitative Ca^{2+} entry in

smooth muscle cells (Zakharov et al., 2004), was a very effective inhibitor of I_{SOC} in A7r5 cells (Fig. 2C). 10 μ M DES applied externally for 10-20 min significantly inhibited $96 \pm 3\%$ ($p < 0.001$, paired Student's t test) of Gd-sensitive I_{SOC} with $93 \pm 15\%$ recovery during washout for 15-25 min ($n=4$, Fig. 2C).

Potential activators of non-selective cation channels were also tested for effects on I_{SOC} . Flufenamate has been used to distinguish among members of the TRP channel family and diacylglycerol analogs have been reported to activate non-selective cation channels, including store-operated channels in some cases (see below). I_{SOC} activated by passive depletion of Ca^{2+} stores either with thapsigargin ($n=3$) or with BAPTA ($n=4$, Fig. 2D) was partially inhibited by 100 μ M flufenamate ($80 \pm 16\%$ inhibition of thapsigargin-activated I_{SOC} , $p < 0.05$, and $44 \pm 7\%$ inhibition of BAPTA-activated I_{SOC} , $P < 0.05$, paired Student's t test). Inhibition of thapsigargin-activated I_{SOC} by flufenamate was not significantly different from inhibition of BAPTA-activated I_{SOC} by flufenamate ($p > 0.05$, Student's t test). I_{SOC} was not activated by the soluble diacylglycerol analog, 1-oleoyl-2-acetyl-sn-glycerol in perforated patch recordings (OAG, 100 μ M, $n=4$, Fig. 2E).

We previously found I_{SOC} to be highly selective for Ca^{2+} over Na^{+} (Brueggemann et al., 2005). In the present study the selectivity of I_{SOC} was evaluated for Ca^{2+} , Ba^{2+} , Sr^{2+} and Mn^{2+} , divalent cations commonly used for fura-2 fluorescence measurements of CCE. In response to equimolar substitution of 20 mM Ba^{2+} , Sr^{2+} , or Mn^{2+} for Ca^{2+} in NMDG-containing external solution, the I_{SOC} amplitudes measured at -115 mV were in the following order relative to Ca^{2+} : Sr^{2+} (1.7 ± 0.2 , $n=6$) $>$ Ba^{2+} (1.3 ± 0.1 , $n=5$) \geq Ca^{2+} (1) \gg Mn^{2+} (0.13 ± 0.10 , $n=4$) (Fig. 3). The mean amplitude of the current in 20 mM Ca^{2+} was 0.35 ± 0.07 pA/pF ($n=15$).

Fura-2 fluorescence studies

To further investigate whether I_{SOC} measured by whole cell patch clamp corresponds to the capacitative Ca^{2+} entry pathway (CCE) previously described (Byron and Taylor, 1995; Broad et al.,

1999; Moneer and Taylor, 2002; Moneer et al., 2003; Brueggemann *et al.*, 2005; Moneer et al., 2005), we used fura-2 fluorescence techniques to compare the pharmacological properties of the Ca^{2+} entry pathway under the same ionic conditions used for I_{SOC} measurements.

In A7r5 cells pretreated with 1 μM thapsigargin for 30-60 min to deplete intracellular Ca^{2+} stores, and then using the same external solutions used for electrophysiological detection of I_{SOC} , we found that capacitative Ca^{2+} entry (CCE) was reversibly blocked by 2-APB (10 μM , Fig. 4A). The decline in $[\text{Ca}^{2+}]_i$ was reproducibly slower and to a slightly lesser extent than that induced by Gd^{3+} , mimicking the pattern of inhibition of I_{SOC} by these same treatments (Fig. 2B). In several experiments, a transient increase in $[\text{Ca}^{2+}]_i$ preceded the decline in $[\text{Ca}^{2+}]_i$ on addition of 2-APB (for examples see Figs. 5B and 6A, representative of 5 of 13 experiments). A similar pattern was occasionally observed during measurements of 2-APB effects on I_{SOC} (not shown). The concentration-dependence for 2-APB was evaluated in cells exposed to thapsigargin in Ca^{2+} -free medium followed by re-exposure to external Ca^{2+} (Fig. 4B,C). 2-APB had no effect on basal $[\text{Ca}^{2+}]_i$ or release of intracellular Ca^{2+} , but it inhibited the sustained phase of the $[\text{Ca}^{2+}]_i$ response after re-exposure to external Ca^{2+} with an IC_{50} of $\approx 2.2 \mu\text{M}$ (Fig. 4C).

Gd^{3+} (approximately 5 μM free Gd^{3+} in the aspartate-based solution, see *METHODS*) inhibited CCE reversibly (Fig. 4A). The concentration-dependence for inhibition of thapsigargin-stimulated CCE by Gd^{3+} is shown in Fig. 5A ($\text{IC}_{50} \approx 600 \text{ nM}$). The effects of Gd^{3+} on CCE were previously reported to be irreversible (Broad et al., 1999). The latter studies utilized a chloride-based external solution for recording fura-2 fluorescence. When we tested the effects of 5 μM Gd^{3+} in a Cl-based external solution (control medium used for loading cells with fura-2; see *METHODS*), we also found that its effects on both I_{SOC} (not shown) and CCE were very poorly reversible (Fig. 5B). Reversibility of Gd^{3+} -block in the aspartate-based solution may be due to chelation of Gd^{3+} by aspartate.

Diethylstilbestrol (DES) is an estrogen analog, which has been recently reported to potentially block CCE in vascular smooth muscle cells (Zakharov et al., 2004). DES also effectively inhibited CCE in A7r5 cells (Fig. 6A), with an IC_{50} of ≈ 800 nM (Fig. 6B).

There is some controversy related to the contributions of different Ca^{2+} entry pathways to AVP-activated Ca^{2+} signals (Brueggemann et al., 2005). We therefore evaluated the effects of various I_{SOC} /CCE inhibitors on the sustained phase of the Ca^{2+} response to 1 nM AVP under the same external ionic conditions used for recording I_{SOC} . As shown in Fig. 7, 2-APB, Gd^{3+} , and DES were similarly effective in reducing the AVP-stimulated Ca^{2+} signals as they were for CCE activated by thapsigargin pretreatment (Fig. 6A). These results suggest a substantial contribution of CCE to the AVP-induced Ca^{2+} response.

The decrease in the sustained phase of the AVP-induced Ca^{2+} signal by DES was transient. In the continued presence of DES, $[Ca^{2+}]_i$ gradually rebounded to greater than its initial level (Fig. 7A). This did not appear to be due to transient block of CCE because the same effect was observed when DES was applied in the presence of Gd^{3+} at a concentration (5 μ M free Gd^{3+}) that stably blocks both I_{SOC} and CCE (Fig. 7B). Increasing external Gd^{3+} concentration to achieve a free concentration of approximately 100 μ M (7 mM total $GdCl_3$) was also ineffective in preventing the DES-induced increase in $[Ca^{2+}]_i$ (Fig. 7C). A similar slow rebound of $[Ca^{2+}]_i$ was apparent when DES was added to thapsigargin-pretreated cells in the presence of Gd^{3+} (not shown).

TRP channel expression

The identities of the channels responsible for non-selective cation currents in A7r5 cells may be related to expression of TRPC homologues. Using RT-PCR, we have detected TRPC1, TRPC4, and TRPC6 mRNA in A7r5 cells. As shown in Fig. 8A, total RNA prepared from A7r5 cells (upper panel), freshly isolated rat mesenteric artery smooth muscle cells (MASMC, middle panel), and adult rat brain/testis (lower panel) was reverse transcribed and subjected to PCR using primers specific for

TRPC1 through TRPC7 (lanes 1-7). Using rat testis as a positive control for TRPC2 or rat brain as a positive control for the other TRPC homologues, TRPC2, TRPC3, TRPC5, and TRPC7 were undetectable in A7r5 cells, whereas TRPC3 and TRPC5 were faintly detected in MASMC. Western blot analysis using commercially available anti-TRPC antibodies (Fig. 8B), revealed that TRPC1, TRPC4, and TRPC6 proteins are detected in both A7r5 cell and MASMC lysates.

TRPC1 has previously been implicated in store-operated Ca^{2+} entry in vascular smooth muscle cells (Golovina et al., 2001; Xu and Beech, 2001; Sweeney et al., 2002). We used an siRNA strategy previously demonstrated by Wu et al. (2004) to selectively depress TRPC1 expression in H19-7 hippocampal neuronal cells. In A7r5 cells selected with G418 for stable expression of the TRPC1 siRNA construct (see *METHODS*), expression of TRPC1 mRNA was significantly decreased (by $42 \pm 4\%$, $n=3$, $p=0.01$, Student's t test, Fig. 9A), and TRPC1 protein levels, based on Western blot analysis, were significantly reduced (by $62 \pm 5\%$, $p<0.001$, $n=6$, Student's paired t test compared with control cells maintained in parallel cultures without G418). In 3 experiments, Western blotting for all three TRPC homologues (TRPC1, TRPC4, and TRPC6, Fig. 9B) revealed no significant reductions in TRPC6 or TRPC4 expression (reductions of $14 \pm 27\%$ and $14 \pm 13\%$, respectively, compared with control cultures). In the same samples there was a consistent and significant reduction in TRPC1 protein expression ($56 \pm 6\%$ less than controls, $p<0.05$). Store-operated current (I_{SOC}) was also significantly decreased (by $51 \pm 6\%$, $n=8$, $p<0.001$, Student's t test, Fig. 9C). Similarly, A7r5 cells infected with an adenovirus to express TRPC1 antisense sequences and green fluorescent protein (AS-TRPC1-GFP) exhibited significantly depressed I_{SOC} compared with non-GFP-expressing cells in the same cultures (Fig. 9D-F). 20 to 30% of A7r5 cells infected for 4 to 6 days with AS-TRPC1-GFP (30 MOI) exhibited green fluorescence (Fig 9D). TRPC1 immunostaining (detected using either a commercial anti-TRPC1 antibody or anti-TRPC1 antibodies obtained from M.L. Villereal—see *METHODS*) was reduced in the GFP-positive cells (Fig. 9D) and I_{SOC} was significantly reduced (by

44%) in cells with GFP fluorescence versus non-fluorescent cells (n=15 and 16, respectively, $p<0.001$, Student's t test, Fig. 9E,F).

DISCUSSION:

Our electrophysiological and pharmacological studies suggest that I_{SOC} , activated either by AVP or by passive depletion of Ca^{2+} stores, corresponds to CCE in A7r5 cells. I_{SOC} is a relatively Ca^{2+} -selective current, which is insensitive to LOE 908, but blocked by micromolar Gd^{3+} (Brueggemann et al., 2005). These properties are consistent with the CCE pathway previously measured in A7r5 cells using fura-2 fluorescence techniques (Broad et al., 1999; Moneer and Taylor, 2002) or in cerebral arteriolar myocytes using fura-PE3 or fluo-4 fluorescence (Flemming *et al.*, 2003). Additional pharmacological agents, 2-APB and DES, that have been used as selective blockers of store-operated currents or Ca^{2+} entry in A7r5 or other cell systems were also effective blockers of both I_{SOC} and CCE measured under identical external ionic conditions in A7r5 cells.

I_{SOC} in A7r5 cells compared with other VSMC

I_{SOC} recorded in A7r5 cells displays inward rectification of its *I-V* relationship in high external Ca^{2+} and a positive reversal potential (Fig. 1B) like that recorded in choroidal arteriolar myocytes (Curtis and Scholfield, 2001), but unlike the linear *I-V* relationship and 0 mV reversal of thapsigargin-activated I_{SOC} from mouse or rabbit aortic smooth muscle cells (Trepakova et al., 2001) or pulmonary artery myocytes (Golovina et al., 2001; McDaniel et al., 2001). Differences in *I-V* relationships may be attributable to different natures of I_{SOC} from different VSM preparations or to different recording conditions used by different groups. Ca^{2+} -selective and non-selective store-operated channels may even exist within the same cells (Bolotina, 2004; Vanden Abeele et al., 2004; Bugaj et al., 2005), though in A7r5 cells we find that depletion of Ca^{2+} stores activates only an inwardly rectifying, relatively Ca^{2+} -selective current measured by whole cell voltage clamp (Brueggemann et al., 2005). We have also detected a similar current in freshly isolated mesenteric artery smooth muscle cells using the same recording conditions (Fig. 1C).

In rabbit portal vein myocytes, a moderately Ca^{2+} -selective I_{SOC} (Helliwell and Large, 1997; Albert and Large, 2002a; Albert and Large, 2002b) is activated by OAG and pharmacological studies have implicated protein kinase C (PKC) as an essential step in its activation by either passive store depletion or by noradrenaline (Albert and Large, 2002b). In contrast, I_{SOC} in A7r5 cells is not activated by OAG (Fig. 2E), nor do PKC inhibitors prevent store-operated Ca^{2+} entry (Broad et al., 1999), suggesting that distinct channels or regulatory mechanisms may be involved.

I_{SOC} identification and relationship to CCE

Considering that currents identified as “store-operated” in various vascular smooth muscle preparations vary significantly in their electrophysiological characteristics, it is important to consider the basis for their identification as store-operated currents. In A7r5 cells, I_{SOC} is activated by both physiological and pharmacological treatments that lead to store depletion and its activation is reversed under conditions in which the stores are allowed to refill (Brueggemann *et al.*, 2005). Furthermore, pharmacological agents that distinguish store-operated from non-store-operated divalent cation influx measured with fura-2 similarly distinguish I_{SOC} from other cation currents. Few other studies have met these criteria using vascular smooth muscle cells.

In A7r5 cells, CCE activation has been demonstrated for both passive and active depletion of Ca^{2+} stores using fura-2 fluorescence to examine Ca^{2+} , Ba^{2+} , Mn^{2+} , and Sr^{2+} influx (Byron and Taylor, 1995). Its activation is reversed by refilling of Ca^{2+} stores (Byron and Taylor, 1995; Brueggemann *et al.*, 2005). Cyclopiazonic acid (CPA) is a reversible inhibitor of the SR Ca^{2+} -ATPase and we have shown that I_{SOC} in A7r5 cells is activated by CPA and deactivates after washout of CPA (Brueggemann *et al.*, 2005). Other studies have used CPA to activate I_{SOC} in vascular myocytes but failed to demonstrate that this effect reverses on removal of CPA (Albert and Large, 2002a,b; Ng and Gurney, 2001; Sweeney *et al.*, 2002; Bergdahl *et al.*, 2005).

In some cases, pharmacological inhibition of CPA- or thapsigargin-stimulated Ca^{2+} entry characterized using fluorescent indicators (Flemming *et al.*, 2003) does not match the effects of the same agents on store-operated currents in cells from the same tissue (Bergdahl *et al.*, 2005). In other cases (e.g. Ng and Gurney, 2001; McDaniel *et al.*, 2001), CCE and inward current were similarly inhibited by agents that may not distinguish the store-operated influx pathway from other cation entry pathways that may be active in the vascular myocytes.

Physiological activation of I_{SOC}

It is generally accepted that Ca^{2+} influx via the store-operated pathway may contribute to the refilling of intracellular stores following InsP_3 -mediated Ca^{2+} release. It is important therefore to confirm that I_{SOC} is induced by agonists that activate phospholipase C and that this current corresponds to currents activated by other treatments that deplete intracellular Ca^{2+} stores.

In rabbit portal vein myocytes, noradrenaline was found to activate a current with properties similar to a CPA-activated current (Albert and Large, 2002a), but the current was activated under conditions that were not consistent with store depletion and the authors concluded that the effects of noradrenaline were store-independent. In pulmonary artery myocytes, phenylephrine (PE) and CPA activated similar currents (McDaniel *et al.*, 2001). However, the conditions used by McDaniel *et al.* (2001) for recording CPA- and PE-activated currents would favor development of magnesium-inhibited currents (I_{MIC}), which, as noted by Bakowski and Parekh (2002), had likely been a contaminant of store-operated currents in some earlier studies.

DES has been recently described as an effective inhibitor of store-operated currents and CCE at concentrations which have no effect on I_{MIC} mediated by TRPM7 (Zakharov *et al.*, 2004). Our results suggest a somewhat less potent inhibitory effect of DES on CCE in A7r5 cells and indicate that at concentrations that maximally inhibit CCE (10 μM), DES has additional effects on Ca^{2+} transport that may complicate its use as a CCE inhibitor. DES was also previously reported to inhibit L-type Ca^{2+}

currents in A7r5 cells with an $IC_{50} < 10 \mu M$, and, at a concentration of $30 \mu M$, DES inhibited AVP-activated non-selective cation currents and K^+ currents (Nakajima et al., 1995), providing additional evidence that the effects of DES are not particularly selective.

I_{SOC} , I_{CAT} , and AVP-stimulated Ca^{2+} entry in A7r5 cells

In A7r5 cells, AVP was previously found to activate both CCE and NCCE (Byron and Taylor, 1995; Broad et al., 1999). Previous electrophysiological studies identified a non-selective cation current (I_{CAT}) in A7r5 cells activated by high [AVP] ($100 nM$; Van Renterghem et al., 1988; Krautwurst et al., 1994; Nakajima et al., 1996; Iwasawa et al., 1997; Jung et al., 2002), which may relate to AVP-activated NCCE. I_{CAT} is clearly distinct from I_{SOC} in several ways: 1) OAG, was reported to directly activate I_{CAT} in A7r5 cells (Jung et al., 2002), but OAG did not activate I_{SOC} (Fig. 2C); 2) I_{CAT} activated by $100 nM$ AVP in A7r5 cells was inhibited by LOE 908 (Krautwurst et al., 1994), but LOE 908 does not inhibit I_{SOC} (Brueggemann et al., 2005); 3) AVP-activated I_{CAT} is enhanced by flufenamate (Jung et al., 2002), whereas I_{SOC} is inhibited by flufenamate (Fig. 2B); 4) I_{CAT} is a non-selective cation current (Iwasawa et al., 1997; Jung et al., 2002), whereas I_{SOC} appears to be highly selective for Ca^{2+} over Na^+ (Brueggemann et al., 2005).

Using fura-2 fluorescence to examine divalent cation entry pathways in A7r5 cells, Byron and Taylor (1995) found that both thapsigargin- and AVP-stimulated Ca^{2+} entry were inhibited by other divalent cations ($Be^{2+} > Zn^{2+} > Ni^{2+} > Sr^{2+}$) with the same relative effectiveness. They concluded that either CCE was the predominant contributor to the sustained phase of the AVP-activated Ca^{2+} response or both CCE and NCCE have similar sensitivities to these cation blockers. In the present study we have found that both thapsigargin- and AVP-stimulated Ca^{2+} entry are inhibited similarly by pharmacological agents that are reportedly selective for the store-operated pathway, leading us to conclude that CCE contributes substantially to the sustained phase of the AVP-induced Ca^{2+} response. These findings are consistent with our earlier results (Brueggemann et al., 2005), but directly challenge

the model put forward by Moneer and Taylor (2002), which suggests that AVP inhibits CCE in A7r5 cells.

Divalent cation selectivity: electrophysiology (I_{SOC}) versus fura-2 fluorescence (CCE):

Initial studies of the multiple divalent cation entry pathways in A7r5 cells using fura-2 fluorescence suggested that the CCE pathway was relatively less permeable to Sr^{2+} , but more permeable to Mn^{2+} compared with the NCCE pathway (Byron and Taylor, 1995). More recent publications have extended the original interpretation to assert that the CCE pathway is impermeable to Sr^{2+} and that the NCCE pathway is impermeable to Mn^{2+} (Moneer and Taylor, 2002; Moneer et al., 2005). However, fura-2 fluorescence cannot be used to assess absolute conductance. A poorly detectable Sr^{2+} signal compared with a large Ca^{2+} signal when CCE is active (e.g. Fig. 2 of Byron and Taylor (1995)) may reflect the large differences in affinities of Sr^{2+} and Ca^{2+} for fura-2 (Schilling et al., 1989; Kwan and Putney, 1990). Our ionic selectivity measurements for I_{SOC} suggest that this pathway conducts Sr^{2+} at least as well as it does Ca^{2+} (Fig. 3A,C). Mn^{2+} currents, on the other hand, were barely detectable by whole cell patch clamp with 20 mM external Mn^{2+} (Fig. 3B,C), whereas Mn^{2+} entry is readily detected by fura-2 fluorescence even with 200-fold less Mn^{2+} in the external medium (Byron and Taylor, 1995). The greater sensitivity of fura-2 compared with electrophysiology may be attributable to the very high affinity of fura-2 for Mn^{2+} ($K_D = 8.64$ nM; Kwan and Putney, 1990). These differences in sensitivity unfortunately prevent conducting fura-2 Mn^{2+} quench and electrophysiological measurements of I_{SOC} under the same ionic conditions.

TRPC channels

Several TRPC homologues are expressed in A7r5 cells. Jung et al. (2002) detected TRPC1 and TRPC6 by Northern blot. Our own results using RT-PCR and Western blotting (Fig. 8) confirm that finding and identify TRPC4 as another homologue expressed in this cell line and in MASMC. A recent study by Moneer et al. (2005) has used quantitative RT PCR to evaluate TRPC expression in several

strains of A7r5 cells and reported expression of TRPC1 and TRPC6, with less expression of TRPC2, TRPC3, TRPC4, and TRPC5; TRPC7 expression was undetectable.

Given the potential for heterotetrameric assembly of TRPC homologues (Hofmann et al., 2002; Strübing et al., 2003), we should be cautious in assigning a particular current or Ca^{2+} entry pathway to a particular TRPC homologue. Nonetheless, previous studies have suggested that TRPC1 may be involved in capacitative Ca^{2+} entry in VSM (Golovina et al., 2001; Sweeney et al., 2002; Xu and Beech, 2001; Bergdahl et al., 2005). TRPC1 has also been reported to form heterotetramers with TRPC4 (Hofmann et al., 2002) and studies in vascular endothelial cells (reviewed by Cioffi et al. (2003)) suggest that TRPC4 may be required for normal targeting of TRPC1 to the plasma membrane and formation of functional store-operated channels. Interestingly, store-operated currents recorded from those cells exhibit inward rectification and Ca^{2+} selectivity similar to I_{SOC} recorded in A7r5 cells (Cioffi et al., 2003). Our results using siRNA or antisense constructs to reduce expression of TRPC1 (Fig. 9) support a role of TRPC1 in store-operated Ca^{2+} entry in A7r5 cells.

Enhancement of an AVP-stimulated non-selective cation current (I_{CAT}) by flufenamate, its activation by OAG (Hofmann et al., 1999), and evidence of TRPC6 mRNA expression have been cited as evidence that TRPC6 proteins may form the channels through which this current passes in A7r5 cells (Jung et al., 2002). The differences in pharmacological profiles (activation by flufenamate and OAG) and cation selectivity suggest that TRPC6 may form channels that are distinct from those that mediate I_{SOC} /CCE in A7r5 cells. Nonetheless, it seems likely that TRPC6 mediates I_{CAT} and contributes to NCCE in A7r5 and other vascular smooth muscle cells (Inoue et al., 2001). These results emphasize the importance of establishing electrophysiological recording conditions and pharmacological profiles that distinguish among different agonist-stimulated Ca^{2+} entry pathways in vascular smooth muscle cells.

Potential functional significance of Ca^{2+} -permeable cation channel activation by AVP

It is increasingly apparent that Ca^{2+} -permeable cation channels may be important targets for development of new therapeutic agents for the treatment of cardiovascular diseases (Stevens, 2001). Store-operated Ca^{2+} currents were recently found to be correlated with TRPC1 expression in cerebral arteries and store-depletion-induced contractile responses were inhibited by anti-TRPC1 antibodies (Bergdahl et al., 2005). In the same study, TRPC1 expression was found to be upregulated in human internal mammary arteries after dilation by a balloon used for angioplasty. Several studies have suggested a role for TRPC1 in store-operated Ca^{2+} entry and growth of vascular smooth muscle cells (Golovina et al., 2001; Xu and Beech, 2001; Sweeney et al., 2002). Plasticity of TRPC expression and the resulting alterations of cation permeability may relate to pathological conditions such as the proliferation of vascular smooth muscle cells that contributes to restenosis following angioplasty. Pharmacological, electrophysiological, and molecular characterization of store-operated or receptor-activated non-selective cation channels may facilitate the development of more effective therapeutic regimens for the prevention or reversal of cardiovascular diseases.

ACKNOWLEDGEMENTS:

The authors gratefully acknowledge the technical assistance of Patrycja Galazka, John Barakat, and Beverly Martin.

REFERENCES:

- Albert AP and Large WA (2002a) Activation of store-operated channels by noradrenaline via protein kinase C in rabbit portal vein myocytes. *J Physiol* **544**:113-125.
- Albert AP and Large WA (2002b) A Ca^{2+} -permeable non-selective cation channel activated by depletion of internal Ca^{2+} stores in single rabbit portal vein myocytes. *J Physiol* **538**:717-728.
- Bakowski D, and Parekh AB (2002) Permeation through store-operated CRAC channels in divalent-free solution: potential problems and implications for putative CRAC channel genes. *Cell Calcium* **32**:379–391.
- Bergdahl A, Gomez MF, Wihlborg AK, Erlinge D, Eyjolfson A, Xu SZ, Beech DJ, Dreja K and Hellstrand P (2005) Plasticity of TRPC expression in arterial smooth muscle: correlation with store-operated Ca^{2+} entry. *Am J Physiol - Cell Physiology* **288**:C872-880.
- Bolotina VM (2004) Store-operated channels: diversity and activation mechanisms. *Science's Stake [Electronic Resource]: Signal Transduction Knowledge Environment* **2004**:pe34.
- Broad LM, Cannon TR and Taylor CW (1999) A non-capacitative pathway activated by arachidonic acid is the major Ca^{2+} entry mechanism in rat A7r5 smooth muscle cells stimulated with low concentrations of vasopressin. *J Physiol* **517**:121-134.
- Brueggemann LI, Markun DR, Barakat JA, Chen H and Byron KL (2005) Evidence against reciprocal regulation of Ca^{2+} entry by vasopressin in A7r5 rat aortic smooth-muscle cells. *Biochem J* **388**:237-244.
- Bugaj V, Alexeenko V, Zubov A, Glushankova L, Nikolaev A, Wang Z, Kaznacheyeva E, Bezprozvanny I and Mozhayeva GN (2005) Functional properties of endogenous receptor- and store-operated calcium influx channels in HEK293 cells. *J Biol Chem* **280**:16790-16797.
- Byron K and Taylor CW (1995) Vasopressin stimulation of Ca^{2+} mobilization, two bivalent cation entry pathways and Ca^{2+} efflux in A7r5 rat smooth muscle cells. *J Physiol* **485**:455-468.

- Byron KL and Lucchesi PA (2002) Signal transduction of physiological concentrations of vasopressin in A7r5 vascular smooth muscle cells. A role for PYK2 and tyrosine phosphorylation of K^+ channels in the stimulation of Ca^{2+} spiking. *J Biol Chem* **277**:7298-7307.
- Byron KL and Taylor CW (1993) Spontaneous Ca^{2+} spiking in a vascular smooth muscle cell line is independent of the release of intracellular Ca^{2+} stores. *J Biol Chem* **268**:6945-6952.
- Chomczynski P and Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* **162**:156-159.
- Cioffi DL, Wu S and Stevens T (2003) On the endothelial cell I(SOC). *Cell Calcium* **33**:323-336.
- Curtis TM and Scholfield CN (2001) Nifedipine blocks Ca^{2+} store refilling through a pathway not involving L-type Ca^{2+} channels in rabbit arteriolar smooth muscle. *J Physiol* **532**:609-623.
- Fan J and Byron KL (2000) Ca^{2+} signalling in rat vascular smooth muscle cells: a role for protein kinase C at physiological vasoconstrictor concentrations of vasopressin. *J Physiol* **524 Pt 3**:821-831.
- Flemming R, Xu SZ and Beech DJ (2003) Pharmacological profile of store-operated channels in cerebral arteriolar smooth muscle cells. *Br J Pharmacol* **139**: 955-965.
- Golovina VA, Platoshyn O, Bailey CL, Wang J, Limsuwan A, Sweeney M, Rubin LJ and Yuan JX (2001) Upregulated TRP and enhanced capacitative Ca^{2+} entry in human pulmonary artery myocytes during proliferation. *Am J Physiol - Heart & Circulatory Physiology* **280**:H746-755.
- He TC, Zhou S, da Costa LT, Yu J, Kinzler KW and Vogelstein B (1998) A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci USA* **95**:2509-2514.
- Helliwell RM and Large WA (1997) Alpha 1-adrenoceptor activation of a non-selective cation current in rabbit portal vein by 1,2-diacyl-sn-glycerol. *J Physiol* **499**:417-428.
- Hofmann T, Obukhov AG, Schaefer M, Harteneck C, Gudermann T and Schultz G (1999) Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature* **397**:259-263.

- Hofmann T, Schaefer M, Schultz G and Gudermann T (2002) Subunit composition of mammalian transient receptor potential channels in living cells. *Proc Natl Acad Sci USA* **99**:7461-7466.
- Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S, Ito Y and Mori Y (2001) The transient receptor potential protein homologue TRP6 is the essential component of vascular $\alpha(1)$ -adrenoceptor-activated Ca^{2+} -permeable cation channel. *Circ Res* **88**:325-332.
- Iwamuro Y, Miwa S, Zhang XF, Minowa T, Enoki T, Okamoto Y, Hasegawa H, Furutani H, Okazawa M, Ishikawa M, Hashimoto N and Masaki T (1999) Activation of three types of voltage-independent Ca^{2+} channel in A7r5 cells by endothelin-1 as revealed by a novel Ca^{2+} channel blocker LOE 908. *Br J Pharmacol* **126**:1107-1114.
- Iwasawa K, Nakajima T, Hazama H, Goto A, Shin WS, Toyooka T and Omata M (1997) Effects of extracellular pH on receptor-mediated Ca^{2+} influx in A7r5 rat smooth muscle cells: involvement of two different types of channel. *J Physiol* **503**:237-251.
- Jung S, Strotmann R, Schultz G and Plant TD (2002) TRPC6 is a candidate channel involved in receptor-stimulated cation currents in A7r5 smooth muscle cells. *Am J Physiol - Cell Physiology* **282**:C347-359.
- Krautwurst D, Degtiar VE, Schultz G and Hescheler J (1994) The isoquinoline derivative LOE 908 selectively blocks vasopressin-activated nonselective cation currents in A7r5 aortic smooth muscle cells. *Naunyn-Schmiedeberg's Arch Pharmacol* **349**:301-307.
- Kwan CY and Putney JW, Jr. (1990) Uptake and intracellular sequestration of divalent cations in resting and methacholine-stimulated mouse lacrimal acinar cells. Dissociation by Sr^{2+} and Ba^{2+} of agonist-stimulated divalent cation entry from the refilling of the agonist-sensitive intracellular pool. *J Biol Chem* **265**:678-684.
- McDaniel SS, Platoshyn O, Wang J, Yu Y, Sweeney M, Krick S, Rubin LJ and Yuan JX (2001) Capacitative Ca^{2+} entry in agonist-induced pulmonary vasoconstriction. *Am J Physiol - Lung Cellular & Molecular Physiology* **280**:L870-880.

- Moneer Z, Dyer JL and Taylor CW (2003) Nitric oxide co-ordinates the activities of the capacitative and non-capacitative Ca^{2+} -entry pathways regulated by vasopressin. *Biochem J* **370**:439-448.
- Moneer Z, Pino I, Taylor EJ, Broad LM, Liu Y, Tovey SC, Staali L and Taylor CW (2005) Different phospholipase C-coupled receptors differentially regulate capacitative and non-capacitative Ca^{2+} entry in A7r5 cells. *Biochem J* DOI: **10.1042/BJ20050145**.
- Moneer Z and Taylor CW (2002) Reciprocal regulation of capacitative and non-capacitative Ca^{2+} entry in A7r5 vascular smooth muscle cells: only the latter operates during receptor activation. *Biochem J* **362**:13-21.
- Montell C (2005) The TRP superfamily of cation channels. *Science's Stake [Electronic Resource]: Signal Transduction Knowledge Environment* **2005**:re3.
- Nakajima T, Hazama H, Hamada E, Wu SN, Igarashi K, Yamashita T, Seyama Y, Omata M and Kurachi Y (1996) Endothelin-1 and vasopressin activate Ca^{2+} -permeable non-selective cation channels in aortic smooth muscle cells: mechanism of receptor-mediated Ca^{2+} influx. *J Mol Cell Cardiol* **28**:707-722.
- Nakajima T, Kitazawa T, Hamada E, Hazama H, Omata M and Kurachi Y (1995) 17beta-Estradiol inhibits the voltage-dependent L-type Ca^{2+} currents in aortic smooth muscle cells. *Eur J Pharmacol* **294**:625-635.
- Ng LC and Gurney AM (2001) Store-operated channels mediate Ca^{2+} influx and contraction in rat pulmonary artery. *Circ Res* **89**:923-929.
- Parekh AB and Putney JW, Jr. (2005) Store-operated calcium channels. *Physiol Rev* **85**:757-810.
- Plane F, Johnson R, Kerr P, Wiehler W, Thorneloe K, Ishii K, Chen T and Cole W (2005) Heteromultimeric Kv1 channels contribute to myogenic control of arterial diameter. *Circ Res*. **96**:216-224.

- Schilling WP, Rajan L and Strobl-Jager E (1989) Characterization of the bradykinin-stimulated calcium influx pathway of cultured vascular endothelial cells. Saturability, selectivity, and kinetics. *J Biol Chem* **264**:12838-12848.
- Stevens T (2001) Is there a role for store-operated calcium entry in vasoconstriction? *Am J Physiol - Lung Cellular & Molecular Physiology* **280**:L866-869.
- Strübing C, Krapivinsky G, Krapivinsky L and Clapham DE (2003) Formation of novel TRPC channels by complex subunit interactions in embryonic brain. *J Biol Chem* **278**:39014-39019.
- Sweeney M, Yu Y, Platoshyn O, Zhang S, McDaniel SS and Yuan JX (2002) Inhibition of endogenous TRP1 decreases capacitative Ca^{2+} entry and attenuates pulmonary artery smooth muscle cell proliferation. *Am J Physiol - Lung Cellular & Molecular Physiology* **283**:L144-155.
- Trepakova ES, Gericke M, Hirakawa Y, Weisbrod RM, Cohen RA and Bolotina VM (2001) Properties of a native cation channel activated by Ca^{2+} store depletion in vascular smooth muscle cells. *J Biol Chem* **276**:7782-7790.
- Van Renterghem C, Romey G and Lazdunski M (1988) Vasopressin modulates the spontaneous electrical activity in aortic cells (line A7r5) by acting on three different types of ionic channels. *Proc Nat Acad Sci USA* **85**:9365-9369.
- Vanden Abeele F, Lemonnier L, Thebault S, Lepage G, Parys JB, Shuba Y, Skryma R and Prevarskaya N (2004) Two types of store-operated Ca^{2+} channels with different activation modes and molecular origin in LNCaP human prostate cancer epithelial cells. *J Biol Chem* **279**:30326-30337.
- Wang Y, Chen J, Wang Y, Taylor CW, Hirata Y, Hagiwara H, Mikoshiba K, Toyo-oka T, Omata M, Sakaki Y (2001) Crucial role of type 1, but not type 3, inositol 1,4,5-trisphosphate (IP_3) receptors in IP_3 -induced Ca^{2+} release, capacitative Ca^{2+} entry, and proliferation of A7r5 vascular smooth muscle cells. *Circ Res* **88**:202-209.

- Wu X, Zagranichnaya TK, Gurda GT, Eves EM and Villereal ML (2004) A TRPC1/TRPC3-mediated increase in store-operated calcium entry is required for differentiation of H19-7 hippocampal neuronal cells. *J Biol Chem* **279**:43392-43402.
- Xu SZ and Beech DJ (2001) TrpC1 is a membrane-spanning subunit of store-operated Ca^{2+} channels in native vascular smooth muscle cells. *Circ Res* **88**:84-87.
- Zagranichnaya TK, Wu X and Villereal ML (2005) Endogenous TRPC1, TRPC3, and TRPC7 Proteins Combine to Form Native Store-operated Channels in HEK-293 Cells. *J Biol Chem* **280**:29559-29569.
- Zakharov SI, Smani T, Dobryднеva Y, Monje F, Fichandler C, Blackmore PF and Bolotina VM (2004) Diethylstilbestrol is a potent inhibitor of store-operated channels and capacitative Ca^{2+} influx. *Mol Pharmacol* **66**:702-707.

FOOTNOTES

- *This work was supported by the John and Marian Falk Trust for Medical Research, and the National Heart, Lung, and Blood Institute (R01HL70670 (KLB)).
- ** To whom correspondence should be addressed.
- A preliminary account of this work has been presented and accepted for publication as an abstract:

Brueggemann LI, Markun DR, Cribbs LL, and Byron KL Electrophysiological and pharmacological characterization of store-operated currents and capacitative Ca^{2+} entry in rat vascular smooth muscle cells. *Journal of Physiology*, in press.

LEGENDS FOR FIGURES

Figure 1. I_{SOC} in A7r5 and Mesenteric Artery Smooth Muscle Cells. A. Representative time course of I_{SOC} development in an A7r5 cell (perforated patch, capacitance (C) = 95.5 pF) during application of 100 nM AVP (5 min, black box) followed by washout of AVP. Vasopressin was applied after stable recording of resting current for 5 min. Current was recorded with a ramp voltage protocol (from +85 mV to -115 mV) from -15 mV holding potential every 10 sec. Inward currents measured at -115 mV (open circles) and outward current measured at +85 mV (filled circles) are presented. Dotted line indicates 0 current level. B. Current-voltage relationship of AVP-activated I_{SOC} recorded at time point b on panel A after subtraction of resting current (time point a in panel A). Results are representative of 4 similar experiments. C. Representative time course of I_{SOC} development in an MASMC cell in perforated patch configuration (C = 25.4 pF) during application of 1 μ M thapsigargin (10 min, white box). D. Current-voltage relationship of thapsigargin-activated I_{SOC} recorded with the ramp voltage protocol (8 ramps recorded at the end of thapsigargin application (time point labeled b on panel C) were averaged and mean resting current (at time point a on panel C) was subtracted). Results are representative of 4 similar experiments.

Figure 2. Pharmacology of I_{SOC} . A. Reversibility of $GdCl_3$ block. Representative time course of I_{SOC} activation by 1 μ M thapsigargin (15 min) recorded in a single A7r5 cell (C=269.5 pF) with the perforated patch configuration as described in *METHODS* and measured at -115 mV. Drug applications are indicated by boxes. External solution contained 100 mM Na^+ and 20 mM Ca^{2+} . 100 μ M $GdCl_3$ (about 5 μ M free Gd^{3+}) was applied for 5 min followed by washout. Dotted lines indicate 0 current level. Similar results were obtained in 6 experiments. B. 2-APB reversibly inhibited I_{SOC} . Application of 10 μ M 2-APB for 10 min reversibly inhibited I_{SOC} activated with 1 μ M thapsigargin (20 min) in perforated patch configuration (C=467.5 pF, representative of 4 similar experiments), followed

by washout and reversible inhibition of current by 100 μM GdCl_3 . C. Reversible inhibition of I_{SOC} by DES. Thapsigargin-activated I_{SOC} (1 μM thapsigargin for 15 min in perforated patch recording, $C=177.2$ pF, representative of 4 similar experiments) was reversibly blocked by external application of 10 μM DES for 10 min. D. Inhibition of I_{SOC} by flufenamic acid. Representative time course of I_{SOC} activation by dialyzing a cell ($C=195$ pF) with 10 mM BAPTA in ruptured patch configuration (external solution contained 100 mM NMDG^+ and 20 mM Ca^{2+}) followed by application of 100 μM flufenamate. Similar results were obtained in 4 experiments with BAPTA dialysis and in 3 experiments when thapsigargin was used to activate I_{SOC} . E. OAG did not activate I_{SOC} . No measurable inward current developed at -115 mV in perforated patch recording upon application of 100 μM OAG for 10 min in external solution contained 100 mM NMDG^+ and 20 mM Ca^{2+} ($C=186$ pF, representative of 4 similar experiments).

Figure 3. Divalent selectivity of I_{SOC} . A, B. Representative time courses of I_{SOC} development at -115 mV (ruptured patch configuration) recorded in 20 mM CaCl_2 followed by replacement of CaCl_2 with an equimolar concentration of SrCl_2 (A, $C = 102$ pF) or MnCl_2 (B, $C = 114$ pF). C. Relative current densities of Gd^{3+} -sensitive I_{SOC} measured at -115 mV in 20 mM CaCl_2 , 20 mM SrCl_2 ($n=6$), 20 mM BaCl_2 ($n=5$) and 20 mM MnCl_2 ($n=4$) normalized by current densities in CaCl_2 as described in *METHODS*. Data are presented as mean \pm S.E. * $p<0.05$ versus current densities in CaCl_2 .

Figure 4. Inhibition of CCE by 2-APB. A. Reversible inhibition of CCE by 2-APB and Gd^{3+} . Fura-2-loaded A7r5 cells were pretreated with 1 μM thapsigargin and exposed to the same external solution used for recording I_{SOC} (see *METHODS*) until the fluorescence ratio stabilized (approximately 40 min). 2-APB (10 μM) was added as indicated by the white box, followed by washout of 2-APB and application of Gd^{3+} (black box; 100 μM total Gd^{3+} to give a free concentration of approximately 5

μM). Results are representative of 13 similar experiments. B. The concentration-dependence for 2-APB inhibition of CCE was evaluated in A7r5 cells exposed to thapsigargin ($1\ \mu\text{M}$) in Ca^{2+} -free medium followed by readdition of extracellular Ca^{2+} (final concentration, $1.5\ \text{mM}$). Traces are representative of 4 experiments performed in triplicate and demonstrate control responses (in the absence of 2-APB) and responses recorded with $10\ \mu\text{M}$ 2-APB in all solutions. C. Concentration-dependence for 2-APB inhibition of Ca^{2+} entry. Ca^{2+} entry was estimated by calculating the maximal change in fura-2 fluorescence ratio on addition of extracellular Ca^{2+} . Results from 4 experiments, presented as mean \pm S.E., are normalized to the control responses in the absence of 2-APB recorded from cells on the same 96-well plate. * $p < 0.05$ versus control.

Figure 5. **Inhibition of CCE by Gd^{3+} .** A. Concentration-dependence for inhibition of CCE by Gd^{3+} was evaluated as for 2-APB (see legend for Fig. 4). Data from 4 experiments performed in triplicate are presented as mean \pm S.E. * $p < 0.05$ versus control. B. In chloride-based external solution (control medium—see *METHODS*), the steady-state Ca^{2+} plateau in thapsigargin-pretreated cells was reversibly inhibited by 2-APB ($10\ \mu\text{M}$, white box), whereas the inhibition by Gd^{3+} ($2.5\ \mu\text{M}$, black box) was poorly reversed on washout of Gd^{3+} . Results are representative of 4 similar experiments.

Figure 6. **Inhibition of CCE by DES.** A. Effects of DES on CCE were evaluated in fura-2-loaded A7r5 cells pretreated with thapsigargin in external solution used for recording I_{SOC} . A representative recording is shown demonstrating sequential inhibition of CCE by Gd^{3+} ($5\ \mu\text{M}$, black box), 2-APB ($10\ \mu\text{M}$, white box), and DES ($10\ \mu\text{M}$, gray box). Similar results were obtained in 5 experiments. B. Concentration-dependence for inhibition of thapsigargin-stimulated CCE by DES was determined as for 2-APB and Gd^{3+} (see legend for Fig. 4). Data from 3 experiments performed in triplicate are presented as mean \pm S.E. * $p < 0.05$ versus control.

Figure 7. AVP-induced Ca^{2+} entry: inhibition by 2-APB, Gd^{3+} , and DES. A7r5 cells were loaded with fura-2 in control medium (see *METHODS*). At the start of recording, the cells were exposed to the external solution used for recording I_{SOC} . All subsequent recordings were in this solution. A. After $[\text{Ca}^{2+}]_{\text{i}}$ stabilized in the external solution, the cells were exposed to 1 nM AVP (light gray box). 2-APB (10 μM , white box), Gd^{3+} (5 μM , black box), and DES (10 μM , dark gray box) were applied sequentially. Responses are representative of 6 similar experiments. Note that after inhibition of the Ca^{2+} response by DES, the Ca^{2+} signal begins to rebound in the continued presence of DES. B. Application of DES in the presence of low $[\text{Gd}^{3+}]$ (5 μM) does not prevent the rebound of $[\text{Ca}^{2+}]_{\text{i}}$. Trace is representative of 3 similar experiments. C. A similar rebound of $[\text{Ca}^{2+}]_{\text{i}}$ is observed on application of DES in the presence of high $[\text{Gd}^{3+}]$ (100 μM) (representative of 2 similar experiments).

Figure 8. TRPC homologue expression in A7r5 and MASMC. A. total RNA prepared from A7r5 cells (upper panel), freshly isolated MASMC (middle panel), and adult rat brain (lower panel) was reverse transcribed and subjected to PCR using primers specific for TRPC1 through TRPC7 (lanes 1-7). Molecular weight marker (M) is 100-base pair ladder, and 800 bp is indicated by the arrow head on the left. B. Western blot analysis of TRPC1, TRPC4, and TRPC6 protein expression in A7r5 cells and cultured MASMC.

Figure 9. Knock-down of TRPC1 expression and I_{SOC} in A7r5. A. Level of TRPC1 mRNA measured with real-time PCR as described in *METHODS* was significantly decreased by $42 \pm 4\%$ (white bar, $n=3$, $*p<0.05$ versus control, paired Student's *t*-test) in A7r5 cells stably expressing TRPC1 siRNA, compared with control cells maintained in parallel cultures (black bar). B. Representative Western blot of TRPC expression in control or TRPC1 siRNA stable cell line. A single blot was probed simultaneously with antibodies for TRPC1, TRPC4, and TRPC6. On average the density of the TRPC1 band was reduced by $56 \pm 6\%$ in the TRPC1 siRNA stable cell line compared with controls

($p < 0.05$, paired Student's t -test, $n=3$). The TRPC4 and TRPC6 bands were not significantly different.

C. Current densities of Gd^{3+} -sensitive I_{SOC} measured at -115 mV with ruptured patch configuration in control untransfected cells (-0.49 ± 0.06 pA/pF, $n=8$, black bar) and in A7r5 cells stably transfected with TRPC1 siRNA (-0.22 ± 0.02 pA/pF, $n=8$, white bar). * $p < 0.001$ versus control. D.

Immunostaining for TRPC1 (left panels) or GFP fluorescence (right panels) in A7r5 cells infected with AS-TRPC1-GFP adenovirus (100 MOI, 5 days). Two different anti-TRPC1 antibodies were used (I, Villereal antibody; III, Santa Cruz antibody) to demonstrate that TRPC1 immunoreactivity was reduced in GFP-positive cells (II & IV, cells outlined in red) compared with non-GFP-expressing cells in the same field. E. Representative time courses of I_{SOC} development in A7r5 cells treated with AS-TRPC1-GFP adenovirus (30 MOI, 4-6 days). Currents were measured at -115 mV with ruptured patch configuration in non-fluorescent cells (filled circles), and fluorescent cells expressing TRPC1 antisense (open circles). F. Summary of current densities of Gd^{3+} -sensitive I_{SOC} in control (non-fluorescent) cells (-0.55 ± 0.04 pA/pF, $n=16$, black bar) and fluorescent cells expressing TRPC1 antisense (-0.31 ± 0.04 pA/pF, $n=15$, white bar). * $p < 0.001$ versus control.




Figure 1




Figure 2




Figure 3





Figure 4A,B




Figure 4C





Figure 5




Figure 6




Figure 7





Figure 8




Figure 9