

**Phenobarbital treatment inhibits the formation of estradiol-dependent mammary tumors
in the ACI rat**

Sonia Mesia-Vela, Rosa I. Sanchez, Kenneth R. Reuhl, Allan H. Conney, Frederick C. Kauffman

Laboratory for Cellular and Biochemical Toxicology (S.M-V., R.I.S. F.C.K.), Laboratory for
Neurotoxicology (K.R.R.), Department of Pharmacology and Toxicology, and Susan Lehman
Cullman, Laboratory for Cancer Research , Department of Chemical Biology (A.H.C.), Ernest
Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

Running title: Inhibition of adenocarcinomas by Phenobarbital in ACI rats

Corresponding author: Dr. Frederick C. Kauffman
Rutgers University
41 Gordon Rd
Piscataway, NJ 08854
Phone: (732) 445 6900
Fax: (732) 445 6905
Email: kauffma@rci.rutgers.edu

Number of text pages: 20

Number of tables: 6

Number of figures: 3

Number of references: 40

Number of words in Abstract: 238

Number of words in Introduction: 522

Number of words in Discussion: 1348

List of Non-Standard abbreviations:

PB, phenobarbital; NQO1, NADPH: quinone oxidoreductase; GST, glutathione S-transferase

Recommended Section Assignment: Absorption, Distribution, Metabolism and Excretion

Abstract

Exposure of female ACI rats for 28 weeks to 3 mg estradiol (E_2) contained in cholesterol pellets elevated blood E_2 levels and caused palpable mammary tumors in all animals. Co-administration of phenobarbital (PB) in their drinking water reduced the incidence, number and size of MT but did not reduce blood E_2 levels. Inhibition of MT by PB was accompanied by significant changes in total hepatic metabolism of E_2 measured *in vitro*. PB treatment caused approximately a 4-fold increase in hepatic metabolism of E_2 in control and E_2 -treated rats. The major NADPH-dependent metabolites of E_2 were 2-OH E_2 and estrone (E_1). PB, either alone or together with E_2 , increased microsomal 2-hydroxylation of E_2 ; formation of E_1 was either unaffected or decreased slightly. PB also increased microsomal metabolism of E_2 to minor metabolites (4-OH- E_2 , 6 α -OH- E_2 , 6 β -OH- E_2 , 14 α -OH- E_2 , 6-keto E_1 and 2-OH- E_1) and reduced the formation of the E_2 -17 β -oleoyl ester and the E_2 -3- and 17-glucuronides. In contrast, when given in combination with E_2 , PB increased the formation of both glucuronides. Co-treatment of animals with PB and E_2 increased activities of NADP(H):quinone oxidoreductase and glutathione-S-transferase to a greater extent than either compound alone. Collectively, these results show that the multiple actions of PB on hepatic metabolism of E_2 including induction of E_2 -hydroxylation, glucuronidation, and antioxidant defense enzymes along with inhibition of E_2 esterification in livers of female ACI rats accompany a marked reduction of E_2 -dependent mammary tumors in this model.

Introduction

Phenobarbital (PB) is a known inducer of microsomal hydroxylation of drugs and steroids in rodents and in humans (Conney et al., 1973). Treatment of rodents with PB induces the synthesis of hepatic microsomal enzymes that hydroxylate progesterone, estradiol (E₂), estrone (E₁), deoxycorticosterone, testosterone, Δ^4 -androstene-3,17-dione and cortisol (reviewed by Zhu and Conney, 1998). Endogenous estrogens are hydroxylated at multiple positions by several hepatic and non-hepatic microsomal monooxygenase systems (reviewed by Martucci and Fishman, 1993; Zhu and Conney, 1998). The alteration of microsomal hydroxylation of steroids by PB is reflected *in vivo* by enhanced metabolism and altered actions of steroids. For example, pretreatment of rats with PB decreases the uterotrophic action of E₂ and E₁ and enhances their metabolism *in vivo* (Levin et al., 1967, Levin et al., 1968). Treatment of rats with PB also inhibits the growth-promoting effect of testosterone on the seminal vesicles (Levin et al., 1974) and decreases the anesthetic action of progesterone and deoxycorticosterone in rodents (Conney et al., 1965).

Oxidative stress arising from redox cycling of catechol estrogens formed during E₂ metabolism has been suggested as an important factor in initiation and progression of many cancers including mammary carcinogenesis (Cavalieri et al., 1997, 2004). In mammals, the liver contains high levels of cytochromes P450 (CYP) which catalyze NADPH-dependent oxidation of estrogens to various hydroxylated or keto metabolites (Martucci and Fishman, 1993; Zhu and Conney, 1998). Reaction of endogenous catechol estrogens with DNA causes formation of depurinating DNA adducts (Cavalieri et al., 1997; Li et al., 2004; Cavalieri et al., 2004) which has been proposed to cause oncogenic mutations (Chakravarti et al., 1995). The carcinogenic

effects of 4-catechol estrogens (E_2 and E_1) in the kidney of castrated male Syrian hamsters (Liehr et al., 1986) reinforce this hypothesis. However, administration of high doses of 2-OH- E_2 , 4-OH- E_2 or 4-OH- E_1 to ACI rats failed to cause mammary tumors under conditions where E_2 was highly active (Turan et al., 2004). In addition, the direct injection of estrone-3,4-quinone (the chemically reactive ortho-quinone derived from 4-OH- E_1) under the nipples of the mammary glands in rats failed to cause mammary tumors, whereas injection of a positive control, a diol epoxide of benzo[c] phenanthrene was highly active (El Bayoumy et al., 1996).

In the present study we investigated the effects of PB on the formation of mammary tumors induced by E_2 in ACI rats, an estrogen-sensitive strain which is considered a unique model because of its high sensitivity to estrogen dependent mammary ductal adenocarcinomas (80 to 100% incidence) within a relatively short time period (Harvell et al., 2000; Shull et al., 1997; Li et al., 2002). The main objective of the present study was to determine whether PB-induced increases in E_2 -metabolism in the liver corresponded with alterations in the formation of E_2 -induced mammary tumors. The liver is a major site of estrogen metabolism, and estrogen circulates primarily as inactive conjugates of E_1 and E_2 produced in the liver, i.e., conjugated E_2 is a transport form of estrogen which is converted back to active estrogen in target tissues such as mammary gland (reviewed in: Pasqualini 2004; Reed et al., 2005; Zhu and Conney, 1998)

Methods

Chemicals. E₂, ascorbic acid, NADPH, β-glucuronidase (EC. 3.2.1.3.1), UDP-glucuronic acid (UDPGA), sodium phenobarbital, saccharic acid 1,4-lactone, *p*-nitrophenol, 1-chloro-2,4-dinitrobenzene (CDNB), reduced glutathione, cytochrome c, oleoyl coenzyme A, sodium azide, and Tris-base were obtained from Sigma Chemicals Co. (St. Louis, MO). Glutathione reductase (140.7 U/mg protein) was purchased from Fluka Biochemica (Switzerland), ³⁵S-PAPS from New England Nuclear (Boston, MA), and PAPS (>99% pure) from H. Glatt and R. Landseidel, German Institute of Nutrition (Potsdam, Germany). [2,4,6,7,16,17-³H(N)]-Estradiol (*s.a.* 110-170 Ci/mmol) was purchased from NEN Life Science Products Inc. (Boston, MA). Pellets with or without E₂ were purchased from Hormone Pellets Press (Shawnee Mission, KS). AIN-76A diet was purchased from Dyets Inc. (Bethlehem, PA). All other chemicals used were of the highest grade from standard sources.

Animals and treatments. Female ACI rats (7 to 8 weeks old) were obtained from Harlan Sprague-Dawley Laboratory (Indianapolis, IN). The animals were housed individually in an AAALAC accredited barrier facility under controlled temperature, humidity, and lighting conditions and were fed with AIN-76A diet (Dyets Inc., Pennsylvania, PA). Treatment protocols started 4 days after arrival of the animals. Rats received water *ad libitum* or 0.05% PB in their drinking water. A single 20-mg pellet containing 3 mg of E₂ plus 17 mg of cholesterol was implanted subpannicularly in the shoulder region, as previously described (Li et al., 2002). Control animals were implanted with 20-mg cholesterol pellets alone. The rats were palpated for mammary tumors twice weekly, and weighed every two weeks for the duration of the experiment. Animals (N=8-15) were killed by decapitation after 6, 12 or 28 weeks. The

geometric volume of the tumors was determined using the formula: length x width x height x 0.5326, assuming an hemi-ellipsoid shape (Shah et al., 1999).

Serum levels of E₂

Trunk blood collected at decapitation was allowed to clot at 4°C for six hours and centrifuged. The serum was collected and stored at -80°C. Circulating levels of E₂ were determined in whole serum by RIA, using Coat-A Count® Estradiol RIA kits (Diagnostic Products Corporation, Los Angeles, CA). According to the manufacture's instructions, this assay measures both bound and free E₂ in serum.

Tissue Processing. All the animals were subjected to macroscopic pathologic examination when killed and the number, volume and localization of mammary tumors recorded. The mammary glands and the tumors were quickly removed. Portions of these tissues were fixed in Carnoy's solution for 4 h and processed for embedding in paraffin. Sections (6 µm) were prepared from each of the Carnoy's-fixed tissues and stained with hematoxylin and eosin. Selected estrogen target organs including pituitary, adrenals, thymus, uterus, kidneys and liver were removed and weighed.

Preparation of hepatic subcellular fractions. Liver cytosols and microsomes were prepared by differential centrifugation as described previously (Thomas et al., 1983) and stored at -80°C until used. The protein concentration was determined with the BCA™ protein assay kit (PIERCE, Rockford, IL) according to the supplier's instructions using bovine serum albumin as a standard.

Enzyme Assays

NADPH- dependent oxidation of E₂ (CYP450 assay) was carried out using liver microsomes

incubated with 5 mM ascorbic acid, 3 mM magnesium chloride, 50 μ M sodium phosphate buffer (pH 7.4) and 25 μ M [3 H]-E₂ (0.5 μ Ci) for 20 min at 37 °C. The enzyme reaction was initiated with 2 mM NADPH and terminated by the addition of 5 ml of ethyl acetate and vortexing. The ethyl acetate extracts were evaporated to dryness under nitrogen. The residue was dissolved in methanol and analyzed for metabolite composition by HPLC as described previously (Suchar et al., 1996; Mesia-Vela et al., 2002).

Fatty acyl-CoA:estradiol acyltransferase was assayed in reaction mixtures containing 50 μ M [3 H]-E₂ (1 μ Ci), 100 μ M fatty acyl-CoA, 5 mM magnesium chloride in 0.1 M sodium acetate buffer (pH 5.5) in a final volume of 0.5 ml. The reaction was initiated by the addition of liver microsomes (1 mg of protein/ml). After incubation at 37°C for 30 min, the reaction was arrested by placing the tubes on ice, followed by addition of 0.2 ml of ice-cold sodium acetate buffer (pH 5.5) and extraction by vortexing with 4 ml of ethyl acetate (HPLC grade from Fisher Scientific). Dry extracts were redissolved and 90 μ l aliquots analyzed by HPLC as described previously (Xu et al., 2002). Metabolite quantification was based on the amount of radioactivity in the metabolite peak as compared to the total radioactivity collected from the HPLC column from each sample.

Glucuronosyltransferase activity was assayed using a modification of a previously described method (Sanchez et al., 2003). The reaction mixture contained 1.0 mg of microsomal protein, 2 mM UDPGA, 5mM MgCl₂, 100 μ M [3 H]-E₂ (0.15 μ Ci) and 50 mM Tris-HCl buffer pH 8.5 in a final volume of 150 μ l. The reaction was initiated by addition of UDPGA. Incubations proceeded at 37°C for 15 min and were terminated by placing them on ice and adding 50- μ l ice-cold acetonitrile. The reaction mixtures were then vortexed and centrifuged at 3000 x g for 5

min. 10 μ l of the supernatants were used for the determination of E₂ and E₂-glucuronides by HPLC. The HPLC system consisted of a Shimadzu SCL-10A system controller with a Shimadzu SIL-10A auto injector, two LC-10AD pumps, SPD-10A UV-Vis detector set at 280 nm and an Eclipse XDB C18 (4.6mm \times 150 mm) column (MacMod Analytical, Chadds Ford, PA). The solvent system consisted of solvent A: 0.1% acetic acid in water and solvent B: containing 20% methanol, 80% acetonitrile and 0.1% acetic acid. E₂ and its 3- and 17-hydroxyglucuronides were eluted with a 30 min linear gradient from 25 to 90% B. Metabolite quantification was based on the amount of radioactivity in the metabolite peak as compared to the total radioactivity collected from the HPLC column from each sample. The retention time of metabolites and E₂ agreed with corresponding UV-absorbing peaks of standards. The glucuronides were identified by their co-elution with authentic standards or by determination of E₂ after hydrolysis in the presence of β -glucuronidase (200 U/ml).

Cytosolic sulfotransferase (SULT) activity was determined using PAP³⁵S as cofactor. The incubations were carried out with 20 mM Tris.HCl, pH 7.5, 4 mM MgCl₂, 10 μ M PAP³⁵S (0.02 μ Ci) and 5 μ M *p*-nitrophenol in 100 μ l (Foldes and Meek, 1973).

Cytosolic NADPH quinone oxidoreductase (NQO1) was measured by reduction of cytochrome c (50 μ M) in the presence of liver cytosol, 10 μ M menadione and 1 mM NADPH. The reaction was monitored at 550 nm. The reactions were carried out in 100 mM potassium phosphate buffer pH 7.7 containing 0.04% triton X-100 at 25°C (Jaiswal et al., 1988). Activity of dicumarol-inhibitable menadione reductase was determined using an extinction coefficient of 21/mM/cm for cytochrome c.

Cytosolic glutathione S-transferase (GST) activity was measured in the presence of liver cytosolic protein, 1 mM 1-chloro-2,4-dinitrobenzene (CDNB), 1 mM reduced glutathione and 100 mM potassium phosphate buffer pH 6.5, at 25°C. Conjugation of CDNB with glutathione was monitored at 340 nm. Specific activity was calculated using an extinction coefficient of 9.6/mM/cm (Habig et al., 1974).

Cytosolic glutathione peroxidase (GPx) activity was measured using hydrogen peroxide as the substrate and was monitored by the decrease in absorbance at 340 nm. Specific activity was calculated using an extinction coefficient for NADPH of 6.22/mM/cm (Flohe and Gunzler, 1984).

Statistical analysis of data

Data are presented as the mean \pm S.E. Differences between means were assessed by ANOVA followed by Bonferroni hoc post test, $P < 0.05$.

Results

Tissue wet weights and histopathology in PB and E₂ –treated ACI rats. Administration of E₂ increased pituitary weight by 3.5 and 4.8-fold after 6 and 12 weeks of treatment, respectively. These E₂-dependent increases in pituitary weights were not reduced by co-administration of PB (Table 1). A small increase in kidney weight (36%) and a reduction of thymus weight was also observed in the group treated with E₂ for 6-12 weeks (Table 1). Relative liver weight was increased about 1.5-fold by E₂ at both experimental periods (Table 2). Administration of PB also enhanced the E₂-induced increase in the relative liver weight, indicating differential effects on liver by both drugs. A 2 to 2.6-fold increase in the microsomal protein/liver after 6 and 12 weeks of PB treatment reflected the induction of microsomal protein by PB treatment (Table 2). PB alone or in combination with E₂ slightly increased adrenal weights at 6 weeks (Table 1). No alteration of body or uterine weight was seen during PB or E₂ treatment. It is noteworthy that PB did not alter E₂- dependent increases in pituitary weights (Table 1). Significant losses in body weight were noted at about 20 weeks when large palpable mammary tumors were seen in E₂-treated rats. In addition, pathology studies showed key changes in liver and mammary tissue. In liver, there was an increase of mitotic figures and scattered single cell degeneration caused by E₂ treatment alone. These effects diminished with time of exposure. Administration of PB alone induced enlargement of hepatocytes and vacuolar degeneration that increased markedly with the time of exposure to the drug. In contrast, for co-administered PB and E₂, the effects of PB were predominant with a reduction of the scattered single cell degeneration induced by E₂. In mammary samples, pronounced hyperplasia of mammary ductal cells due to E₂- treatment was observed. No quantitative difference could be observed in the extent of hyperplasia induced by

E_2 in relation to time of exposure, but some of the samples from the 12 week exposure group showed major ductal changes and atypia. PB treatment alone did not alter the morphology of mammary tissue. Mammary tumors classified as mammary ductal adenocarcinomas (MDAs) were histologically similar in animals treated with E_2 alone or E_2 plus PB.

Effect of PB on the incidence, multiplicity and size of mammary tumors in E_2 -treated ACI

rats. The first mammary tumor appeared during the 15th week in E_2 - treated rats. Fifty percent of the animals had mammary tumors by the 26th week and 100% of the animals had mammary tumors by week 28. Although PB administration did not substantially alter the initial onset of E_2 -induced mammary tumors, it reduced to 53% the number of rats with mammary tumors at the end of the experiment (28 weeks) (Fig. 1). Treatment of the rats with PB had a dramatic inhibitory effect on the number and size of E_2 -induced mammary tumors observed at 28 weeks (96-97 % inhibition; Fig. 2).

Serum E_2 levels. Treatment of female ACI rats with E_2 increased normal serum E_2 levels by 9 and 6-fold after 6 and 12 weeks of treatment, respectively. PB alone did not alter the serum level of total E_2 in normal or E_2 -treated rats at any of the periods of the study (Fig 3).

Effect of PB administration on the NADPH-dependent oxidative metabolism of E_2 by liver

microsomes. 2-OH- E_2 and E_1 were the major metabolites formed during the incubation of E_2 with control liver microsomes and NADPH (Table 3). Treatment of the rats with PB for 6 and 12 weeks increased the formation of 2-hydroxy- E_2 per liver by 4.7 and 6.6 fold, respectively when compared with E_2 -treated rats. Treatment of the rats with PB for 6 and 12 weeks prevented the increase in formation E_1 induced by chronic E_2 treatment. In addition to the above results, PB

administration also caused substantial increases in the metabolism of E₂ to 6 α -hydroxy-E₂, 6 β -hydroxy-E₂, 14 α -hydroxy-E₂, 4-hydroxy-E₂, 6-keto-E₂ and 2-hydroxy-E₁ per liver (calculated from tables 2 and 3). Since the later compounds are relative minor metabolites, the overall major effect of PB was to enhance the 2-hydroxylation of E₂ and also the 2-hydroxylation of E₁.

Effect of PB administration on the esterification of E₂ by fatty acylCoA: E₂-acyltransferase in liver microsomes. Treatment of rats with E₂ for 6 or 12 weeks had little or no effect on liver microsomal formation of E₂-17 β -oleoyl ester when compared with control rats (calculated from Tables 2 and 4); however, co-administration of PB and E₂ reduced the esterification of E₂ by fatty acylCoA:E₂-acyltransferase significantly at both 6 and 12 weeks (Table 4). PB treatment alone decreased esterification of E₂ only after 12 weeks of treatment.

Effect of PB administration on the glucuronidation of E₂ by liver microsomes.

Treatment of rats with E₂ for 6 or 12 weeks did not alter microsomal glucuronidation of E₂ per mg microsomal protein at any experimental time; however, treatment with PB reduced the formation of both E₂-glucuronides (3- and 17-) by 45- and 34%, respectively, at 12 weeks. When given in combination with E₂, PB increased formation of both glucuronides by 1.2- to 1.5-fold per mg protein after 6 and 12 weeks, respectively (calculated from Tables 2 and 4).

Effect of E₂ and PB administration on sulfotransferase and antioxidant enzymes in the liver.

Hepatic sulfotransferase: E₂ administration for 6 or 12 weeks decreased p-nitrophenol sulfonation per mg cytosolic protein by ~ 30% (Table 5). Administration of PB had little or no effect on p-nitrophenol sulfonation (SULT1A1) activity when given alone or together with E₂

(Table 5).

JPET #96867

14

Antioxidant enzymes: Although administration of PB caused a modest decrease in hepatic GPx activity per mg cytosolic protein and E₂ treatment was inactive, the administration of E₂ or PB for 6 or 12 weeks caused large increases in hepatic NQO1 and GST activity (Table 6).

Administration of PB for 6 or 12 weeks caused a 3.7- to 4.2- fold increase in NQO1 activity per mg cytosolic protein, and administration of E₂ alone increased this activity 2.6- to 3.7-fold (Table 6). Administration of PB for 6 and 12 weeks elevated GST activity 2.7- to 2.9-fold per mg cytosolic protein; and administration of E₂ alone increased this activity 1.9- to 2-fold (Table 6).

The administration of a combination of PB and E₂ increased in NQO1 activity 68 to 84% (4.9- to 6.3-fold) when compared with the activity observed after administration of E₂ alone.

Administration of a combination of PB and E₂ also increased GST activity beyond that seen with either compound alone (3.1- to 3.9-fold per mg cytosolic protein, Table 6). In summary, administration of E₂ or PB alone caused large increases in the levels of the antioxidant enzymes, NQO1 and GST per mg cytosolic protein, and the effects of the combined administration of PB and E₂ were even larger than after administration of E₂ or PB alone.

Discussion

The frequency, size and number of mammary tumors induced in female ACI rats chronically treated with E₂ were reduced dramatically by PB given in the drinking water (Figs 1 and 2). The reduction of E₂-dependent mammary tumors noted in animals exposed to PB indicates that very potent mechanisms of protection are activated by this compound in ACI rats. Surprisingly, the dramatic decrease in mammary tumors in PB treated rats did not correlate with levels of serum E₂ measured in animals given E₂ chronically (Fig. 3). The failure of pituitary and uterine weights, which normally change in response to the “estrogenicity” of serum *in vivo*, to change in rats co-treated with E₂ and PB is in accord with the finding that total E₂ levels were essentially the same in serum from both groups of animals. Further studies to characterize bound and free forms of E₂ and its metabolites in serum and mammary tissue of rats treated with E₂ alone or in combination with PB over the course of mammary tumor generation are needed in future studies employing the ACI rat model.

The inhibition of E₂-dependent tumorigenesis by PB was accompanied by significant changes in both the oxidation of E₂ and by smaller changes in conjugation reactions. In addition, PB treatment had a remarkable stimulatory effect on two antioxidant defense enzymes, NQO1 and GST, which may be particularly important for the protection of PB against mammary tumorigenesis in the ACI rat model. Since blood E₂ levels were unchanged after PB, it is possible that E₂ synthesis was enhanced to compensate for increased E₂ metabolism. Further studies to clarify the relationship between alterations in hepatic metabolism of E₂ and the profile of E₂ metabolites in blood over the course of mammary tumor formation in the ACI rat model are

clearly warranted.

Chemoprevention of E₂-induced mammary tumors and modulation of microsomal oxidation of E₂ by PB.

The chemopreventive effect of PB likely involves multiple mechanisms including an increase in microsomal oxidation triggered by PB. The strong stimulatory effect of PB on the hepatic formation of E₂-hydroxylated E₂-metabolites including some that are formed only in trace amounts if at all in control animals (4-hydroxy-E₂, 6 α -hydroxy-E₂, 14 α -hydroxy-E₂, 6 β -hydroxy-E₂ and 6-keto-E₁) is noteworthy. The formation and pattern of induction of hydroxylated metabolites of E₂ by PB was similar to findings in other rat strains (Suchar et al., 1996) which differ from the ACI rat in sensitivity to E₂-induced mammary tumors. The major hydroxylated metabolite produced in the liver of the ACI rat is 2-hydroxy-E₂. This metabolite, which is potentially chemopreventive against mammary tumors, was markedly induced in liver microsomes by PB in rats also treated with E₂. A number of studies indicated that the 2-methoxy derivative of this metabolite formed by catechol O-methyl transferase has strong antiproliferative and proapoptotic actions in a variety of human cancer cell lines including human breast cancer lines *in vitro* (Lottering et al., 1992; Pribluda et al., 2000; Liu and Zhu, 2004). Furthermore, 2-methoxy-E₂ has strong antiangiogenic and inhibitory effects on growth of mammary tumors in mice *in vivo* (Klauber et al., 1997; Fotsis et al., 1994). Recently, synergistic inhibitory effects of 2-methoxy-E₂ and microtubule-disrupting agents were observed on the proliferation of human breast cancer cells (Han et al., 2005). Thus, the marked stimulatory effect of PB on the 2-hydroxylation of E₂ may contribute to the inhibitory effect of PB on E₂ induced mammary tumors in the ACI rat. The other hydroxylated metabolites, some of which (e.g. 4-hydroxy-E₂)

may be tumorigenic (Liehr 1997), were also increased to a variable extent by PB. Transport of some of these metabolites either in their free or conjugated form to mammary tissue may contribute to mammary tumorigenesis either by direct genomic effects or via oxidative stress in mammary tissue. It is unlikely that circulating 4-hydroxy-E₂, 4-hydroxy-E₁ or 16 α -hydroxy-E₂ contribute to the carcinogenic effects of E₂ in the ACI rat since these metabolites of E₂ were not tumorigenic under conditions described above where E₂ caused mammary tumors (Turan et al., 2004). In addition, the direct injection of estrone-3,4-quinone (the chemically reactive ortho-quinone derived from 4-OH-E₂) under the nipples of the mammary gland in CD rats failed to cause mammary tumors (El-Bayoumy et al., 1996). It is of interest that injection of 4-hydroxy-E₂ or E₂-3,4-quinone into the mammary gland of the female ACI rat resulted in 4-hydroxy-E₂-1-N₃-adenosine and 4-hydroxy-E₂-1-N₇-guanine adducts in DNA (Li et al., 2004). It will be of interest to determine whether these injections result in mammary cancer.

Chemoprevention of E₂-induced mammary tumors and modulation of E₂ conjugation by PB

E₂-3- and 17- glucuronidation, major inactivation pathways of E₂, were modestly increased when PB was co-administered with E₂. In contrast, glucuronidation of E₂ was reduced by PB-treatment alone at 12 weeks. Reduction in the capacity of liver to form E₂-glucuronides following chronic exposure to PB may increase amounts of active E₂ available for transport to E₂-target tissues such as mammary tissue. However, when PB was given concomitantly with E₂, the formation of E₂-3- and 17-glucuronide was modestly increased suggesting enhanced inactivation of E₂ in these animals. Reduction of microsomal fatty acylCoA: E₂ acyltransferase activity in PB-treated rats would reduce the formation of fatty acyl esters of E₂, which are

postulated to be long term storage forms of E₂ in fatty tissues such as the breast (Xu et al., 2002). Further, hepatic cytosolic sulfotransferase (SULT1A1), that may be involved in the conversion of E₂ to conjugates that serve as transport and storage forms, was also decreased slightly by treatment with combinations of PB and E₂.

Chemoprevention of E₂-induced mammary tumors and modulation of antioxidant pathways by PB and E₂.

The observation that administration of PB alone or together with E₂ strongly induced the activity of NQO1 and GST in liver after 6 and 12 weeks of treatment is intriguing because these activities may be linked to detoxification of reactive oxygen species generated by redox cycling of E₂-catechols; and increased levels of these antioxidant enzymes have been associated with anticarcinogenic effects (Ramos-Gomez et al., 2001). It is very important to determine whether the actions of PB observed in liver also occur in mammary tissue. The effects of PB on NQO1 and GST in liver were opposite to those on GPx and were in accord with the idea that regulation of expression of GPx differs from that of NQO1 and GST (Radjendirane et al., 1998; Esposito et al., 2000). It is also of considerable interest that administration of E₂ alone increases the level of antioxidant enzymes, a finding observed earlier by our group (Sanchez et al., 2003; Mesia-Vela et al., 2004). Moreover, the induction of NQO1 and GST in response to chronic treatment with E₂ seems to be specific for the ACI rat (Sanchez et al., 2003). The stimulatory effect of PB administration alone or together with E₂ on NQO1 and GST activity suggests that PB induction of antioxidant enzymes may play a role in the protective effect of PB on E₂-induced breast cancer.

In conclusion, the growth and multiplicity of E₂-dependent mammary tumors in the ACI

rat is markedly reduced by PB in the drinking water. This reduction of mammary tumors in the ACI rat is accompanied by multiple effects of PB on hepatic metabolism including: induction of E₂-hydroxylation via NADPH-dependent oxidations (predominantly an increase in 2-hydroxy E₂ formation), alteration in the formation of E₂ conjugates, and by the induction of important antioxidant defense activities. Such changes induced by PB may reduce the incidence of mammary tumors by: 1) enhancing the formation of a metabolite (2-methoxy E₂) that is inhibitory to mammary tumor growth, 2) decreasing the formation of fatty acid esters of E₂ metabolites that serve as long term depot forms of hormone in mammary tissue, 3) maintaining the formation E₂ glucuronidation in animals treated chronically with the hormone, and 4) elevation of antioxidant defense enzymes. Thus, PB-induced alterations in hepatic E₂-metabolism leading to the inactivation of E₂ and to the increased activity of antioxidant enzymes need to be considered as determinants for E₂-dependent mammary tumor formation in the ACI rat model.

Acknowledgement

We thank Mrs. Kathy Piano for secretarial assistance and her generous help in maintaining the bibliographic database used in this work.

References

- Cavalieri EL, Stack DE, Devanesan PD, Todorovic R, Dwivedy I, Higginbotham S, Johansson SL, Patil KD, Gross ML, Gooden JK, Ramanathan R, Cerny RL, Rogan EG (1997) Molecular origin of cancer: catechol estrogen-3,4-quinones as endogenous tumor initiators. *Proc Natl Acad Sci USA* 99:10937-10942.
- Cavalieri EL, Rogan EG (2004) A unifying mechanism in the initiation of cancer and other diseases by catechol quinones. *Ann NY Acad Sci* 1028:247-257.
- Conney AH, Jacobson M, Levin W, Kuntzman R (1965) Decreased central depressant effect of progesterone and other steroids in rats pretreated with drugs and insecticides. *J Pharmacol Exp Ther* 154: 310-318, 1966.
- Conney AH, Levin W, Jacobson M, Kuntzman R (1973) Effects of drugs and environmental chemicals on steroid metabolism. Proc Symp drugs and the unborn child. *New York Clin Pharmacol Ther* 14:727-741
- Chakravati D, Pelling JC, Cavalieri EL, Rogan EG (1995) Relating aromatic hydrocarbon-induced DNA adducts and c-Harvey-ras mutations in mouse skin papillomas: The role of apurinic sites. *Proc Natl Acad Sci USA* 92:10422-10426
- El-Bayoumy K, Ji BY, Upadhyaya P, Chae YH, Kurtzke C, Rivenson A, Reddy BS, Amin S, Hecht SS. (1996) Lack of tumorigenicity of cholesterol epoxides and estrone-3,4-quinone in the rat mammary gland. *Cancer Res* 56:1970-1973
- Esposito LA, Kokoszka JE, Waymire KG, Cottrell B, MacGregor GR, Wallace DC (2000) Mitochondrial oxidative stress in mice lacking the glutathione peroxidase-1 gene. *Free Rad*

Biol Med 28:754-766.

Flohe L, Gunzler WA (1984) Assay of glutathione peroxidase. *Meth Enzymol* 105:114-121.

Foldes A, Meek JL (1973) Rat brain phenolsulfotransferase-partial purification and some properties. *Biochem Biophys Acta* 327:365-374.

Fotsis T, Zhang Y, Pepper MS, Adlercreutz H, Montesano R, Nawroth PP, Schweigerer L (1994) The endogenous oestrogen metabolite 2-methoxyestradiol inhibits angiogenesis and suppresses tumor growth. *Nature* 368:237-239

Habig WH, Pabst MJ, Jakoby WB (1974) Glutathione S-transferases: The first enzymatic step in mercapturic acid formation. *J Biol Chem* 249:7130-7139.

Han GZ, Liu ZJ, Shimoi K, Zhu BT (2005) Synergism between the anticancer actions of 2-methoxyestradiol and microtubule-disrupting agents in human breast cancer. *Cancer Res* 65: 387-393.

Harvell DM, Strecker TM, Tochacek M, Xie B, Pennington KL, McComb RD, Roy SK, Shull JD (2000) Rat strain-specific actions of 17-beta-estradiol in the mammary gland: correlation between estrogen-induced lobuloalveolar hyperplasia and susceptibility to estrogen-induced mammary cancers. *Proc Natl Acad Sci USA* 97:2779-2784.

Jaiswal AK, McBride OW, Adesnik M, Nebert DW (1988) Human dioxin-inducible cytosolic NADP(H):menadione oxidoreductase. cDNA sequence and localization of gene to chromosome. *J Biol Chem* 263:3572-13578.

Klauber N, Parangi S, Flynn E, Hamel E, D'Amato RJ (1997) Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and taxol. *Cancer*

Res 57:81-86

Levin W, Welch RM, Conney AH. (1967) Effect of chronic phenobarbital treatment on the liver microsomal metabolism and uterotrophic action of 17-beta-estradiol. *Endocrinology* 80:135-140.

Levin W, Welch RM, Conney AH (1968) Effect of phenobarbital and other drugs on the metabolism and uterotrophic action of estradiol-17 β and estrone. *J Pharmacol Exp Ther* 159:362-371

Levin W, Welch RM, Conney AH (1974) Increased liver microsomal androgen metabolism by phenobarbital: Correlation with decreased androgen action on the seminal vesicles of the rat. *J Pharmacol Exp Ther* 188:287-292

Li SA, Weroha SJ, Tawfik O, Li JJ (2002) Prevention of solely estrogen-induced mammary tumors by Tamoxifen: Evidence for estrogen receptor mediation. *J Endocrinol* 175:297-305

Li KM, Todorovic R, Devanesan P, Higginbotham S, Kofeler H, Ramanathan R, Gross ML, Rogan EG, Cavalieri EL (2004) Metabolism and DNA binding studies of 4-hydroxyestradiol and estradiol-3,4-quinone in vitro and in female ACI rat mammary gland in vivo. *Carcinogenesis* 25:289-297.

Liehr JG, Fang WF, Sirbasku DA, Ari-Ulubelen A (1986) Carcinogenicity of catecholestrogens in Syrian hamsters. *J Steroid Biochem* 24:353-356.

Liehr JG (1997) Hormone-associated cancer: mechanistic similarities between human breast cancer and estrogen-induced kidney carcinogenesis in hamsters. *Environ Health Perspect* 105:565-569.

Liu ZJ, Zhu BT (2004) Concentration-dependent mitogenic and antiproliferative actions of 2-

methoxyestradiol in estrogen receptor-positive human breast cancer cells. *J Steroid Biochem Mol Biol* 88:265-275

Lottering ML, Haag M, Seegers JC (1992) Effects of 17 β -estradiol metabolites on cell cycle events in MCF-7 cells. *Cancer Res* 52:5926-5932

Martucci CP, Fishman J (1993) P450 enzymes of estrogen metabolism. *Pharmacol Ther* 57:237-257

Mesia-Vela S, Sanchez RI, Li J, Li S, Conney AH, Kauffman FC (2002) Liver microsomes from ACI and Sprague-Dawley rats catalyze the 2-hydroxylation of estradiol to a much extent than the 4-hydroxylation. *Carcinogenesis* 23:1369-1372

Mesia-Vela S, Sanchez, RI, Reuhl, K, Conney AH, Kauffman FC (2004) Dietary clofibrate inhibits estrogen-induced antioxidant enzymes in female ACI rats. *Toxicology* 200:103-111

Pasqualini JR (2004) The selective estrogen modulators in breast cancer: a review. *Biochimica Biophysica Acta* 1654:123-143

Pribluda VS, Gubish ER, Lavallo TM, Treston A, Swartz GM, Green SJ (2000) 2-methoxyestradiol: an endogenous antiangiogenic and antiproliferative drug candidate. *Cancer Metastasis Rev* 19:173-179

Radjendirane V, Joseph P, Lee YH, Kimura S, Klein-Szanto AJ, Gonzalez FJ, Jaiswal AK (1998) Disruption of the DT diaphorase (NQO1) gene in mice leads to increased menadione toxicity. *J Biol Chem* 273:7382-7389.

Ramos-Gomez M, Kwak MK, Dolan PM, Itoh K, Yamamoto M, Talalay P, Kensler TW (2001) Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers

- is lost in nrf2 transcription factor-deficient mice. *Proc Natl Acad Sci USA*. 98:3410-3415.
- Reed MJ, Purohit A, Woo, LWL, Newman SP, Potter BVL (2005) Steroid Sulfatase: Molecular Biology, Regulation, and Inhibition. *Endocrine Reviews* 26: 171-202.
- Sanchez RI, Mesia-Vela S, Conney AH, Kauffman FC (2003) Induction of NAD(P)H : quinone oxidoreductase and glutathione S-transferase activities in livers of female August-Copenhagen Irish and Sprague-Dawley rats treated chronically with Estradiol. *J Steroid Biochem Mol Biol* 87:199–206.
- Shah N, Antony T, Haddad S, Amenta P, Shirahata A, Thomas TJ, Thomas T (1999) Antitumor effects of bis(ethyl)polyamine analogs on mammary tumor development in FVB/NTgN (MMTVneu) transgenic mice. *Cancer Lett* 146:15-23
- Shull JD, Spady TJ, Snyder MC, Johansson SL, Pennington KL (1997) Ovary-intact, but not ovariectomized female ACI rats treated with 17-beta-estradiol rapidly develop mammary carcinoma. *Carcinogenesis* 18:1595-1601.
- Suchar LA, Chang RL, Thomas PE, Rosen RT, Lech J, Conney AH (1996) Effects of phenobarbital, dexamethasone and 3-methylcholanthrene administration on the metabolism of 17 β -estradiol by liver microsomes from female rats. *Endocrinology* 137:663-676.
- Thomas PE, Reick LM, Ryan DE, Levin W (1983) Induction of two immunochemically related rat liver cytochrome P450 isozymes cytochromes P450c and P450d, by structurally diverse xenobiotics. *J Biol Chem* 258:4590-4598.
- Turan V, Sanchez RI, Li JJ, Li SA, Reuhl KR, Thomas PE, Conney AH, Gallo MA, Kauffman FC, Mesia-Vela S (2004) The effects of steroidal estrogens in ACI rat mammary carcinogenesis: 17 beta-estradiol, 2-hydroxyestradiol, 4-hydroxyestradiol, 16 alpha-

hydroxyestradiol, and 4-hydroxyestrone. *J Endocrinol* 183: 91-99.

Xu S, Zhu BT, Conney AH (2002) Effect of clofibrate administration on the esterification and deesterification of steroid hormones by liver and extrahepatic tissues in rats. *Biochem Pharmacol* 163:985-992

Zhu BT, Conney AH (1998) Functional role of estrogen metabolism in target cells: review and perspectives. *Carcinogenesis* 19:1-27

Footnotes

Send reprint requests to: F.C. Kauffman, kauffma@rci.rutgers.edu

Financial Support: Provided, in part, by NIH Grant ES05022.

Legends for Figures

Fig. 1. Inhibition of mammary adenocarcinomas by phenobarbital in the ACI rat. E₂ was delivered in cholesterol pellet implants. Animals (8-11 per group) were examined twice a week for tumors and were killed when the tumors reached 3 cm² or at the end of the experiment (28 weeks). PB treatment reduced the incidence of mammary tumors by 47 %.

Fig. 2. Reduction of multiplicity and growth of E₂-induced mammary adenocarcinomas by phenobarbital in the ACI rat. Phenobarbital reduced the multiplicity (tumors/rat; **A**) and growth (tumor size/rat; **B**) of E₂-induced mammary tumors by >95%. The data presented are average values ± S.E. from 8-11 animals per group killed at 28 weeks of treatment.

* *Statistically significant (P < 0.05) when compared to E₂ treated group (unpaired Student t Test).*

Fig. 3. Serum E₂-levels after chronic treatment of ACI rats with E₂. Data are presented as means ± S.E. of 8-11 animals per group.

* *Statistically significant (P < 0.001) when compared with groups not treated with E₂ (unpaired Student's t Test).*

JPET #96867

Table 1. Time course-effects on organ weights of female ACI rats receiving phenobarbital alone or in combination with estradiol

Treatment	<i>Weight</i>						
	Body (g)	Pituitary (mg)	Adrenal s (mg)	Thymus (mg)	Uterus (mg)	Kidneys (g)	Liver (g)
<i>6 weeks</i>							
Water	159 ± 3	9.1 ± 0.6	52 ± 2	251 ± 12	397 ± 43	1.1 ± 0.05	5.3 ± 0.12
0.05% Phenobarbital	157 ± 3	8.3 ± 0.5	66 ± 3*	243 ± 17	373 ± 16	1.2 ± 0.01	6.2 ± 0.12
E ₂	164 ± 3	32 ± 0.7*	55 ± 1	142 ± 4*	418 ± 33	1.5 ± 0.03*,†	7.8 ± 0.31*
0.05% Phenobarbital + E ₂	158 ± 5	29 ± 2*	66 ± 1*,#	127 ± 8*	453 ± 28	1.5 ± 0.04*,†	8.0 ± 0.30*
<i>12 weeks</i>							
Water	173 ± 3	8.6 ± 0.2	59 ± 2	191 ± 5	406 ± 25	1.2 ± 0.03	5.2 ± 0.2
0.05% Phenobarbital	173 ± 5	8.8 ± 1	70 ± 3*	204 ± 15	403 ± 24	1.3 ± 0.05	6.0 ± 0.2
E ₂	171 ± 3	44 ± 5*	58 ± 2	107 ± 13*	532 ± 47	1.5 ± 0.04*	6.6 ± 0.07
0.05% Phenobarbital + E ₂	168 ± 2	38 ± 2*	59 ± 2	112 ± 5*,†	431 ± 11	1.4 ± 0.03*	7.3 ± 0.1

3 mg of E₂ was delivered in a 20 mg cholesterol pellet. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and a 20 mg cholesterol pellet only. Data are presented as means ± S.E. of 8-11 animals per group.

* Statistically different from water control group,

Statistically different from E₂ treated group, † Statistically different from PB treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

JPET #96867

Table 2. Effect of treatment of female ACI rats with PB or E₂ for 6 or 12 weeks on liver weight and on liver microsomal protein per rat.

<i>Treatment</i>	<i>Percent liver weight</i>	<i>Microsomal protein per g liver (mg/g)</i>	<i>Microsomal protein per liver (mg/liver)</i>
6 weeks			
Water	3.4 ± 0.05	13 ± 1	67 ± 6
0.05% Phenobarbital	4.0 ± 0.05*	22 ± 2*	137 ± 12*
E ₂	4.7 ± 0.1*	10 ± 0.7	75 ± 6
0.05% Phenobarbital + E ₂	5.1 ± 0.08* [#]	18 ± 1* [#]	146 ± 20* [#]
12 weeks			
Water	2.9 ± 0.07	12 ± 0.7	62 ± 4
0.05% Phenobarbital	3.5 ± 0.005*	27 ± 3*	162 ± 15*
E ₂	3.9 ± 0.07*	10 ± 2	64 ± 10
0.05% Phenobarbital + E ₂	4.4 ± 0.07* [#]	25 ± 1* [#]	184 ± 10* [#]

3 mg of E₂ was delivered in a 20 mg cholesterol pellets. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and a 20 mg cholesterol pellet only. Percent liver weight = (liver weight /body weight) x 100

Data are presented as means ± S.E. of six animals per group

* Statistically different from water control group

[#] Statistically different from E₂ treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

Table 3. Effect of phenobarbital administration on the NADPH-dependent oxidation of estradiol by liver microsomes from ACI rats

Treatment	Estradiol metabolites formed (pmol/mg protein/min)									E ₂ metabolized	
	<i>6α-OH-E₂</i>	<i>14α-OH-E₂</i>	<i>E₃</i>	<i>6β-OH-E₂</i>	<i>6-keto-E₁</i>	<i>4-OH-E₂</i>	<i>2-OH-E₂</i>	<i>2-OH-E₁</i>	<i>E₁</i>	pmol/mg/min (%) ^a	nmol/liver/min
6 weeks											
Water	0 ± 0	6 ± 0.6	16 ± 1	2 ± 0.7	7 ± 3	9 ± 2	123 ± 11	24 ± 7	134 ± 13	392 ± 23 (17%)	28 ± 3
0.05% Phenobarbital	6 ± 2*	19 ± 2	27 ± 6	33 ± 3*	25 ± 3*	59 ± 1*	314 ± 13*	114 ± 18*	122 ± 20	884 ± 56* (29%)	115 ± 11*
E ₂	0 ± 0	5 ± 1	15 ± 2	1 ± 0.5	8 ± 2	11 ± 2	149 ± 10	32 ± 6	232 ± 7*	530 ± 28 (25%)	40 ± 4
0.05% Phenobarbital + E ₂	4 ± 1* [#]	34 ± 5* [#]	28 ± 5	42 ± 3 [#]	29 ± 3*	36 ± 2 [#]	434 ± 37* [#]	98 ± 8* [#]	108 ± 15 [#]	1053 ± 39* [#] (38%)	145 ± 12* [#]
12 weeks											
Water	0 ± 0	9 ± 1	17 ± 1	2 ± 1	10 ± 3	7 ± 1	169 ± 12	41 ± 7	140 ± 28	473 ± 33 (24%)	30 ± 3
0.05% Phenobarbital	8 ± 1*	21 ± 4*	26 ± 2	33 ± 3*	26 ± 3*	32 ± 4*	418 ± 27*	114 ± 19*	94 ± 10*	1002 ± 33* (31%)	166 ± 16*
E ₂	0 ± 0	6 ± 1	17 ± 2	1 ± 0.5	13 ± 3	11 ± 3	182 ± 36	102 ± 26	309 ± 42*	644 ± 64 (32%)	51 ± 10
0.05% Phenobarbital + E ₂	4 ± 1* [#]	26 ± 4* [#]	27 ± 1	38 ± 2* [#]	25 ± 5*	36 ± 3* [#]	427 ± 2* [#]	156 ± 14* [#]	90 ± 15* [#]	1096 ± 42* [#] (37%)	179 ± 15* [#]

3 mg of E₂ was delivered in a 20 mg cholesterol pellets. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and a 20 mg cholesterol pellet. Liver microsomes (0.5 mg protein) were incubated at 37 °C for 20 min with 50 μM [³H]-E₂, 2 mM NADPH, 5 mM ascorbic acid. Each value is mean ± S.E. obtained from liver microsomes from 6 ACI rats.

Percentage of E₂ metabolized

* Statistically different from water control group

[#] Statistically different from the E₂ treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

JPET #96867

Table 4. Effect of treatment of ACI rat with PB or E₂ on the microsomal conjugation of estradiol by liver microsomes.

Treatment	<i>Conjugate formed (pmol/mg/min)</i>		
	<i>Esterification</i>	<i>Glucuronidation</i>	
	<i>E₂-oleoyl-ester</i>	<i>E₂-3-glucuronide</i>	<i>E₂-17-glucuronide</i>
6 weeks			
Water	13 ± 1	138 ± 7	91 ± 8
0.05% Phenobarbital	21 ± 10	126 ± 14	117 ± 17
E ₂	17 ± 4	118 ± 6	64 ± 2
0.05% Phenobarbital + E ₂	8 ± 0.2* [#]	171 ± 16 [#]	138 ± 14 [#]
12 weeks			
Water	16 ± 3	117 ± 11	82 ± 7
0.05% Phenobarbital	5 ± 1*	64 ± 8*	54 ± 4*
E ₂	14 ± 2	132 ± 12	75 ± 9
0.05% Phenobarbital + E ₂	7 ± 1* [#]	139 ± 18 [†]	120 ± 14 ^{#,†}

3 mg of E₂ was delivered in a 20 mg cholesterol pellet. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and a 20 mg cholesterol pellet. Liver microsomes (1 mg/ml protein) were incubated as described in Material and Methods section. Data are mean ± S.E. obtained from microsomes from six animals per group

* Statistically different from water control group,

[#] Statistically different from E₂ treated group,

[†] Statistically different from PB treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

JPET #96867

Table 5. Effect of treatment of ACI rats with PB or E₂ on sulfotransferase (SULT 1A1) activity in liver cytosol

Treatment	<i>p</i> -nitrophenol sulfate formed (<i>pmol/mg/min</i>)
6 weeks	
Water	147 ± 12
0.05% Phenobarbital	134 ± 7
E ₂	93 ± 4 *
0.05% Phenobarbital + E ₂	82 ± 1*
12 weeks	
Water	134 ± 18
0.05% Phenobarbital	111 ± 8
E ₂	98 ± 12*
0.05% Phenobarbital + E ₂	66 ± 5 *

3 mg of E₂ was delivered in a 20 mg cholesterol pellets. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and a 20 mg cholesterol pellet. Liver cytosols (25 µg) were incubated at 37 °C for 30 min with 5 µM *p*-nitrophenol as substrate as described in Material and Methods section. Data are presented as mean ± S.E. of five animals per group.

* Statistically different from water control group

Statistically different from E₂ treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

JPET #96867

Table 6. Effect of treatment of ACI rats with PB or E₂ on antioxidant enzyme activity in liver cytosol

Treatment	Enzyme Activity (nmol/mg/min)		
	<i>NQO1</i>	<i>GPx</i>	<i>GST</i>
6 weeks			
Water	713 ± 21	1527 ± 24	1001 ± 70
0.05% Phenobarbital	2986 ± 155*	1092 ± 49*	2668 ± 191*
E ₂	1888 ± 69*	1471 ± 90	2001 ± 125*
0.05% Phenobarbital + E ₂	3476 ± 152* [#]	1156 ± 39 [#]	3056 ± 160* [#]
12 weeks			
Water	446 ± 33	1020 ± 60	879 ± 59
0.05% Phenobarbital	1670 ± 181*	787 ± 67	2549 ± 379*
E ₂	1664 ± 88*	1029 ± 189	1665 ± 83*
0.05% Phenobarbital + E ₂	2793 ± 116* [#]	738 ± 83	3462 ± 428* [#]

3 mg of E₂ was delivered in a 20 mg cholesterol pellets. 0.05% sodium phenobarbital was given in the drinking water. Control animals received water and received a 20 mg cholesterol pellet only. Data are presented as means ± S.E. of six animals per group

* Statistically different from water control group

[#] Statistically different from E₂ treated group (ANOVA followed by Bonferroni hoc post test) (P < 0.05)

Figure 1




Figure 2




Figure 3

