

JPET #171298

Calpain inhibitor protects cells against light-induced retinal degeneration

Shunsuke Imai[#], Masamitsu Shimazawa[#], Tomohiro Nakanishi, Kazuhiro Tsuruma,
Hideaki Hara

Molecular Pharmacology, Department of Biofunctional Evaluation, Gifu
Pharmaceutical University, 1-25-4 Daigaku-nishi, Gifu 501-1196, Japan (*S.I., M.S., T.N.,
K.T., H.H.*).

JPET #171298

a) Running title: SNJ-1945 Protects against Light-induced Retinal Damage

b) Corresponding author: Hideaki Hara, Molecular Pharmacology, Department of Biofunctional Evaluation, Gifu Pharmaceutical University, 1-25-4 Daigaku-nishi, Gifu 501-1196 Japan. e-mail: hidehara@gifu-pu.ac.jp, Tel. and fax.: +81-58-230-8126

c) Number of text pages: 31

Number of tables: 0

Number of figures: 6

Number of references: 34

Number of words in Abstract: 210

Number of words in Introduction: 431

Number of words in Discussion: 1284

d) Abbreviation: SNJ-1945,

((1S)-1-(((1S)-1-benzyl-3-cyclopropylamino-2,3-di-oxopropyl)amino)carbonyl)-3-met

hylbutyl)carbamic acid 5-methoxy-3-oxapentyl ester; ERG, electroretinogram

e) Recommended section: Cellular and Molecular Pharmacology

JPET #171298

Abstract

Calpains are activated by excessive light exposure, and related to retinal degeneration.

We investigated the protective effects of

((1S)-1-((((1S)-1-benzyl-3-cyclopropylamino-2,3-di-oxopropyl)amino)carbonyl)-3-methylbutyl)carbamic acid 5-methoxy-3-oxapentyl ester (SNJ-1945), a calpain inhibitor, against light-induced retinal degeneration in mice. SNJ-1945 was orally administrated at doses of 100 and 200 mg/kg at 30 min before and just after light exposure.

Light-induced calpain activation was evaluated by using proteolysis of α -spectrin and p35 (a neuron-specific activator for cyclin-dependent kinase 5). The effects of SNJ-1945 against light-induced retinal damage were examined by the thickness of the outer nuclear layer (ONL). Photoreceptor apoptosis was assessed by counting terminal deoxynucleotidyl transferase dUTP nick-end labeling (TUNEL)-positive cells in ONL. Retinal functions were measured in terms of a- and b-wave amplitudes by using an electroretinogram (ERG). As the mechanism of SNJ-1945, caspase-3/7 measurement was carried out. SNJ-1945 inhibited the proteolysis of α -spectrin and p35 by light exposure and also presented a decrease in the numbers of TUNEL-positive cells and ONL atrophy. Furthermore, SNJ-1945 presented a decrease in a- and b-wave amplitude and caspase-3/7 activation induced by light exposure. These findings suggest that the activation of calpain plays a pivotal role in photoreceptor degeneration by light exposure, and SNJ-1945 may be one of the candidates for the effective treatment of diseases related to photoreceptor degeneration.

JPET #171298

Introduction

When retinal photoreceptors are injured by exposure to excessive light (Noell et al., 1966), retinal damage is irreversible and leads to serious visual field loss.

Photoreceptor loss is the primary cause of blindness in degenerative diseases such as age-related macular degeneration (AMD) and retinitis pigmentosa (RP). However, in these diseases, there are few usable therapeutic agents. To search for a candidate compound against retinal diseases, we used a light-induced photoreceptor degeneration model, and studied the mechanism of progressive disease and drug efficacy.

Calpains are calcium-activated cysteine proteases that occur during the apoptosis process (Utz and Anderson, 2000). The calpain family is represented by 15 genes in mammals, and is comprised of two major isozymes (namely, μ - and m-calpain) which have been well-characterized (Saido et al., 1994; Evans and Turner, 2007). The calpain family is related to many human diseases, such as Alzheimer's disease, brain ischemia, and cataracts (Huang and Wang, 2001). Furthermore, the involvement of calpain activation was reported in retinal diseases such as glaucoma, AMD, and RP (Paquet-Durand et al., 2007; Tamada et al., 2007). Pharmacological inhibition of calpain activation protects from retinal neuronal degeneration (Azuma and Shearer, 2008). Therefore, calpains may be a therapeutic target for treatment of retinal disorders.

A calpain inhibitor, (2S)-4-Methyl-2-[[N-[(4-fluorophenyl) sulfonyl]-L-valyl] amino]pentanal (SJA6017), inhibits potently calpain-1 and -2, and showed a protective effect against ischemia-reperfusion induced retinal cell death (Sakamoto et al., 2000). However, SJA6017 had low oral bioavailability and required high-dose administration (500 mg/kg, orally). On the other hand, a novel calpain inhibitor,

JPET #171298

((1S)-1-(((1S)-1-benzyl-3-cyclopropylamino-2,3-dioxopropyl)amino)carbonyl)-3-methylbutyl)carbamic acid 5-methoxy-3-oxapentyl ester (SNJ-1945) was synthesized from SJA6017 (Shirasaki et al., 2005). SNJ-1945 has more favorable retinal pharmacokinetics, such as good retinal penetration, high oral bioavailability, and long half-life (Shirasaki et al., 2006). In our laboratory, it has been reported that SNJ-1945 has a protective effect against cerebral ischemia-induced neuronal cell death (Koumura et al., 2008) and *N*-methyl-D-aspartate (NMDA)-induced retinal cell death (Shimazawa et al., 2010). Interestingly, SNJ-1945 showed protective effects by oral administration in these models. Furthermore, Oka et al. have reported that oral administration of SNJ-1945 protects against *N*-methyl-*N*-nitrosourea (MNU)-induced photoreceptor degeneration in rats. On the other hand, Perche et al. (2009) reported that another calpain inhibitor, intravitreal injection of Mu-Phe-hPhe-FMK, a calpain inhibitor, did not show any neuroprotective effects against light damage. Therefore, we investigated whether calpain activation would participate in photoreceptor degeneration and retinal dysfunction after light exposure.

In the present study, to demonstrate the protective effects of SNJ-1945, a calpain inhibitor, against photoreceptor cell death induced by light exposure, we examined histological and electrophysiological analyses and the underlying mechanism using an *in vivo* mouse model.

JPET #171298

Materials and methods

Animals. All experiments were performed in accordance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research, and they were approved and monitored by the Institutional Animal Care and Use Committee of Gifu Pharmaceutical University. Male albino ddY mice (Japan SLC, Inc., Hamamatsu, Japan), aged 9 to 10 weeks, were used in this study. They were kept under controlled lighting conditions (12 h:12 h light/dark).

Exposure to Light. After dark adaptation for 24 h, the pupils were dilated with 1% cyclopentolate hydrochloride eye drops (Santen, Osaka, Japan) 30 min before exposure to light. Non-anesthetized mice were exposed to 8,000 lux of white fluorescent light (Toshiba, Tokyo, Japan) for 3 h in cages with a reflective interior. The temperature during exposure to light was maintained at $25 \pm 1.5^{\circ}\text{C}$. After the exposure to light, all mice were returned to darkness for 24 h and then placed in the normal light/dark cycle.

Treatment with SNJ-1945. The SNJ-1945, ((1S)-1-(((1S)-1-benzyl-3-cyclopropylamino-2,3-di-oxopropyl)amino)carbonyl)-3-methylbutyl)carbamic acid 5-methoxy-3-oxapentyl ester, was a gift from Senju Pharmaceutical Co. Ltd. (Kobe, Japan) and it was suspended in distilled water containing 0.5% carboxymethyl cellulose (CMC). SNJ-1945 was orally administrated at 100 or 200 mg/kg, or intraperitoneally administrated at 100 mg/kg, at 30 min before and just after exposure to light. The vehicle group was treated with 10 mL/kg CMC.

Electroretinogram. Electroretinograms (ERG) were recorded at 5 days after light

JPET #171298

exposure (Mayo, Aichi, Japan). Mice were maintained in a totally dark room for 24 h. They were intraperitoneally anesthetized with a mixture of ketamine (120 mg/kg) (Daiichi-Sankyo, Tokyo, Japan) and xylazine (6 mg/kg) (Bayer Health Care, Tokyo, Japan). Pupils were dilated with 1% tropicamide and 2.5% phenylephrine (Santen). Flash ERG was recorded in the left eyes of dark-adapted mice by placing a gold ring electrode (Mayo) in contact with the cornea and a reference electrode (Nihon Kohden, Tokyo, Japan) through the tongue. A neutral electrode (Nihon Kohden) was inserted subcutaneously near the tail. All procedures were performed in dim red light, and the mice were kept warm during the entire procedure. The amplitude of the a-wave was measured from the baseline to the maximum a-wave peak, and the b-wave was measured from the maximum a-wave peak to the maximum b-wave peak.

Histological Analysis. In mice under anesthesia produced by an intraperitoneal injection of sodium pentobarbital (80 mg/kg) (Nakalai Tesque, Kyoto, Japan), each eye was enucleated and kept immersed in a fixative solution containing 4% paraformaldehyde for at least 24 h at 4°C. Six paraffin-embedded sections (thickness 5 μ m) cut through the optic disc of each eye were prepared in a standard manner, and stained with hematoxylin and eosin. The damage induced by light exposure was then evaluated, with six sections from each eye being used for the morphometric analysis described below. Light-microscope images were photographed, and the thickness of the ONL from the optic disc was measured at 240 μ m intervals by photograph in a masked fashion by a single observer (S.I.). Data from three sections (selected randomly from the six sections) were averaged for each eye.

JPET #171298

TUNEL Staining. TdT-mediated biotin-dUTP nick end labeling (TUNEL) staining was performed according to the manufacturer's protocols (In Situ Cell Death Detection Kit; Roche Biochemicals, Mannheim, Germany) to detect retinal cell death induced by exposure to light. The mice were anesthetized with pentobarbital sodium at 80 mg/kg, i.p. at 48 h after exposure to light for 3 h. The eyes were enucleated, fixed overnight in 4% paraformaldehyde, and immersed for 2 days in 25% sucrose with phosphate-buffered saline (PBS). The eyes were then embedded in a supporting medium for frozen-tissue specimens (OCT compound; Tissue-Tek, IL, USA). Retinal sections were cut at 10- μ m thick on a cryostat at -25°C, and stored at -80°C until staining. After washing twice with PBS, sections were incubated with terminal deoxyribonucleotidyl transferase (TdT) enzyme at 37°C for 1 h. The sections were washed 3 times in PBS for 1 min at room temperature. After washing twice with PBS, fluorescence images were photographed, and the intensity was measured in the ONL at a distance between 480 and 720 μ m from the optic disc obtained from the superior area of the retina. The intensity of TUNEL-positive cells was averaged for these superior areas.

Western Blot Analysis. *In vivo*, mice were euthanized using sodium pentobarbital at 80 mg/kg, i.p. and their eyeballs were quickly removed. The retinas were carefully separated from the eyeballs and quickly frozen in dry ice. For protein extraction, the tissue was homogenized in cell-lysis buffer [RIPA buffer (R0278; Sigma-Aldrich, St. Louis, MO, USA) with protease (P8340; Sigma-Aldrich) and phosphatase inhibitor cocktails (P2850 and P5726; Sigma-Aldrich), and 1 mM EDTA] using a homogenizer (Physcotron; Microtec Co. Ltd., Chiba, Japan). The lysate was centrifuged at

JPET #171298

12,000×g for 20 min and the supernatant was used for this study. The protein concentration was measured by comparison with a known concentration of bovine serum albumin using a BCA Protein Assay Kit (Pierce Biotechnology, Rockford, IL, USA). A mixture of equal parts of an aliquot of protein and sample buffer with 10% 2-mercaptoethanol was subjected to 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The separated protein was then transferred onto a polyvinylidene difluoride membrane (Immobilon-P; Millipore Corporation, Bedford, MA, USA). For immunoblotting, the following primary antibodies were used: mouse anti- α -spectrin monoclonal antibody (clone AA6; Millipore, Billerica, MA, USA) (1:2000), p35/25 rabbit monoclonal antibody (Cell Signaling, Danvers, MA, USA) (1:1000), β -actin mouse monoclonal antibody (Sigma-Aldrich) (1:4000). The secondary antibody used was either goat anti-rabbit HRP-conjugated (1:2000) or goat anti-mouse HRP-conjugated (1:2000). The immunoreactive bands were visualized using SuperSignal West Femto Maximum Sensitivity Substrate (Pierce Biotechnology). The band intensity was measured using a Lumino Imaging Analyzer (Fujifilm, Osaka, Japan).

Caspase-3/7 Assay. *In vivo*, mice were euthanized using sodium pentobarbital at 80 mg/kg, i.p. and their eyeballs were quickly removed. The retinas were carefully separated from the eyeballs and quickly frozen in liquid nitrogen. For protein extraction, the tissue was homogenized in cell-lysis buffer using a homogenizer (Phycotron; Microtec Co. Ltd., Chiba, Japan). The lysate was centrifuged at 12,000×g for 20 min and the supernatant used for this study. The protein concentration was measured by comparison with a known concentration of bovine serum albumin using a BCA Protein Assay Kit (Pierce Biotechnology). Caspase-3/7

JPET #171298

was measured by using Caspase-Glo[®] 3/7 Assay (Promega, Madison, WI, USA) according to the manufacturer's instructions. The luminescence of each sample was measured in a plate-reading microplate.

JPET #171298

Results

Effects of SNJ-1945 on the activation of calpain by light exposure

To investigate the activation of calpain after light exposure, proteolysis of the cytoskeletal protein α -spectrin was measured by Western blot (Fig. 1A and 1B). At 24 h after light exposure, the proteolyzed α -spectrin band at 150 kDa became clear, and a new band of proteolyzed α -spectrin at 145 kDa appeared. Just after light exposure, calpain tended to be activated by light injury. Treatment with SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg, i.p.) significantly inhibited the two cleaved products (145 kDa and 150 kDa) at 24 h after light exposure (Fig. 1C and 1D).

Effects of SNJ-1945 against the degradation of p35 by light exposure

To investigate calpain-induced photoreceptor degeneration, p35 (Cdk5 regulator protein) was measured in light-exposed mouse retinas. Time-dependent degradation of p35 after light exposure was shown in Fig. 2A. In non-treated normal retinas, p35 protein was expressed, and in light-exposed retinas, p35 was decreased at 48 and 72 h after light exposure; p35 was significantly reduced (Fig. 2B). Treatment with SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg, i.p.) inhibited the degradation of p35 at 48 h after light exposure (Fig 2C, D).

Effects of SNJ-1945 on light-induced photoreceptor degeneration

In histological evaluation, Figure 3A-3E shows representative retinal images between 480 μ m to 720 μ m from the optic nerve in the superior area at 5 days after light exposure. The ONL thickness was remarkably thinned in the vehicle-treated group (Fig. 3B) versus the non-treated normal group (Fig. 3A). The group treated with

JPET #171298

SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg i.p.) showed significantly suppressed photic damage (Fig. 3D and 3E) versus the vehicle-treated group (Fig. 3F). SNJ-1945, when administered 30 min before and just after light exposure, significantly protected the retinal superior area from 240 μm to 1440 μm , and the inferior area from 240 μm to 960 μm (Fig. 3F).

Light-induced expression of TUNEL-positive cells

To show light-induced apoptotic cell death and the effect of SNJ-1945 against cell death, we investigated the expression of TUNEL-positive cells at 48 h after light exposure (Fig. 4A to 4E). TUNEL-positive cells were observed only at ONL, but not at other retinal areas (Fig. 4A to 4D). On the other hand, in non-treated retina, TUNEL-positive cells were not shown in any retinal area (Fig. 4A). Quantitative analysis showed that exposure to light in mouse retina significantly increased the number of TUNEL-positive cells in ONL at 48 h (vs. non-treated normal retina) (Fig. 4E). SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg, i.p.) significantly reduced the number of TUNEL-positive cells (Fig. 5E).

Electroretinogram (ERG)

The effects of SNJ-1945 on light-induced photoreceptor degeneration were examined by electrophysiologic analyses. Both a- and b-wave amplitudes were significantly reduced at 5 days after 8,000 lux white light exposure for 3 h, and a- and b-wave decreased by 85% and 83%, respectively, as compared with non-treated retina at 0.98 log cds/m² (Fig. 5B, C). At 0.98 log cds/m², the SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg, i.p.) treated group showed significantly less reduction of a- and b-wave

JPET #171298

amplitudes, when compared with the saline-treated group (Fig. 5B, C).

Effect of SNJ-1945 against light-induced caspase-3/7 activation

We investigated the mechanism of photoreceptor degeneration and the effect of SNJ-1945. We measured time-dependent changes in caspase-3/7 activity after light exposure. Caspase-3/7 activity was increased just after light exposure, and decreased at 24 h. However, at 48 h, caspase-3/7 activity was elevated again (Fig. 6A). At 48 h after light exposure, we investigated the effect of SNJ-1945 against light-induced caspase-3/7 activation. Systemic (oral or intraperitoneal) treatment with SNJ-1945 inhibited caspase-3/7 activity (Fig. 6B).

JPET #171298

Discussion

In the present study, we showed the protective effects of SNJ-1945, a novel calpain inhibitor, against light-induced retinal degeneration in mice.

As a result of light irradiation, retinal photoreceptors trigger apoptotic cell death (Abler et al., 1996). One of the causes is elevation of intracellular calcium concentrations and calpain activation. It has been reported that intracellular calcium influx and calpain activation were increased by light exposure, and a calcium channel blocker, flunarizine, was protective from retinal damage by light exposure (Edward et al., 1991; Isayama et al., 1991; Donovan et al., 2001). Furthermore, in other retinal degeneration models such as the retinal degeneration (rd) 1 mouse and MNU-induced retinal degeneration, calpain activation was increased in photoreceptor cells (Paquet-Durand et al., 2006; Oka et al., 2007). In the present study, we examined calpain activation by measuring calpain-specific α -spectrin fragments at 145 kDa in mouse retina after light exposure. The 150 kDa fragments are broken by the effects of calpain and caspase-3, while the 145 kDa fragment is broken only by calpain (Nath et al., 1996). In the retina, α -spectrin is localized in inner and outer retina (Isayama et al., 1991), and the proteolysis of spectrin is thought to be a cause of retinal cell death. We showed that the cleaved products of α -spectrin at 145 kDa increased at 24 h after light exposure, and SNJ-1945 treatment (200 mg/kg, p.o. or 100 mg/kg, i.p.) decreased the proteolysis of α -spectrin. As the other substrate of calpain, we evaluated p35 degradation. The p35 protein is a regulator of cyclin-dependent kinases 5 (Cdk5), and p35 is degraded to p25 by calpain (Lee et al., 2000). Proteolytic cleavage of p35 by calpain produces p25, resulting in prolonged activation of Cdk5 with p25 and hyperphosphorylation of tau protein (Kusakawa et al., 2000). Accumulation of

JPET #171298

hyperphosphorylated tau protein causes neurofibrillary tangles and neuronal cell death. In retina, p35 was proteolyzed to p25 by several factors, such as hypoxia, ocular hypertension, photoreceptor cell death, and ganglion cell death (Tamada et al., 2005; Oka et al., 2006; Oka et al., 2007; Shimazawa et al., 2010). In the present study, p35 was decreased at 48 h after light exposure, and SNJ-1945 treatment (200 mg/kg, p.o. or 100 mg/kg, i.p.) preserved p35 by calpain inhibition. These results indicate that calpain was activated by light exposure, and SNJ-1945 inhibited calpain activation with systemic administration. Interestingly, degradation of α -spectrin and p35 was mildly increased within a few hours after light exposure; afterwards their degradations increased rapidly. These biphasic changes were reported by Perche and colleagues (Perche et al., 2009), and attributed to the endogenous calpain inhibitor, calpastatin, being increased with activation of μ -calpain and m-calpain. In a previous report, photoreceptor degeneration was associated with the difference in the time of μ -calpain and m-calpain activation (Oka et al., 2007). The reason was considered that μ -calpain was activated first, and m-calpain was followed by activation of μ -calpain. Moreover, the results in this study were correlated with the proteolysis of α -spectrin. In the present study, it was confirmed that light exposure induced proteolysis of α -spectrin and p35 by calpain, and therefore we examined whether SNJ-1945 protects from light-induced retinal damage or not. In histological analysis, we evaluated the ONL thickness as photoreceptor atrophy. SNJ-1945 treatment (200 mg/kg, p.o. or 100 mg/kg, i.p.) reduced the loss of ONL, and this result suggests that calpain inhibitors have a protective effect against photic retinal damage. Moreover, the effect of SNJ-1945 was evaluated electrophysiologically by using ERG. At 5 days after light exposure, amplitudes of a- and b-waves were decreased, and SNJ-1945 administration

JPET #171298

(200 mg/kg, p.o. or 100 mg/kg, i.p.) inhibited the reduction in both a- and b-waves. A-waves show the function of photoreceptor, and b-waves show the function of Müller cells and bipolar cells. The results indicated that SNJ-1945 prevented decreased visual function by light exposure. On the other hand, Perche et al. (2009) reported that another calpain inhibitor, Mu-Phe-hPhe-FMK, did not show any neuroprotective effects against light damage. They confirmed that Mu-Phe-hPhe-FMK inhibited calpain activation during light exposure, but it did not show any protective effects on photoreceptor loss, increasing apoptotic cells, or retinal dysfunction. As the reason for the inefficacy, it was suggested that Mu-Phe-hPhe-FMK may have a short-half life in plasma and/or retina. On the other hand, oral administration of SNJ-1945 was reported to show good penetration and long half-life ($T_{1/2} = 4.3$ h) in rat retina. (Shirasaki et al., 2006). Furthermore, Oka et al. (Oka et al., 2007) have reported that oral administration of SNJ-1945 at 200 mg/kg/day, but not at 100 mg/kg/day, protects against MNU-induced photoreceptor degeneration. These data were consistent with our results in Figure 3 and, therefore, SNJ-1945 at 200 mg/kg, p.o. may be an effective dose in photoreceptor degeneration of mice and rats. On the other hand, there was no report about pharmacokinetic information after intraperitoneal administration of SNJ-1945. However, the maximum plasma concentration (C_{max}) after intraperitoneal administration of SNJ-1945 at 100 mg/kg is predicted to be higher than that after the oral administration at 100 mg/kg, and the half-life is shorter than that. Therefore, it is considered that SNJ-1945 at 100 mg/kg, i.p., but not at 100 mg/kg, p.o., was effective in the ONL atrophy after light exposure.

Retinal diseases in human and animal models lead to apoptotic cell death (Nickells and Zack, 1996; Reme et al., 2000) and, in light-induced retinal degeneration, apoptosis

JPET #171298

is a common final pathway (Wenzel et al., 2005). We investigated photoreceptor apoptosis by using TUNEL staining. Light-induced TUNEL positive cells were significantly expressed at 48 h after the light exposure in ONL. SNJ-1945 (200 mg/kg, p.o. or 100 mg/kg, i.p.) inhibited increased expression of TUNEL-positive cells in ONL. This result indicates that light exposure activates calpain, and calpain activation triggers photoreceptor apoptosis. Furthermore, oral (200 mg/kg) and intraperitoneal (100 mg/kg) treatments with SNJ-1945 suppressed apoptotic DNA damage.

During apoptosis processes, caspase-3 has been identified as a key protease in the execution of apoptosis, whereas calpains have been implicated in neuronal death (Blomgren et al., 2001). In both inherited retinal degeneration and light-induced retinal degeneration, caspase-3 inhibitors showed a protective effect, suggesting that photoreceptor degeneration is related to caspase-3 activation (Perche et al., 2007; Perche et al., 2008). For these reasons, we confirmed that caspase-3/7 was activated by light injury, being significant at 48 h after light exposure. Costa et al. (2008) also reported that caspase-3 was up-regulated at 48 h after light exposure, and our result corresponded with those results. In the present study, SNJ-1945 inhibited caspase-3/7 activation at 48 h after light exposure. Taken together, these results suggest that caspase-3 activation is partially mediated by calpain activation after light exposure. However, we do not have any data about specificity of SNJ-1945 against calpain. Some investigators have reported that SNJ-1945 inhibits μ -calpain and m-calpain activities with IC_{50} values of 0.062 μ M and 0.045 μ M, respectively (Shirasaki et al., 2006), and that it also exerts inhibitory actions on Ca^{2+} -independent proteinase and cathepsin L and B (Yoshikawa et al., 2010). SNJ-1945 was synthesized from SJA-6017, a potent calpain inhibitor, for improving its metabolic stability and water

JPET #171298

solubility (Shirasaki et al., 2006). SJA-6017 is well known to inhibit cathepsin B and L in addition to calpain, but does not inhibit other cysteine proteases (interleukin 1 β converting enzyme), serine proteases (trypsin, chymotrypsin, thrombin, factor VIIa, factor Xa), or proteasome (Inoue et al., 2003). Furthermore, a docking study supported that SNJ-1945 bound in the active site pocket of μ -calpain (Azuma and Shearer, 2008). Accordingly, SNJ-1945 is also predicted to have similar profiles with SJA-6017, but further studies will be needed.

In conclusion, we demonstrated that SNJ-1945, a calpain inhibitor, has neuroprotective effects against light-induced retinal damage. SNJ-1945 may be a candidate compound for photoreceptor degeneration-related diseases.

JPET #171298

Acknowledgements

The authors wish to express their gratitude to Dr. Yoshiyuki Tamada, Senju Laboratory of Ocular Sciences, Senju Pharmaceutical Co., Ltd., Kobe, Japan for the kind gift of SNJ-1945 and useful advice.

JPET #171298

References

- Abler AS, Chang CJ, Ful J, Tso MO and Lam TT (1996) Photic injury triggers apoptosis of photoreceptor cells. *Res Commun Mol Pathol Pharmacol* **92**:177-189.
- Azuma M and Shearer TR (2008) The role of calcium-activated protease calpain in experimental retinal pathology. *Surv Ophthalmol* **53**:150-163.
- Blomgren K, Zhu C, Wang X, Karlsson JO, Leverin AL, Bahr BA, Mallard C and Hagberg H (2001) Synergistic activation of caspase-3 by m-calpain after neonatal hypoxia-ischemia: a mechanism of "pathological apoptosis"? *J Biol Chem* **276**:10191-10198.
- Costa BL, Fawcett R, Li GY, Safa R and Osborne NN (2008) Orally administered epigallocatechin gallate attenuates light-induced photoreceptor damage. *Brain Res Bull* **76**:412-423.
- Donovan M, Carmody RJ and Cotter TG (2001) Light-induced photoreceptor apoptosis in vivo requires neuronal nitric-oxide synthase and guanylate cyclase activity and is caspase-3-independent. *J Biol Chem* **276**:23000-23008.

JPET #171298

Edward DP, Lam TT, Shahinfar S, Li J and Tso MO (1991) Amelioration of
light-induced retinal degeneration by a calcium overload blocker. Flunarizine.
Arch Ophthalmol **109**:554-562.

Evans JS and Turner MD (2007) Emerging functions of the calpain superfamily of
cysteine proteases in neuroendocrine secretory pathways. *J Neurochem*
103:849-859.

Huang Y and Wang KK (2001) The calpain family and human disease. *Trends Mol Med*
7:355-362.

Inoue J, Nakamura M, Cui YS, Sakai Y, Sakai O, Hill JR, Wang KK and Yuen PW
(2003) Structure-activity relationship study and drug profile of
N-(4-fluorophenylsulfonyl)-L-valyl-L-leucinal (SJA6017) as a potent calpain
inhibitor. *J Med Chem* **46**:868-871.

Isayama T, Goodman SR and Zagon IS (1991) Spectrin isoforms in the mammalian
retina. *J Neurosci* **11**:3531-3538.

Koumura A, Nonaka Y, Hyakkoku K, Oka T, Shimazawa M, Hozumi I, Inuzuka T and

JPET #171298

Hara H (2008) A novel calpain inhibitor,
((1S)-1((((1S)-1-benzyl-3-cyclopropylamino-2,3-di-oxopropyl)amino)carbonyl)-
3-methylbutyl) carbamic acid 5-methoxy-3-oxapentyl ester, protects neuronal
cells from cerebral ischemia-induced damage in mice. *Neuroscience*
157:309-318.

Kusakawa G, Saito T, Onuki R, Ishiguro K, Kishimoto T and Hisanaga S (2000)

Calpain-dependent proteolytic cleavage of the p35 cyclin-dependent kinase 5
activator to p25. *J Biol Chem* **275**:17166-17172.

Lee MS, Kwon YT, Li M, Peng J, Friedlander RM and Tsai LH (2000) Neurotoxicity

induces cleavage of p35 to p25 by calpain. *Nature* **405**:360-364.

Nath R, Raser KJ, Stafford D, Hajimohammadreza I, Posner A, Allen H, Talanian RV,

Yuen P, Gilbertsen RB and Wang KK (1996) Non-erythroid alpha-spectrin
breakdown by calpain and interleukin 1 β -converting-enzyme-like protease(s) in
apoptotic cells: contributory roles of both protease families in neuronal
apoptosis. *Biochem J* **319 (Pt 3)**:683-690.

JPET #171298

Nickells RW and Zack DJ (1996) Apoptosis in ocular disease: a molecular overview.

Ophthalmic Genet **17**:145-165.

Noell WK, Walker VS, Kang BS and Berman S (1966) Retinal damage by light in rats.

Invest Ophthalmol **5**:450-473.

Oka T, Nakajima T, Tamada Y, Shearer TR and Azuma M (2007) Contribution of

calpains to photoreceptor cell death in *N*-methyl-*N*-nitrosourea-treated rats. *Exp*

Neurol **204**:39-48.

Oka T, Tamada Y, Nakajima E, Shearer TR and Azuma M (2006) Presence of

calpain-induced proteolysis in retinal degeneration and dysfunction in a rat

model of acute ocular hypertension. *J Neurosci Res* **83**:1342-1351.

Paquet-Durand F, Azadi S, Hauck SM, Ueffing M, van Veen T and Ekstrom P (2006)

Calpain is activated in degenerating photoreceptors in the rd1 mouse. *J*

Neurochem **96**:802-814.

Paquet-Durand F, Johnson L and Ekstrom P (2007) Calpain activity in retinal

degeneration. *J Neurosci Res* **85**:693-702.

JPET #171298

Perche O, Doly M and Ranchon-Cole I (2007) Caspase-dependent apoptosis in light-induced retinal degeneration. *Invest Ophthalmol Vis Sci* **48**:2753-2759.

Perche O, Doly M and Ranchon-Cole I (2008) Transient protective effect of caspase inhibitors in RCS rat. *Exp Eye Res* **86**:519-527.

Perche O, Doly M and Ranchon-Cole I (2009) Calpains are activated by light but their inhibition has no neuroprotective effect against light-damage. *Exp Eye Res* **89**:989-994.

Reme CE, Grimm C, Hafezi F, Wenzel A and Williams TP (2000) Apoptosis in the Retina: The Silent Death of Vision. *News Physiol Sci* **15**:120-124.

Saido TC, Nagao S, Shiramine M, Tsukaguchi M, Yoshizawa T, Sorimachi H, Ito H, Tsuchiya T, Kawashima S and Suzuki K (1994) Distinct kinetics of subunit autolysis in mammalian m-calpain activation. *FEBS Lett* **346**:263-267.

Sakamoto YR, Nakajima TR, Fukiage CR, Sakai OR, Yoshida YR, Azuma MR and Shearer TR (2000) Involvement of calpain isoforms in ischemia-reperfusion injury in rat retina. *Curr Eye Res* **21**:571-580.

JPET #171298

Shimazawa M, Suemori S, Inokuchi Y, Matsunaga N, Nakajima Y, Oka T, Yamamoto T

and Hara H (2010) A novel calpain inhibitor,

((1S)-1-(((1S)-1-Benzyl-3-cyclopropylamino-2,3-di-oxopropyl)amino)carbonyl

l)-3-methylbutyl)carbamic acid 5-methoxy-3-oxapentyl ester (SNJ-1945),

reduces murine retinal cell death in vitro and in vivo. *J Pharmacol Exp Ther*

332:380-387.

Shirasaki Y, Miyashita H, Yamaguchi M, Inoue J and Nakamura M (2005) Exploration

of orally available calpain inhibitors: peptidyl alpha-ketoamides containing an

amphiphile at P3 site. *Bioorg Med Chem* **13**:4473-4484.

Shirasaki Y, Yamaguchi M and Miyashita H (2006) Retinal penetration of calpain

inhibitors in rats after oral administration. *J Ocul Pharmacol Ther* **22**:417-424.

Tamada Y, Nakajima E, Nakajima T, Shearer TR and Azuma M (2005) Proteolysis of

neuronal cytoskeletal proteins by calpain contributes to rat retinal cell death

induced by hypoxia. *Brain Res* **1050**:148-155.

Tamada Y, Walkup RD, Shearer TR and Azuma M (2007) Contribution of calpain to

JPET #171298

cellular damage in human retinal pigment epithelial cells cultured with zinc chelator. *Curr Eye Res* **32**:565-573.

Utz PJ and Anderson P (2000) Life and death decisions: regulation of apoptosis by proteolysis of signaling molecules. *Cell Death Differ* **7**:589-602.

Wenzel A, Grimm C, Samardzija M and Reme CE (2005) Molecular mechanisms of light-induced photoreceptor apoptosis and neuroprotection for retinal degeneration. *Prog Retin Eye Res* **24**:275-306.

Yoshikawa Y, Zhang GX, Obata K, Ohga Y, Matsuyoshi H, Taniguchi S and Takaki M (2010) Cardioprotective effects of a novel calpain inhibitor SNJ-1945 for reperfusion injury after cardioplegic cardiac arrest. *Am J Physiol Heart Circ Physiol* **298**:H643-651.

JPET #171298

Footnotes

Shunsuke Imai and Masamitsu Shimazawa contributed equally to this article.

JPET #171298

Figure Legends

Figure 1. Changes in proteolysis of α -spectrin after light exposure in the mouse retina and the effects of SNJ-1945.

(A) Representative immunoblots showing proteolysis of α -spectrin in retinal extracts at 0, 3, 6, 12, 24, 48, and 72 h after light exposure in mice. (B) Quantitative analysis of the band density of the cleaved products at 145 and 150 kDa of α -spectrin (280 kDa).

Data are shown as mean \pm S.E.M. (n = 5 or 6). The number in parentheses above each column represents the number of animals. *p < 0.05 vs. non-treated group (Normal).

(C) Representative immunoblots showing proteolysis of α -spectrin in mouse retinal extracts at 24 h after light exposure with or without treatment with SNJ-1945.

SNJ-1945 (at 200 mg/kg, p.o. and 100 mg/kg, i.p.) or an identical volume (10 ml/kg) of vehicle (0.5% sodium carboxymethyl cellulose) was administered at 30 min before and just after light exposure. (D) Quantitative analysis of the band density at

145kDa/150kDa (proteolyzed from α -spectrin at 280 kDa). Data are shown as means \pm S.E.M., n = 4 to 6. The number in parentheses above each column represents the number of animals. *p < 0.05 vs. light exposure plus vehicle-treated group, ##p < 0.01 vs. non-treated group (Normal).

Figure 2. Changes in proteolysis of p35 after light exposure in the mouse retina and the effect of SNJ-1945.

(A) Representative immunoblots showing p35 protein levels in retinal extracts at 48 h after light exposure in mice. (B) Quantitative analysis of the band density at 35 kDa. Data are shown as mean \pm S.E.M. (n = 5 or 6). The number in parentheses above each column represents the number of animals. *p < 0.05 vs. non-treated group (Normal).

JPET #171298

(C) Representative immunoblots showing proteolysis of p35 in mouse retinal extracts at 48 h after light exposure with or without treatment with SNJ-1945. SNJ-1945 (at 200 mg/kg, p.o. and 100 mg/kg, i.p.) or an identical volume (10 ml/kg) of vehicle (0.5% sodium carboxymethyl cellulose) was administered at 30 min before and just after light exposure. (D) Quantitative analysis of the band density at p35/actin. Each column represents the mean \pm S.E.M. ($n = 5$ or 6). The number in parentheses above each column represents the number of animals. * $p < 0.05$ vs. light exposure plus vehicle-treated group; # $p < 0.05$ vs. non-treated group (Normal).

Figure 3. Effects of SNJ-1945 on retinal damage induced by exposure to light in mice. (A) Nontreated, (B) light exposure (8,000 lux) plus vehicle-treated, and (C, D, and E) light exposure plus SNJ-1945 treated (100 mg/kg, p.o., 200 mg/kg, p.o. and 100 mg/kg, i.p., respectively) retinal cross sections at 5 days after light exposure in mice. (F) Measurement of thickness in the outer nuclear layer (ONL) at 5 days after light exposure. Data are shown as means \pm SEM, $n = 6$ or 8 . * $p < 0.05$, ** $p < 0.01$ vs. light exposure plus the vehicle-treated group. The scale bar represents 25 μm .

Figure 4. Effects of SNJ-1945 on expression of TUNEL-positive cells at 48 h after light exposure.

Representative photographs of TUNEL staining showing (A) non-treated normal retina, (B) light exposure (8,000 lux) plus vehicle-treated, (C) light exposure plus SNJ-1945 at 200 mg/kg, p.o. and (D) light exposure plus SNJ-1945 at 100 mg/kg, i.p. (E) Quantitative analysis of intensity of TUNEL-positive cells in outer nuclear layer at 48 h after light exposure. Data are shown as means \pm S.E.M., $n = 7$ or 8 . The number in

JPET #171298

parentheses above each column represents the number of animals. * $p < 0.05$ vs. light exposure plus vehicle-treated group. ## $p < 0.01$ vs. non-treated group (Normal).

Scale bar represents 25 μm . ONL: outer nuclear layer.

Figure 5. Effects of SNJ-1945 on changes in the dark adapted-ERG amplitudes after exposure to light in the mouse retina.

(A) Typical traces of dark-adapted ERG responses measured at 5 days after exposure to light. Stimulus flashes were used at 0.98 log cds/m². (B and C) Amplitudes of a- and b-waves of light exposure (8,000 lux) plus vehicle-treated group vs. light exposure plus SNJ-1945 at 200 mg/kg, p.o. and at 100 mg/kg, i.p. treated group. Data are shown as means \pm S.E.M., $n = 6$ or 8 . The number in parentheses above each column represents the number of animals. * $p < 0.05$, ** $p < 0.01$ vs. light exposure plus vehicle-treated group.

Figure 6. Effects of SNJ-1945 on light-induced expression of caspase-3 in the mouse retina.

(A) Quantitative analysis of the fluorescent intensity of caspase-3 activation. Data are shown as mean \pm S.E.M. ($n = 5$ or 6). The number in parentheses above each column represents the number of animals. * $p < 0.05$, ** $p < 0.01$ vs. non-treated group

(Normal). (B) Quantitative analysis of the effect of SNJ-1945 against activation of caspase-3 at 48 h after light exposure. SNJ-1945 (at 200 mg/kg, p.o. and 100 mg/kg, i.p.) or an identical volume (10 ml/kg) of vehicle (0.5% sodium carboxymethyl

JPET #171298

cellulose) was administered at 30 min before and just after light exposure. Each column represents the mean \pm S.E.M. (n = 5 or 6). The number in parentheses above each column represents the number of animals. ##p < 0.01 vs. non-treated group (Normal), * p < 0.05 vs. vehicle.











