# CONTENTS

Number 1, I	EBRUARY,	1921
-------------	----------	------

I. On the Influence of Colloids on the Action of Non-Colloidal Drugs. I. By W. Storm van Leeuwen	1
II. Effect of Some Antipyretics on the Behavior of Rats in the Circular	2
Maze. By D. I. Macht and Wm. Bloom	21
Cushny	41
By Torald Sollmann	63 103
VI. Some Observations upon the Behavior of a Fixed Oil (Peanut Oil) Injected Intraperitoneally. By Erich W. Schwartze	118
VII. On the Influence of Colloids on the Action of Non-Colloidal Drugs. II.  By W. Storm van Leeuwen and J. Zeijdner	
VIII. A Comparison of the Action of Certain Drugs upon Muscular Work	
in Frogs. By Eleanor M. Scarborough	
By Herbert Olivecrona.	141
. Number 3, April, 1921	
<ul> <li>X. Studies on the Influence of Sedatives on Animals. I. A Method for Measuring the Influence of Stimulating Drugs and of Sedatives on the Activity of Animals (Activitymeter). By W. Storm van Leeuwen</li> <li>XI. On the Duration of Construction of Bloodvessels by Epinephrine. By S. J. Meltzer and John Auer</li> <li>XII. Studies of Chronic Intoxications on Albino Rats. IV. Fluorid, Chlorid and Calcium, Including Sodium Fluorid, Sodium Chlorid, "Phosphate Rock," Calcium Phosphate (Precipitated) and Calcium Carbonate (Precipitated). By Torald Sollmann, O. H. Schettler and N. C. Wetzel.</li> <li>XIII. The Action of Drugs upon the Output of Epinephrin from the Adrenals. VII. Physostigmine. By G. N. Stewart and J. M. Rogoff</li> <li>XIV. Observations on the Effect of Ipecac in the Treatment of Infectious Entero-Hepatitis (Blackhead) in Turkeys. By Harry M. Wegeforth and Paul Wegeforth</li> </ul>	177 197 227
Number 4, May, 1921	
XV. The Comparative Toxicity of Thymol and Carvacrol (Isothymol).  By A. E. Livingston	261
. Dooley	277
iii .	

### CONTENTS

XVIII. Scientific Proceedings of the American Society for Pharmacology	289 325
Number 5, June, 1921	
XIX. The Toxicity of Some Thioureas and Thiuramdisulphides. By P. J. Hanzlik and A. Irvine	
Salicylate in the Circulation. By P. J. Hanzlik	395
Number 6, July, 1921	
XXV. Route of Administration of Drugs in Relation to Toxicity in Chemotherapeutic Investigations with Special Reference to Intrapleural Injections of Ethylhydrocuprein Hydrochloride. By John A. Kolmer	431
XXVI. Comparative Effects of Morphin and Alkaloids of the Benzyliso- quinolin Group on Cardiac Muscle. By P. J. Hanzlik	
Glands of Individuals with Exophthalmic Goiter and Other Conditions Involving the Thyroid. By J. M. Rogoff and H. Goldblatt	473 481

# ILLUSTRATIONS

Concentration-action curve, type I (Fig. 1)	2
action curve, type II (Fig. 2)	2
Adsorption curves of benzoic acid and succinic acid (Fig. 3)	3
	4
— of nicotine by animal charcoal (Fig. 5)	5
isotherm (logarithmated) of pilocarpine by animal charcoal (Fig. 6)	6
Strip of cat gut suspended in 75 cc. of Tyrode solution (Fig. 7, a-d)	9
Isolated strip of cat gut suspended in 75 cc. Tyrode solution (Fig. 8, a-e)	10
Amount and concentration of pilocarpine (Fig. 9)	13
Isolated cat gut suspended in 75 cc. of Tyrode solution (Fig. 10, a-c)	15
Circular maze viewed from above (Fig. 1)	22
— maze with camera lucida attachment (Fig. 2)	23
Graphs of the secretion of saliva by a dog from 5 mgm. of pilocarpine nitrate	
injected subcutaneously at $P$ , preceded by hyoscine hydrobromide at $H$	
(Fig. 1)	45
— of the secretion of saliva by another dog from 5 mgm. pilocarpine nitrate	
injected subcutaneously at $P$ , preceded by hyoscine hydrobromide at $H$	
(Fig. 2)	46
Effect of hyoscines on the movements of the surviving bowel in 50 cc. of	
oxygenated Ringer's solution (exp. 1), (Fig. 3)	49
Tracing of surviving intestine of rabbit in 50 cc. of Tyrode's solution (Fig. 4).	49
Records of movements of a piece of surviving intestine of rabbit in Ringer's	
solution (Fig. 5)	51
Time factor in solution of tannin-protein compounds (Fig. 1)	81
— factor in solution of tannin-protein compounds (Fig. 2)	82
—— factor in hydrolysis of acetannins (Fig. 3)	90
Length of life (in 48-hour periods) of rabbits given potassium arsenite alone,	
and of animals given potassium arsenite and later magnesium sulphate	
(Fig. 1)	111
Influence of colloids on non-colloidal drugs (Fig. 1, a-h)	
Action of drugs upon muscular work of frogs (Fig. 1)	
of drugs upon muscular work of frogs (Fig. 2)	
of drugs upon muscular work of frogs (Fig. 3)	
of drugs upon muscular work of frogs (Fig. 4)	
— of drugs upon muscular work of frogs (Fig. 5)	
Control fatigue curve; perfusion with Ringer's solution (Fig. 6)	134
Fatigue curve; perfusion with Ringer's solution (Fig. 7)	
curve; perfusion with Ringer's solution (Fig. 8)	
curve; perfusion with Ringer's solution (Fig. 9)	
curve; perfusion with Ringer's solution (Fig. 10)	
curve; perfusion with Ringer's solution (Fig. 11)	136

### ILLUSTRATIONS

Small intestine of cat (Fig. 1)	146
—— intestine of cat (Fig. 2)	147
—— intestine of cat (Fig. 3)	148
—— intestine of cat (Fig. 4)	
—— intestine of cat (Fig. 5)	
— intestine of cat (Fig. 6)	152
—— intestine of cat (Fig. 7)	153
intestine of rabbit (Fig. 8)	157
intestine of rabbit (Fig. 9)	157
— intestine of rabbit (Fig. 10)	157
, 0 /	
intestine of rabbit (Fig. 12)	158
intestine of cat (Fig. 13)	160
—— intestine of cat (Fig. 14)	160
Uterus of guinea-pig (Fig. 15)	161
Isolated loop of small intestine of cat (Fig. 16)	162
Small intestine of rabbit (Fig. 17)	163
— intestine of rabbit (Fig. 18)	163
— intestine of rabbit (Fig. 19)	163
— intestine of rabbit (Fig. 20)	164
— intestine of rabbit (Fig. 21)	
Concentration-action curve type I (Fig. 1)	171
—— curve type II (Fig. 2)	170
Apparatus for measuring the influence of drugs on the activity of animals	
	172
	174
of dog during night after a large dose of coffee (containing about 200	
J 1 1 0 1	174
	202
— fluorid on growth of female rats (Fig. 2)	
—— fluorid on growth of male rats (Fig. 3)	
— fluorid on food consumption (Fig. 4)	
— fluorid on food consumption (Fig. 5)	
Choice of fluorid and unpoisoned food (Fig. 6)	
Sodium chlorid on growth of female rats (Fig. 7)	
	211
	211
1	214
rock on growth of female rats (Fig. 11)	
,	216
Choice of phosphate rock and unpoisoned food (Fig. 13)	
Calcium phosphate on growth of male rats (Fig. 14)	
	220
— phosphate on food consumption (Fig. 16)	
carbonate on growth of male rats (Fig. 17)	
carbonate on growth of female rats (Fig. 18)	
carbonate on food consumption (Fig. 19)	
Intestine tracings Rloads from cat 468 (Fig. 1)	231

ILLUSTRATIONS	vii
Intestine tracings. Bloods from cat 468 (Fig. 2)	. 231
— tracings. Bloods from cat 469 (Fig. 3)	
— tracings. Bloods from cat 472 (Fig. 6)	
— tracings. Bloods from cat 472 (Fig. 7)	
— tracings. Bloods from cat 471 (Fig. 9)	
Toxicity of thymol and carvacrol (Fig. 1)	
— of thymol and carvacrol (Fig. 2)	
Microphotograph from a fresh frozen section of the kidney of the norma	
control animal of experiment 2, table 1 (Fig. 1)	
from a fresh frozen section of the kidney of the normal control anim	
of experiment 5, table 1 (Fig. 2)	
—from a fresh frozen section of the kidney of the animal of experiment	
table 1 (Fig. 3)	
table 1 (Fig. 4)	
Colored microphotograph from a fresh frozen section of the kidney of the	
control naturally nephropathic animal of experiment 1, table 2 (Fig.	
— microphotograph from a fresh frozen section of the naturally nephr	
pathic animal of experiment 10 (table 2) (Fig. 6)	
Effect of benzyl benzoate on leucocytes (Charts I and II)	
— of benzyl benzoate on leucocytes (Chart III)	
— of benzyl benzoate on leucocytes (Chart IV)	
— of benzyl benzoate on leucocytes (Chart V)	
of benzyl benzoate on leucocytes (Chart V)	
Highest tolerated doses of ethylhydrocuprein hydrochlorid per kilogram	
white rat (Chart 1)	
per kilogram of white rat (Chart 2)	
tolerated doses of mercurophen per kilogram of white rat (Chart 3)	
tolerated doses of solutions of disodium arsphenamin per kilogram	
white rat (Chart 4)	
tolerated doses of neoarsphenamin per kilogram of white rat (Chart 5).	
tolerated doses of ethylhydrocuprein hydrochlorid by intravenou	
injection (Chart 6)	
tolerated doses of ethylhydrocuprein hydrochlorid by subcutaneous	
injection (Chart 7)	
Effects of morphin on untreated turtle heart, of chelidonin after morphin an	
of cotarnin hydrochlorid (stypticin) after chelidonin on the same hea	
(Fig. 1)	
of chelidonin on frog's heart (Fig. 2)	<del>1110</del> 459
or energonin on mor story success (Fig. 2)	452 453

## ILLUSTRATIONS

Effects of hydrastinin hydrochlorid after hydrastin hydrochlorid on atro-	
pinized frog's heart (Fig. 4)	454
— of narcotin hydrochlorid on atropinized frog's heart; antagonism by	
morphin hydrochlorid (Fig. 5)	455
of chelidonin sulphate on untreated turtles, heart of morphin hydro-	
chlorid after chelidonin and of cotarnin hydrochlorid (stypticin) after	
morphin (Fig. 6)	457
— of morphin hydrochlorid after hydrastin hydrochlorid on atropinized	
frog's heart (Fig. 7)	459
of hydrastin on turtle's heart and antagonism by cotarnin hydrochlorid	
(stypticin) (Fig. 8)	460
— of different concentrations of chelidonin sulphate in Ringer's solution	
on frog's gastrocnemius muscle (Fig. 9)	463
Tadpoles after four doses ranging from 0.1 mgm. to 50 mgm. of standard	
desiccated cattle thyroid (Fig. 1)	477
after four 20 mgm. doses of desiccated thyroid (Fig. 2)	477
—— fed with 100 mgm. doses of dried blood every other day from June 13 to	
July 10 (Fig. 3)	478
Depression of the hemoglobin curve after the injection in ten to eleven min-	
utes of 25 cc. isotonic saline per kilogram in eight normal dogs (Fig. 1)	483
— of the hemoglobin curve in two dogs in which the liver has been removed	
	484
— of the hemoglobin curve in two normal dogs after the injection in ten	
to eleven minutes of 25 cc. isotonic saline per kilogram to which 0.9 mgm.	
per kilogram of epinephrin has been added (Fig. 3)	485
Twenty cubic centimeters per kilogram of isotonic saline plus 0.9 mgm. of	
epinephrin were injected into two dogs after the removal of the liver	
from the circulation (Fig. 4)	487
Twenty-five cubic centimeters per kilogram of isotonic saline was injected	
into three rabbits showing the depression of the hemoglobin curve	
(Fig. 5)	488
— cubic centimeters of isotonic saline plus 0.9 mgm. of epinephrin per	
kilogram was injected into three rabbits (Fig. 6)	489
Diagram showing the portal and systemic circulation (Fig. 7)	491
If the liver is removed from the circulation and the portal blood shunted	
around it by means of an Eck fistula it will be seen that there is no ob-	
struction to the venous return in the entire body (Fig. 8)	<b>49</b> 2