

JPET # 262063

Title Page

Relative Selectivity of Covalent Inhibitors Requires Assessment of Inactivation Kinetics and Cellular Occupancy: A Case Study of Ibrutinib and Acalabrutinib

Melissa Hopper, Tarikere Gururaja, Taisei Kinoshita, James P. Dean, Ronald J. Hill,
Ann Mongan

*Pharmacyclics LLC, an AbbVie Company, Sunnyvale, CA, USA (M.H., T.G., T.K.,
J.P.D., R.J.H., A.M.)*

Target journal: *The Journal of Pharmacology and Experimental Therapeutics*

Keywords: kinase selectivity, ibrutinib, acalabrutinib

JPET # 262063

Running Title Page

Running title (53 characters; ≤ 60 characters, including spaces): BTK vs TEC

selectivity and relative kinase inhibition

Corresponding author:

Ann Mongan

Pharmacyclics, LLC

999 E. Arques Avenue

Sunnyvale, CA, USA

Phone: (408) 774-0330

Fax: (408) 774-0340

amongan@pcyc.com

Text pages: #

Tables: 3

Figures: 3

References: 14

Abstract (≤ 250 words): 249

Introduction (≤ 750 words): 696

Discussion (≤ 1500 words): 796

JPET # 262063

List of nonstandard abbreviations: BTK, Bruton's tyrosine kinase; CLL, chronic lymphocytic leukemia; DMSO, dimethyl sulfoxide; IC₅₀, half maximal inhibitory concentration; IMAP, immobilized metal ion affinity-based fluorescence polarization; PBMC, peripheral blood mononuclear cells; PBS, phosphate buffered saline.

Recommended section: Drug Discovery and Translational Medicine

Abstract (250-word limit; single, unformatted paragraph)

Kinases form an attractive class of targets for small molecule inhibitors, but similarity among their adenosine triphosphate binding sites presents difficulties for developing selective drugs. Standard methods of evaluating selectivity of most reversibly bound drugs account for binding affinity, but not the two-step process, affinity and inactivation, occurring during covalent inhibition. To illustrate this concept, we assessed the selectivity of Bruton's tyrosine kinase (BTK) over TEC kinases by two novel therapeutics: ibrutinib and acalabrutinib. The two-step process and time-dependent inhibition unique to covalent inhibitors were evaluated with two biochemical assays measuring enzymatic function and inhibition kinetics. The selectivity for BTK over TEC found in these biochemical analyses was 1 -1.5 for ibrutinib and 3.0 - 4.2 for acalabrutinib. To further assess drug selectivity in a more physiologically-relevant context, we developed cell-based occupancy assays that quantify the percentage of drug-inactivated kinases. Cellular selectivity of BTK over TEC was determined after MWCL-1 cells and samples from patients with chronic lymphocytic leukemia (CLL) were treated for durations and concentrations based on human pharmacokinetics of each

JPET # 262063

drug. In MWCL-1 cells, BTK/TEC selectivities measured at 0.5, 1, and 3 hours were 2.53, 1.05, and 1.51 for ibrutinib and 0.97, 1.13, and 2.56 for acalabrutinib, respectively. The equivalent selectivity measured in samples from patients with CLL were 1.31 ± 0.27 and 1.09 ± 0.11 for ibrutinib and acalabrutinib, respectively. Collectively, our data show that when properly accounting for time-dependent factors and relevant cellular context, ibrutinib and acalabrutinib demonstrate similar selectivity for BTK over TEC.

Significance Statement (max: 80 words/currently: 80 words)

This study shows relative selectivity of covalent inhibitors toward different kinase targets should be assessed with both affinity and inactivation kinetics to accurately account for time-dependent effects of covalent binding and assessed in a cellular matrix to reproduce the physiologic context of target inhibition. This is illustrated with a case study of ibrutinib and acalabrutinib for which selectivity assessment with appropriate assays, versus measuring binding affinity with KINOMEscan alone, corroborate emerging clinical data demonstrating similar safety profiles between the therapies.

Introduction (750-word limit)

Bruton's tyrosine kinase (BTK), a member of the TEC family of kinases, mediates B-cell receptor signaling and has emerged over the past decade as an effective clinical target for first-line therapy and for treating relapsed/refractory chronic lymphocytic leukemia (CLL) (Stevenson et al., 2011; Woyach et al., 2012). Ibrutinib is the only once-daily BTK inhibitor that is approved in the United States, Europe, and other countries for the treatment of several B-cell malignancies, including CLL and mantle cell lymphoma (MCL) (Janssen-Cilag International NV, 2018; Pharmacyclics LLC, 2019). Acalabrutinib, another BTK inhibitor taken twice daily, has received conditional approval in the United States only for the treatment of relapsed/refractory MCL and is in development for the treatment of other B-cell malignancies (AstraZeneca Pharmaceuticals LP, 2017). While it has been demonstrated that both ibrutinib and acalabrutinib potently inhibit BTK via covalent bond formation, selectivity for this target over other TEC family kinases has not been rigorously assessed.

In 2017, Barf and colleagues reported the pharmacology of acalabrutinib and highlighted its biochemical and cellular selectivity for BTK over other kinases in the TEC family (Barf et al., 2017). The authors used two different biochemical assays to assess the inhibition selectivity of BTK over TEC for several covalent BTK inhibitors, including ibrutinib and acalabrutinib. In their study, the potency for BTK was measured using the immobilized metal ion affinity-based fluorescence polarization (IMAP) assay and the potency for TEC was measured using a time-resolved fluorescence resonance energy transfer (TR-FRET) assay. The two assays differed in that the IMAP assay informed enzymatic activity of the kinases, whereas the TR-FRET assay measured inhibition via

JPET # 262063

competitive binding and thus only informed affinity. Neither of these assays were optimal for quantifying drug selectivity because they assessed only part of the inactivation mechanism of action. Accordingly, although each assay reported higher potency for both BTK and TEC by ibrutinib relative to acalabrutinib, selectivity between the two kinases could not be inferred from the half maximal inhibitory concentration (IC_{50}) derived from the two different assay platforms. The authors further profiled inhibition via competitive binding of ibrutinib and acalabrutinib against a panel of 395 nonmutant human kinases using the KINOMEscan Assay Platform (Eurofins DiscoverX Corporation, Fremont, CA, USA). This experiment showed that when both drugs were tested at the same concentration, 1 μ M, ibrutinib was more potent against BTK and TEC than acalabrutinib. Notably, physiologically relevant exposures for acalabrutinib are five-fold higher than for ibrutinib (Byrd et al., 2013; Byrd et al., 2016); while these results were consistent with earlier studies (Honigberg et al., 2010; Byrd et al., 2016) in finding ibrutinib as the more potent inhibitor, the arbitrary use of 1 μ M resulted in data that were not informative toward understanding target inhibition of the two drugs at physiologically relevant exposures (Byrd et al., 2013; Byrd et al., 2016).

Kinase inhibition via covalent bond formation is a two-step process that begins with the compound interacting with the kinase driven by affinity (represented by K_i) and ends with an inactivation step (K_{inact}). Accordingly, potency of covalent inhibitors cannot be determined using only traditional IC_{50} measurements because this does not account for the entirety of the two-step process of covalent bond formation (Bauer, 2015). Here, we utilized the ratio of K_{inact}/K_i , which is the preferred metric to rank potency of different covalent inhibitors against a target (Bauer, 2015). In fact, Barf and colleagues discussed

JPET # 262063

the relevance of these two parameters in their article, highlighting the faster inactivation rate for an acrylamide-substituted acalabrutinib and higher target affinity by ibrutinib as the primary factors contributing to the difference in potency of each compound (Barf et al., 2017). Furthermore, both the IMAP and the TR-FRET platforms are biochemical assays that measure the enzymatic reactions in relatively artificial systems that do not account for the complexity of the cellular environment, which has high (millimolar) adenosine triphosphate (ATP) concentrations and hundreds of protein interactions. Collectively, it is clear that selectivity can be accurately assessed only in a comprehensive manner, including analysis of binding kinetics, enzymatic activity, and cellular occupancy at physiologically-relevant drug concentrations. In the present study, we assessed these three parameters to rigorously evaluate the selectivity of ibrutinib and acalabrutinib for BTK and TEC.

Materials and Methods (no word limit)

Biochemical Assay – Enzymatic Function

Biochemical enzymatic IC₅₀ data were generated by Nanosyn (Santa Clara, CA, USA). Determination of inhibitor potency against BTK enzyme was carried out using the microfluidic-based LabChip 3000 Drug Discovery System (Caliper Life Sciences, Hopkinton, MA, USA), which uses capillary electrophoresis to separate phosphorylated and nonphosphorylated peptides. Briefly, the enzyme reaction was started by preincubating inhibitor at 12 different concentrations in dimethyl sulfoxide (DMSO; 1000, 330, 110, 37, 12, 4.1, 1.4, 0.46, 0.15, 0.051, 0.017, and 0.006 nM) with purified BTK for 15 minutes in a buffer system consisting of 100 mM HEPES solution, pH 7.5, 0.1%

JPET # 262063

bovine serum albumin, 0.01% Triton X-100 (Sigma-Aldrich, St. Louis, MO, USA), 1 mM dithiothreitol, and 5 mM magnesium chloride at 25°C. After enzyme-inhibitor preincubation, the kinase reaction was initiated by adding peptide substrate and ATP (50 µM), prepared in the same assay buffer, followed by further incubation of reaction mixture for 3 hours. At the end of the incubation, the reaction was quenched by a buffer containing 50 mM of EDTA. Appropriate control samples (0% inhibition in the absence of inhibitor, DMSO only, and 100% inhibition, in the absence of enzyme) were assembled in replicates of four and used to calculate the percentage of inhibition in the presence of compounds. IC₅₀ values were determined by fitting the inhibition curves using a four-parameter sigmoidal dose-response model using XLfit 4 software (IDBS, Boston, MA, USA). A similar protocol was used to determine the IC₅₀ values for TEC kinase, with minor modifications to optimize for enzyme, peptide, and ATP concentrations.

Biochemical Assay – Kinetics of Target Inhibition

For kinetic studies, similar enzyme assay conditions were employed as described in the Biochemical Assay – Enzymatic Function section. The generation of progress curves for BTK/TEC peptide phosphorylation in the presence of test inhibitor was performed at 11 drug concentrations (10, 8.3, 6.9, 5.8, 4.8, 4.0, 3.4, 2.8, 2.3, 1.9, and 1.6 nM). After initiating the enzyme reaction, the real-time curves were obtained for a total period of 5 hours using a climate-controlled LabChip 3000 Drug Discovery System. Progress curves were then fitted by XLfit 4 software using the time-dependent inhibition equation $[P] = V_s \times t + ((V_i - V_s)/K_{obs}) \times (1 - \exp(-K_{obs} \times t))$, where V_i is the initial velocity, V_s is

JPET # 262063

the steady-state velocity, and K_{obs} represents the rate of inactivation. For time-dependent inhibitors, the obtained K_{obs} values were plotted against the compound concentration using either a hyperbolic or linear fit. The model with the better goodness of fit was selected. These plots enabled determination of K_{inact} and K_i values. The acquisition of all progress curve data points and the calculation of kinetic parameters were performed by Nanosyn.

Cell Treatments

The MWCL-1 cell line, which was established from a patient with Waldenström macroglobulinemia (Hodge et al., 2011), was obtained from Mayo Foundation for Medical Education and Research (Rochester, MN, USA). MWCL-1 treatments were performed in six-well tissue culture plates using 6×10^6 cells per well at a concentration of 2×10^6 cells/mL in Gibco RPMI 1640 (ThermoFisher Scientific, Waltham, MA, USA) supplemented with Gibco 1X Penicillin-Streptomycin (ThermoFisher Scientific), 10% heat-inactivated fetal bovine serum, and 1 mM Gibco Sodium Pyruvate (ThermoFisher Scientific). Cells were treated with inhibitors for 0.5, 1, or 3 hours at 37°C and 5% CO₂, with a final DMSO concentration of 0.1% (v/v). Following treatment, cells were washed two times with 1X phosphate-buffered saline (PBS) to remove excess compound. Cell pellets were lysed in 1X PBS with 0.1% NP-40 detergent and 1X Protease Inhibitor Cocktail (Sigma-Aldrich). Cell lysate protein concentrations were quantified by Pierce BCA Protein Assay (ThermoFisher Scientific) per manufacturer's instruction using a FlexStation 3 Multi-Mode Microplate Reader (Molecular Devices, San Jose, CA, USA). Treatment-free cryopreserved CLL peripheral blood mononuclear cells (PBMC) isolated

JPET # 262063

from four donors were obtained from Discovery Life Sciences (Huntsville, AL, USA). Samples were thawed, washed once with Gibco RPMI 1640 supplemented with 10% heat-inactivated fetal-bovine serum, and resuspended in fresh medium. Cells were plated in a six-well tissue culture plate using 9×10^6 cells per well at a concentration of 3×10^6 cells/mL and incubated at 37°C with 5% CO₂ for 2 hours prior to compound treatment. Treatments and cell lysis were performed using the same procedure as for MWCL-1 cells.

Enzyme Occupancy Assays

Cell-based BTK and TEC occupancy assays were developed to evaluate drug selectivity between the two kinases in a more physiologically-relevant system. An MSD Small Spot High Bind plate (Meso Scale Diagnostics, Rockville, MD, USA) was coated with 35 µL of 10 µg/mL anti-TEC antibody (Sigma-Aldrich) or 35 µL of 1 µg/mL anti-BTK antibody and incubated overnight at 4°C. The following day, the plate was warmed to room temperature, washed three times with 200 µL of 1X MSD Tris Wash Buffer (Meso Scale Diagnostics), and blocked with 200 µL of 3% MSD Blocker A (Meso Scale Diagnostics) for 1 hour. In a 96-well plate, 25 µL of 2 mg/mL cell lysate was incubated with 25 µL of 50 nM proprietary biotinylated probe for 1 hour to allow the probe to bind TEC or BTK kinase not occupied by drug. The MSD plate was then washed three times, and 45 µL of probe-labeled lysate was added to each well and incubated for 2 hours. The plate was then washed three times prior to the addition of 45 µL of MSD SULFO-TAG Streptavidin (Meso Scale Diagnostics) diluted 1:500 in 1X PBS. After 1 hour of incubation, 150 µL of 1X MSD Read Buffer (Meso Scale Diagnostics) was added to

each well and the plate was read immediately on an MSD Sector S 600 plate reader (Meso Scale Diagnostics). Unless otherwise specified, incubation steps were performed at room temperature with 300 rpm shaking.

Results (no word limit)

Inhibition Kinetics of BTK and TEC by Ibrutinib and Acalabrutinib

Inhibition of kinases by covalent binding is a two-step process that is driven by affinity with the target, followed by inactivation by covalent bond formation. Therefore, the assessment of relative selectivity toward different targets for covalent inhibitors should be evaluated using measurement of both binding affinity and time-dependent inactivation (Strelow, 2017). In the current study, biochemical analyses of enzymatic function were performed and kinetics measuring both of these parameters were derived using the LabChip 3000 Drug Discovery System. Biochemical IC₅₀ values for kinase inhibition demonstrated BTK and TEC selectivity ratios of 1.0-fold for ibrutinib and 4.2-fold for acalabrutinib (Table 1). These data demonstrated similar binding affinity of ibrutinib for both BTK and TEC kinase, as indicated by their similar K_i values: 0.95 ± 0.009 nM (BTK) and 1.8 ± 0.05 nM (TEC) (Table 1). Examples of the enzymatic progression curves and hyperbolic fit curves that were used to calculate the kinetic parameters in Table 1 (K_i and K_{inact}) as well as the equations used for the calculations are shown in Figure 1. The complete set of kinetic data for all determinations are included as supplemental material (Supplemental Tables 1–8; Supplemental Figures 1–8). The rates of enzymatic inactivation (K_{inact}) of ibrutinib against both BTK and TEC were also similar: 0.011 ± 0.0004 (BTK) and 0.013 ± 0.0005 (TEC). In contrast,

JPET # 262063

acalabrutinib demonstrated 20-fold higher binding affinity for BTK (8.7 ± 0.5) compared to TEC (160 ± 37), but a five-fold faster rate of TEC inactivation (0.012 ± 0.001) compared to BTK inactivation (0.0021 ± 0.0004). Therefore, despite these large differences in binding and inactivation, selectivity for BTK over TEC, as measured by the K_{inact}/K_i ratio with this method, were within two-fold for the two drugs: a selectivity ratio of 1.5 for ibrutinib and 3 for acalabrutinib (Table 1). This difference in kinetic properties of the two compounds highlights why different approaches are necessary to rank selectivity between reversible and covalent inhibitors. For reversible binding, relative selectivity can be inferred from the IC_{50} of affinity analysis alone because potency is driven by affinity in these cases. For covalent inhibitors, the IC_{50} does not reflect the combined affinity plus inactivation steps. The apparent IC_{50} for covalent inhibitors decreases as a function of time due to the depletion of active enzyme caused by covalent bond formation. Accordingly, this analysis demonstrated how binding affinity measured alone inadequately reflect selectivity among different targets of covalent inhibitors.

Acalabrutinib and Ibrutinib Exhibit Similar Selectivity for BTK Over TEC in MWCL-1 Cells and Human CLL PBMC

MWCL-1 cells were treated with eight different concentrations of either ibrutinib or acalabrutinib for 0.5, 1, and 3 hours (Figure 2). The timepoints were selected to encompass the T_{max} (ie, the time for drug to reach maximal plasma concentration in patients) of both compounds (Advani et al., 2013; Barf et al., 2017). With very short exposure (0.5 hour), ibrutinib showed slightly higher selectivity for BTK over TEC: when

JPET # 262063

50% of BTK was occupied, only 25% of TEC was occupied (Figure 2, left). Acalabrutinib selectivity was not observed after 0.5-hour exposure (ie, there was no separation for BTK and TEC dose-response curves). BTK over TEC selectivity was minimal for both drugs with 1-hour exposure (Figure 2, center) across eight concentrations. With 3 hours of treatment, there was modest selectivity indicated by the small separation between the BTK and TEC dose-response curves for both drugs (Figure 2, right), and this selectivity was nearly equivalent for both drugs. In effect, these cellular results are consistent with the biochemical results; ibrutinib exhibited slightly higher potency on BTK over TEC at 0.5 hours, but similar potency against the two kinases became apparent with longer treatment time (Table 2). The time-dependent effect of kinase inhibition observed here is not unlike the physiological activities of these drugs in vivo. Because the drug-binding reaction is irreversible, the amount of drug-bound targets increases with longer duration of exposure. The current data with MWCL-1 cells show that the two drugs have nearly equivalent selectivity between BTK and TEC, and the ranges of selectivity ratios (BTK/TEC) were similarly small (1.05–2.53 and 0.97–2.56 for ibrutinib and acalabrutinib, respectively) throughout the time window of 0.5 to 3 hours (Table 2) that is required to achieve maximum occupancy resulting in target inhibition (Barf et al., 2017). Protein content and cellular compositions of PBMC from patients with CLL may differ from the MWCL-1 cells, so we further evaluated the differential selectivity of the two drugs in cryopreserved PBMC isolated from previously untreated patients with CLL. Cellular occupancy assays, which measure the percentage of target kinases bound by drugs, were conducted after each patient sample was treated with either ibrutinib or acalabrutinib for 3 hours (Figure 3). Despite the potential biological variation among

JPET # 262063

different donors, IC₅₀ values against BTK and TEC were highly similar for both drugs across four individual patients (Table 3). Ibrutinib was more potent toward both kinases, with IC₅₀ for BTK ranging from 1.3 to 1.5 nM and TEC ranging from 1.3 to 2.1 nM. Acalabrutinib was approximately 10-fold less potent against both kinases, with IC₅₀ for BTK ranging from 7.9 to 11.9 nM and TEC ranging from 10.2 to 11.5 nM. Results of this occupancy assay are consistent with findings from biochemical analyses from the present study and others' work (Bye et al., 2015); ibrutinib is generally a more potent compound than acalabrutinib toward members of the TEC kinase family. However, as observed in MWCL-1 cells, the two drugs' BTK/TEC selectivity ratio observed in samples from patients with CLL was remarkably similar: 1.32 ± 0.27 for ibrutinib and 1.09 ± 0.11 for acalabrutinib (Table 3). These data indicate that the two compounds exerted equivalent selectivity for BTK over TEC in a cellular environment despite having the higher potency of ibrutinib against BTK and TEC kinases based on IC₅₀ values (Table 3, Figure 4). In general, the implication of higher potency is that a drug is more reactive toward its targets and less drug is needed to achieve pharmacologic activity, whereas selectivity is a measure of the ratio of on-target/off-target effects, such as BTK over TEC. Here, the collective data indicate that ibrutinib is a more potent drug than acalabrutinib toward its targets, but acalabrutinib does not offer improved on-target/off target selectivity.

Discussion (1500-word limit)

Kinases form an attractive target class for small molecule inhibitors. Specifically, targeting BTK has resulted in therapies for numerous hematologic malignancies that previously had substantial unmet medical needs. However, the high similarity of the

JPET # 262063

ATP binding site on protein kinases, a typical target of inhibition, mounts a continuous challenge for developing selective small molecule inhibitors for this target class. Our understanding of the selectivity of covalent and noncovalent inhibitors is highly dependent on the assessment methods, which have different implications and interpretations for irreversible and reversible binding modes. Crucial to this is the understanding that covalent kinase inhibition is a two-step process involving initial binding (driven by affinity) and time-dependent inactivation (driven by covalent bond formation). Thus, in order to evaluate biochemical potency and understand target selectivity of covalent inhibitors, both steps must be evaluated.

Published pharmacokinetics for ibrutinib and acalabrutinib report that the two compounds reach peak plasma concentrations between 1 and 2 hours and between 0.6 and 1.1 hours after dosing, respectively (Advani et al., 2013; Byrd et al., 2016). In a recent PK/PD modeling study of a covalent BTK inhibitor, it was shown that target occupancy in B cells and drug plasma exposure reach peak levels at the same time points (Daryaee et al., 2017). C_{\max} for ibrutinib at 420 mg once daily (310 nM) was significantly lower than acalabrutinib at 100 mg twice daily (1.78 μ M) (Byrd et al., 2016; Chen et al., 2018). These data show that direct comparison between the two drugs using in vitro systems should be conducted at concentrations consistent with this five-fold or greater difference in human exposure levels. Without such considerations, the physiological effects of these agents cannot be reasonably discerned. In one example of such oversight, Bye and colleagues argued that acalabrutinib is a less potent inhibitor of TEC phosphorylation based on in vitro analysis using equal concentration of either drug, but subsequently discovered that therapeutic doses of acalabrutinib did inhibit

JPET # 262063

TEC based on their own findings from patient samples (Bye et al., 2015; Bye et al., 2017). These authors concluded that the apparent potency of acalabrutinib toward both BTK and TEC in their in vitro experiments was lower than that observed in patients, as blood samples derived from patients treated with ibrutinib and acalabrutinib exhibited very similar platelet aggregation responses. These inconsistencies highlight the importance of adjusting in vitro exposure according to a drug's therapeutic concentration to obtain experimental results that are informative toward understanding the drugs' effects in humans.

The recent high interest in understanding kinase selectivity profiles of various BTK inhibitors stems from a desire to optimize patient outcomes with treatment, minimize adverse events, and understand how off-target effects may be related to such events. For example, bleeding is an adverse event observed in patients treated with BTK inhibitors, with most cases being grade 1 or 2 in severity (Mock et al., 2018). While clinical data were still being accumulated, an earlier hypothesis based on kinase affinity data proposed that acalabrutinib might lead to lower bleeding rates due to its higher selectivity for BTK relative to TEC (Byrd et al., 2016; Barf et al., 2017; Awan et al., 2019). However, when properly adjusted for clinical exposure, covalent binding mechanism, and enzymatic activity, our data and those of other groups (Bye et al., 2017) now indicate that ibrutinib and acalabrutinib have similar selectivity between BTK and TEC. This similar selectivity is consistent with a recent study suggesting that platelet inhibition and potential hemorrhagic risk, predicted by in vitro closure time, is likely a class effect across five BTK inhibitors: ibrutinib, BGB-3111, acalabrutinib, ONO/GS-4059, and evobrutinib (Denzinger et al., 2019). Furthermore, recent

JPET # 262063

mechanistic investigations of the effects of ibrutinib and acalabrutinib on platelet functions suggested that polymorphism of drug efflux pumps might sensitize some patients toward platelet aggregation impairment more than others (Series et al., 2019). Because polymorphism in drug efflux pumps is an intrinsic factor, platelet aggregation did not improve when the authors re-treated the same patient samples with acalabrutinib. Emerging clinical data now increasingly indicate that the bleeding rates of ibrutinib and acalabrutinib are similar, with more bleeding events reported for patients treated with acalabrutinib: 60% of 99 patients treated with first-line acalabrutinib reporting bleeding events of all grades and two of these patients showed grade 3 events after longer follow-up (median time on study of 33 months) (Byrd et al., 2018). In summary, our study applies fundamental biochemistry principles to demonstrate how standard methods used to evaluate target selectivity for reversible inhibitors fail to fully characterize irreversible inhibitor selectivity. The current case study of the two covalent inhibitors ibrutinib and acalabrutinib highlights how these results, when performed at physiological concentrations and in a more relevant cellular context, better inform the understanding of possible off-target clinical observations of bleeding events with these two drugs.

Acknowledgments

We thank Sany Hoxha for generation of figures and Jo Bairzin, PhD, for medical writing, which was supported by Pharmacyclics LLC, an AbbVie Company.

Authorship Contributions

JPET # 262063

Participated in research design: Mongan, Hill, Gururaja, Kinoshita

Conducted experiments: Hopper, Gururaja

Contributed new reagents or analytic tools: Hopper

Performed data analysis: Hopper, Mongan, Hill, Gururaja

Wrote or contributed to the writing of the manuscript: All authors

References

- Advani RH, Buggy JJ, Sharman JP, Smith SM, Boyd TE, Grant B, Kolibaba KS, Furman RR, Rodriguez S, Chang BY, Sukbuntherng J, Izumi R, Hamdy A, Hedrick E and Fowler NH (2013) Bruton tyrosine kinase inhibitor ibrutinib (PCI-32765) has significant activity in patients with relapsed/refractory B-cell malignancies. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* **31**:88-94.
- Awan FT, Schuh A, Brown JR, Furman RR, Pagel JM, Hillmen P, Stephens DM, Woyach J, Bibikova E, Charuworn P, Frigault MM, Hamdy A, Izumi R, Linghu B, Patel P, Wang MH and Byrd JC (2019) Acalabrutinib monotherapy in patients with chronic lymphocytic leukemia who are intolerant to ibrutinib. *Blood Adv* **3**:1553-1562.
- Barf T, Covey T, Izumi R, van de Kar B, Gulrajani M, van Lith B, van Hoek M, de Zwart E, Mittag D, Demont D, Verkaik S, Krantz F, Pearson PG, Ulrich R and Kaptein A (2017) Acalabrutinib (ACP-196): A Covalent Bruton Tyrosine Kinase Inhibitor with a Differentiated Selectivity and In Vivo Potency Profile. *J Pharmacol Exp Ther* **363**:240-252.
- Bauer RA (2015) Covalent inhibitors in drug discovery: from accidental discoveries to avoided liabilities and designed therapies. *Drug discovery today* **20**:1061-1073.
- Bye AP, Unsworth AJ, Desborough MJ, Hildyard CAT, Appleby N, Bruce D, Kriek N, Nock SH, Sage T, Hughes CE and Gibbins JM (2017) Severe platelet dysfunction in NHL patients receiving ibrutinib is absent in patients receiving acalabrutinib. *Blood Adv* **1**:2610-2623.
- Bye AP, Unsworth AJ, Vaiyapuri S, Stainer AR, Fry MJ and Gibbins JM (2015) Ibrutinib Inhibits Platelet Integrin α IIb β 3 Outside-In Signaling and Thrombus Stability But Not Adhesion to Collagen. *Arterioscler Thromb Vasc Biol* **35**:2326-2335.
- Byrd JC, Furman RR, Coutre SE, Flinn IW, Burger JA, Blum KA, Grant B, Sharman JP, Coleman M, Wierda WG, Jones JA, Zhao W, Heerema NA, Johnson AJ, Sukbuntherng J, Chang BY, Clow F, Hedrick E, Buggy JJ, James DF and O'Brien S (2013) Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. *The New England journal of medicine* **369**:32-42.
- Byrd JC, Harrington B, O'Brien S, Jones JA, Schuh A, Devereux S, Chaves J, Wierda WG, Awan FT, Brown JR, Hillmen P, Stephens DM, Ghia P, Barrientos JC, Pagel JM, Woyach J, Johnson D, Huang J, Wang X, Kaptein A, Lannutti BJ, Covey T, Fardis M, McGreivoy J, Hamdy A, Rothbaum W, Izumi R, Diacovo TG, Johnson AJ and Furman RR (2016) Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia. *The New England journal of medicine* **374**:323-332.
- Byrd JC, Woyach JA, Furman RR, Martin P, Brien SM, Brown JR, Stephens DM, Barrientos JC, Devereux S, Hillmen P, Pagel JM, Chen D-Y, Hamdy A, Izumi R, Patel P, Wang MH, Jain N and Wierda WG (2018) Acalabrutinib in Treatment-Naive (TN) Chronic Lymphocytic Leukemia (CLL): Updated Results from the Phase 1/2 ACE-CL-001 Study. *Blood* **132**:692.
- Chen J, Kinoshita T, Gururaja T, Sukbuntherng J, James D, Lu D, Whang J, Versele M and Chang BY (2018) The effect of Bruton's tyrosine kinase (BTK) inhibitors on collagen-induced platelet aggregation, BTK, and tyrosine kinase expressed in hepatocellular carcinoma (TEC). *European journal of haematology*.

JPET # 262063

- Daryaee F, Zhang Z, Gogarty KR, Li Y, Merino J, Fisher SL and Tonge PJ (2017) A quantitative mechanistic PK/PD model directly connects Btk target engagement and in vivo efficacy. *Chem Sci* **8**:3434-3443.
- Denzinger V, Busygina K, Jamasbi J, Pekrul I, Spannagl M, Weber C, Lorenz R and Siess W (2019) Optimizing Platelet GPVI Inhibition versus Haemostatic Impairment by the Btk Inhibitors Ibrutinib, Acalabrutinib, ONO/GS-4059, BGB-3111 and Evobrutinib. *Thromb Haemost* **119**:397-406.
- Hodge LS, Novak AJ, Grote DM, Braggio E, Ketterling RP, Manske MK, Price Troska TL, Ziesmer SC, Fonseca R, Witzig TE, Morice WG, Gertz MA and Ansell SM (2011) Establishment and characterization of a novel Waldenstrom macroglobulinemia cell line, MWCL-1. *Blood* **117**:e190-197.
- Honigberg LA, Smith AM, Sirisawad M, Verner E, Loury D, Chang B, Li S, Pan Z, Thamm DH, Miller RA and Buggy JJ (2010) The Bruton tyrosine kinase inhibitor PCI-32765 blocks B-cell activation and is efficacious in models of autoimmune disease and B-cell malignancy. *Proceedings of the National Academy of Sciences of the United States of America* **107**:13075-13080.
- Janssen-Cilag International NV (2018) IMBRUVICA (ibrutinib) Summary of Product Characteristics., in, Beerse, Belgium.
- LP AP (2017) CALQUENCE® (acalabrutinib) capsules, for oral use, in, AstraZeneca Pharmaceuticals LP,, Wilmington, DE.
- Mock J, Kunk PR, Palkimas S, Sen JM, Devitt M, Horton B, Portell CA, Williams ME and Maitland H (2018) Risk of Major Bleeding with Ibrutinib. *Clinical lymphoma, myeloma & leukemia* **18**:755-761.
- Pharmacyclics LLC (2019) IMBRUVICA® (ibrutinib) prescribing information,, in, Sunnyvale, CA.
- Series J, Garcia C, Levade M, Viaud J, Sie P, Ysebaert L and Payrastre B (2019) Differences and similarities in ibrutinib and acalabrutinib effects on platelet functions. *Haematologica*.
- Stevenson FK, Krysov S, Davies AJ, Steele AJ and Packham G (2011) B-cell receptor signaling in chronic lymphocytic leukemia. *Blood* **118**:4313-4320.
- Strelow JM (2017) A Perspective on the Kinetics of Covalent and Irreversible Inhibition. *SLAS discovery : advancing life sciences R & D* **22**:3-20.
- Woyach JA, Johnson AJ and Byrd JC (2012) The B-cell receptor signaling pathway as a therapeutic target in CLL. *Blood* **120**:1175-1184.

JPET # 262063

Footnotes

Financial support for this work was provided by Pharmacyclics LLC, an AbbVie Company.

This work was previously presented by M. Hopper et al at the 60th ASH Annual Meeting and Exposition on December 1-4, 2018 (poster 3498).

Conflict of Interest Disclosures: All authors were employed by Pharmacyclics LLC, an AbbVie Company, at the time of study analyses, and have stock ownership in AbbVie. JPD was employed by and has stock ownership in CTI BioPharma Corp. RJH has stock ownership in Principia Biopharma.

Role of the Funder/Sponsor: Pharmacyclics LLC, an AbbVie Company, sponsored and designed the study. Study investigators and their research teams collected the data. The sponsor confirmed data accuracy and performed analysis of the data. Medical writing support was funded by the sponsor.

Figure legends.

Fig. 1 Determination of BTK and TEC inactivation kinetic parameters for (a) ibrutinib and (b) acalabrutinib based upon time-dependent inhibition via reaction progression curves. k_{obs} (pseudo-first-order rate constant) was determined by the following equation: $k_{obs} = k_{-2} + k_2[I]/([I] + K_i^*(1+[S]/K_m))$, where K_i^* is the stable complex forming constant and K_i is the overall final inhibitory constant (for covalent irreversible inhibitors, $k_2 = k_{inact}$ and $k_{-2} = 0$). Upon plotting concentration of inhibitor $[I]$ versus k_{obs} , goodness of fit parameters for the two drugs were determined using either hyperbolic or linear fit from the time-dependent inhibition equation: $[P] = V_s t + ((V_i - V_s)/K_{obs}) * (1 - \exp(-K_{obs} * t))$, where V_i represents initial velocity and V_s is final steady-state velocity (for covalent irreversible inhibitors, $V_s = 0$). From the goodness of fit and statistical analysis data it is clear that the two drugs exhibited a hyperbolic fit (curve fitting software XLfit4 yielded a goodness of fit with lower χ^2 values and F-Test greater than 0.95 in all the cases) indicating that the binding of the two drugs follows a two-step mechanism. The first step includes reversible binding of the inhibitor to the enzyme followed by the second step of covalent bond formation for both BTK and TEC enzymes.

Fig. 2. Dose-response curves of BTK and TEC occupancy in the MWCL-1 cell line. BTK and TEC occupancy (closed and open circles) is shown for ibrutinib (blue lines) and acalabrutinib (red lines). Occupancy is shown for drug exposure at 0.5 hours (left), 1 hour (center), and 3 hours (right).

JPET # 262063

Fig. 3. Dose-response curves of BTK and TEC occupancy in human CLL PBMC.

Occupancy of BTK (blue lines) and TEC (red lines) is shown for human CLL PMBC treated with ibrutinib (left) and acalabrutinib (right). Occupancy was assessed at 3 hours of exposure to ibrutinib or acalabrutinib.

Fig 4. Apparent selectivity of ibrutinib and acalabrutinib. Binding affinity data are reported by Barf and colleagues (Barf et al., 2017). Enzyme function assay data are the IC₅₀ ratios of TEC/BTK and inactivation kinetics data are the K_{inact}/K_i ratios for ibrutinib (blue) and acalabrutinib (red). Cellular occupancy data are mean TEC/BTK selectivity measured by IC₅₀ ratio for human CLL PMBC.

Tables

Table 1. BTK and TEC selectivity based on biochemical functional and kinetic analyses

Drug	IC ₅₀ (nM) ± SD		IC ₅₀ ratio	K _i (nM) ± SD		K _{inact} (s ⁻¹) ± SD		K _{inact} /K _i (nM ⁻¹ *s ⁻¹) ± SD		Ratio of K _{inact} /K _i
	BTK	TEC	TEC/ BTK	BTK	TEC	BTK	TEC	BTK	TEC	BTK TEC
Ibrutinib	0.24 ± 0.2	0.24 ± 0.2	1	0.95 ± 0.009	1.8 ± 0.05	0.011 ± 0.0004	0.013 ± 0.0005	0.012 ± 0.0006	0.008 ± 0.0004	1.5
Acalabrutinib	2.3 ± 1.6	9.7 ± 2.6	4.2	8.7 ± 0.5	160 ± 37	0.0021 ± 0.0004	0.012 ± 0.001	0.00024 ± 0.00003	0.00008 ± 0.00001	3.0

JPET # 262063

Table 2. IC₅₀ derived from dose-response curves of BTK and TEC occupancy in the MWCL-1 cell line. *BTK/TEC selectivity is defined as TEC IC₅₀/BTK IC₅₀

Drug	Time Point	BTK IC ₅₀ (nM)	TEC IC ₅₀ (nM)	BTK/TEC selectivity*
Ibrutinib	0.5 h	2.88	7.28	2.53
	1 h	3.99	4.18	1.05
	3 h	1.06	1.60	1.51
Acalabrutinib	0.5 h	50.89	49.36	0.97
	1 h	26.10	29.38	1.13
	3 h	5.47	14.01	2.56

JPET # 262063

Table 3. IC₅₀ derived from dose-response curves of BTK and TEC occupancy in human CLL PBMC. *BTK/TEC selectivity is defined as TEC IC₅₀/BTK IC₅₀

Drug	PBMC Donor	BTK IC ₅₀ (nM)	TEC IC ₅₀ (nM)	BTK/TEC* Selectivity	Mean ± SD
Ibrutinib	Donor 1	1.39	1.96	1.41	1.32 ± 0.27
	Donor 2	1.45	2.12	1.46	
	Donor 3	1.30	1.29	0.99	
	Donor 4	1.37	1.93	1.41	
Acalabrutinib	Donor 1	7.86	10.25	1.30	1.09 ± 0.11
	Donor 2	11.93	11.53	0.97	
	Donor 3	9.79	10.19	1.04	
	Donor 4	10.84	11.52	1.06	

FIGURES

Figure 1A




Figure 1B




Figure 2




Figure 3





Figure 4



Supplemental Information.

**Relative Selectivity of Covalent Inhibitors Requires Assessment of Inactivation Kinetics and Cellular Occupancy:
A Case Study of Ibrutinib and Acalabrutinib**


Melissa Hopper, Tarikere Gururaja, Taisei Kinoshita, James P. Dean, Ronald J. Hill, Ann Mongan

Journal of Pharmacology and Experimental Therapeutics

Supplemental Table 1. BTK enzyme kinetics data for ibrutinib (0.35 nM BTK)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K_{obs}, s⁻¹	95%-confidence
Ibrutinib	BTK	0.35	1000	10	0.002622204	0.000360793
Ibrutinib	BTK	0.35	1000	8.333333333	0.002546468	0.000330995
Ibrutinib	BTK	0.35	1000	6.944444444	0.002405458	0.000221326
Ibrutinib	BTK	0.35	1000	5.787037037	0.001744919	0.000127889
Ibrutinib	BTK	0.35	1000	4.822530864	0.00177236	0.000185465
Ibrutinib	BTK	0.35	1000	4.01877572	0.001146117	0.000100137
Ibrutinib	BTK	0.35	1000	3.348979767	0.000984421	4.18082E-05
Ibrutinib	BTK	0.35	1000	2.790816472	0.000909377	3.23242E-05
Ibrutinib	BTK	0.35	1000	2.325680394	0.000821419	2.85424E-05
Ibrutinib	BTK	0.35	1000	1.938066995	0.00067625	2.23957E-05
Ibrutinib	BTK	0.35	1000	1.615055829	0.000610541	4.48376E-05


Supplemental Figure 1. BTK enzyme kinetics plots for ibrutinib (1)



Supplemental Table 2. BTK enzyme kinetics data for ibrutinib (0.045 nM BTK)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s ⁻¹	95%-confidence
Ibrutinib	BTK	0.045	1000	5	0.001745425	0.000253643
Ibrutinib	BTK	0.045	1000	4.166666667	0.001578286	0.000140563
Ibrutinib	BTK	0.045	1000	3.472222222	0.00126266	7.3267E-05
Ibrutinib	BTK	0.045	1000	2.893518519	0.001018874	6.78472E-05
Ibrutinib	BTK	0.045	1000	2.411265432	0.000931715	4.25298E-05
Ibrutinib	BTK	0.045	1000	2.00938786	0.000808285	3.47423E-05
Ibrutinib	BTK	0.045	1000	1.674489883	0.000656164	2.93531E-05
Ibrutinib	BTK	0.045	1000	1.395408236	0.00054087	2.39779E-05
Ibrutinib	BTK	0.045	1000	1.162840197	0.000468612	1.93592E-05
Ibrutinib	BTK	0.045	1000	0.969033497	0.000317853	1.73319E-05
Ibrutinib	BTK	0.045	1000	0.807527914	0.000365215	1.46379E-05


Supplemental Figure 2. BTK kinetics plots for ibrutinib (2)



Supplemental Table 3. BTK enzyme kinetics data for acalabrutinib (0.35 nM BTK)


Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s-1	95%-confidence
Acalabrutinib	BTK	0.35	1000	321.97	0.00097791	3.61277E-05
Acalabrutinib	BTK	0.35	1000	268.3083333	0.000956686	3.22938E-05
Acalabrutinib	BTK	0.35	1000	223.5902778	0.000907469	3.34095E-05
Acalabrutinib	BTK	0.35	1000	186.3252315	0.000846832	2.59687E-05
Acalabrutinib	BTK	0.35	1000	155.2710262	0.000714928	2.19069E-05
Acalabrutinib	BTK	0.35	1000	129.3925219	0.000692553	3.28869E-05
Acalabrutinib	BTK	0.35	1000	107.8271016	0.000487165	1.9903E-05
Acalabrutinib	BTK	0.35	1000	89.85591796	0.000462368	1.87039E-05
Acalabrutinib	BTK	0.35	1000	74.87993163	0.000420859	1.36238E-05
Acalabrutinib	BTK	0.35	1000	62.39994303	0.000367365	1.18944E-05
Acalabrutinib	BTK	0.35	1000	51.99995252	0.000290036	7.58079E-06

Supplemental Figure 3. BTK enzyme kinetics plots for acalabrutinib (1)



Supplemental Table 4. BTK enzyme kinetics data for acalabrutinib (0.045 nM BTK)


Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration nM	K _{obs} , s ⁻¹	95%-confidence
Acalabrutinib	BTK	0.045	1000	178.9	0.000997718	5.57243E-05
Acalabrutinib	BTK	0.045	1000	149.0833333	0.000849673	5.84386E-05
Acalabrutinib	BTK	0.045	1000	124.2361111	0.000775032	4.21489E-05
Acalabrutinib	BTK	0.045	1000	103.5300926	0.000655515	3.59637E-05
Acalabrutinib	BTK	0.045	1000	86.27507716	0.000631197	3.11489E-05
Acalabrutinib	BTK	0.045	1000	71.89589763	0.000496656	2.36196E-05
Acalabrutinib	BTK	0.045	1000	59.91324803	0.000462513	1.61551E-05
Acalabrutinib	BTK	0.045	1000	49.92770669	0.000367588	1.62148E-05
Acalabrutinib	BTK	0.045	1000	41.60642224	0.000337013	1.39989E-05
Acalabrutinib	BTK	0.045	1000	34.67201853	0.000267423	1.35051E-05
Acalabrutinib	BTK	0.045	1000	28.89334878	0.000258543	1.29368E-05

Supplemental Figure 4. BTK enzyme kinetics plots for acalabrutinib (2)

Supplemental Table 5. TEC enzyme kinetics data for ibrutinib (0.3 nM TEC)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s ⁻¹	95%-confidence
Ibrutinib	TEC	0.3	1000	15	0.003711357	0.000325344
Ibrutinib	TEC	0.3	1000	12.5	0.003203723	0.000259978
Ibrutinib	TEC	0.3	1000	10.41666667	0.002651187	0.00019961
Ibrutinib	TEC	0.3	1000	8.680555556	0.002183536	0.00018315
Ibrutinib	TEC	0.3	1000	7.233796296	0.001448589	0.000194026
Ibrutinib	TEC	0.3	1000	6.02816358	0.001166187	6.24796E-05
Ibrutinib	TEC	0.3	1000	5.02346965	0.001551147	4.94976E-05
Ibrutinib	TEC	0.3	1000	4.186224709	0.001399701	5.60526E-05
Ibrutinib	TEC	0.3	1000	3.48852059	0.001096532	6.49649E-05
Ibrutinib	TEC	0.3	1000	2.907100492	0.001014172	7.17453E-05
Ibrutinib	TEC	0.3	1000	2.422583743	0.000728765	7.20165E-05


Supplemental Figure 5. TEC enzyme kinetics plots for ibrutinib (1)



Supplemental Table 6. TEC enzyme kinetics data for ibrutinib (0.115 nM TEC)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s ⁻¹	95%-confidence
Ibrutinib	TEC	0.115	1000	8	0.002299512	0.000361184
Ibrutinib	TEC	0.115	1000	6.666666667	0.002103881	0.000225242
Ibrutinib	TEC	0.115	1000	5.555555556	0.002239668	0.000315757
Ibrutinib	TEC	0.115	1000	4.62962963	0.001681009	0.00016689
Ibrutinib	TEC	0.115	1000	3.858024691	0.001154146	8.28214E-05
Ibrutinib	TEC	0.115	1000	3.215020576	0.000939698	6.22118E-05
Ibrutinib	TEC	0.115	1000	2.679183813	0.000944351	7.14332E-05
Ibrutinib	TEC	0.115	1000	2.232653178	0.000731845	0.000149011
Ibrutinib	TEC	0.115	1000	1.860544315	0.000551128	0.000254758
Ibrutinib	TEC	0.115	1000	1.550453596	0.000731739	0.000384208
Ibrutinib	TEC	0.115	1000	1.292044663	0.000330385	0.000190587


Supplemental Figure 6. TEC enzyme kinetics plots for ibrutinib (2)



Supplemental Table 7. TEC enzyme kinetics data for acalabrutinib (0.3 nM TEC)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s ⁻¹	95%-confidence
Acalabrutinib	TEC	0.3	1000	804.9	0.00235	0.000128222
Acalabrutinib	TEC	0.3	1000	670.75	0.001969861	0.000115078
Acalabrutinib	TEC	0.3	1000	558.9583333	0.001576885	9.46784E-05
Acalabrutinib	TEC	0.3	1000	465.7986111	0.001410129	0.000131697
Acalabrutinib	TEC	0.3	1000	388.1655093	0.00100748	8.94748E-05
Acalabrutinib	TEC	0.3	1000	323.4712577	0.000836627	5.11058E-05
Acalabrutinib	TEC	0.3	1000	269.5593814	0.000978956	5.53943E-05
Acalabrutinib	TEC	0.3	1000	224.6328179	0.000885238	5.12951E-05
Acalabrutinib	TEC	0.3	1000	187.1940149	0.000716575	4.57308E-05
Acalabrutinib	TEC	0.3	1000	155.9950124	0.000642043	6.1225E-05
Acalabrutinib	TEC	0.3	1000	129.9958437	0.000441482	4.64066E-05



Supplemental Figure 7. TEC enzyme kinetics plots for acalabrutinib (1)



Supplemental Table 8. TEC enzyme kinetics data for acalabrutinib (0.2 nM TEC)

Compound	Enzyme	Enzyme concentration, nM	ATP concentration, uM	Compound concentration, nM	K _{obs} , s ⁻¹	95%-confidence
Acalabrutinib	TEC	0.2	1000	457.57	0.001630108	0.000129663
Acalabrutinib	TEC	0.2	1000	381.3083333	0.001398252	9.42539E-05
Acalabrutinib	TEC	0.2	1000	317.7569444	0.001188951	8.23696E-05
Acalabrutinib	TEC	0.2	1000	264.7974537	0.000964911	5.898E-05
Acalabrutinib	TEC	0.2	1000	220.6645448	0.000789929	5.28222E-05
Acalabrutinib	TEC	0.2	1000	183.8871206	0.000701655	3.70468E-05
Acalabrutinib	TEC	0.2	1000	153.2392672	0.000613403	2.93045E-05
Acalabrutinib	TEC	0.2	1000	127.6993893	0.000519269	2.58391E-05
Acalabrutinib	TEC	0.2	1000	106.4161578	0.000460735	2.78285E-05
Acalabrutinib	TEC	0.2	1000	88.68013148	0.000371238	2.65775E-05
Acalabrutinib	TEC	0.2	1000	73.90010956	0.00032225	2.50307E-05

Supplemental Figure 8. TEC enzyme kinetics plots for acalabrutinib (2)



BTK (Parameter)	Value	Units
K_{inact} (K2)	0.011573772	s ⁻¹
$K_{\text{iapp}} \cdot (1 + [\text{S}]/K_{\text{m}})$	2804.457364	nM
K-2	0	s ⁻¹
K_{i}	133.5455888	nM
$K_{\text{inact}}/K_{\text{i}}$	0.0000867	nM ⁻¹ *s ⁻¹
Residence Time (1/K-2)	Indefinite	s
FIT Statistics	Goodness of Fit	
Chi ²	7.26207E-09	
F-test	0.964287606	
T-test	0.6587841	