

Title Page

A bicistronic plasmid encoding brain-derived neurotrophic factor and urokinase plasminogen activator stimulates peripheral nerve regeneration after injury

Maxim Karagyaur, Alexandra Rostovtseva, Ekaterina Semina, Polina Klimovich, Vadim Balabanyan, Pavel Makarevich, Vladimir Popov, Dmitry Stambolsky, Vsevolod Tkachuk

Institute of Regenerative Medicine, Lomonosov Moscow State University, 27/10, Lomonosovsky Avenue, 119192, Moscow, Russia (M.K., P.M., V.T.)

Faculty of Medicine, Lomonosov Moscow State University, 27/1, Lomonosovsky Avenue, 119192, Moscow, Russia (M.K., A.R., E.S., V.B., P.M., V.P., V.T.)

Laboratory of Molecular Endocrinology, National Cardiology Research Center, Ministry of Health of the Russian Federation, 15A, 3rd Cherepkovskaya str., 121552, Moscow, Russia (E.S., P.K., V.T.)

Medical Research and Education Center, 27/10, Lomonosovsky Avenue, 119192, Moscow, Russia (D.S.)

Running Title Page

Running title: Bicistronic plasmid stimulates peripheral nerve regeneration

To whom all correspondence should be sent:

Maxim Karagyaur, Ph.D., M.D.

Institute of Regenerative Medicine, Lomonosov Moscow State University

27/10, Lomonosovsky Avenue, 119192, Moscow, Russia

E-mail: m.karagyaur@mail.ru

Tel +7-926-593-6877

Fax +7-499-726-5547

Number of text pages: 37

Number of tables: 0 (+1 in Supplementary Materials)

Number of figures: 6 (+1 in Supplementary Materials)

Number of references: 34

Number of words in abstract: 222 words

Number of words in introduction: 677 words

Number of words in discussion: 1399 words

ABBREVIATIONS: AD-converter, analog-to-digital converter; b-Gal, b-Galactosidase; CCD camera, charge-coupled device camera; CMV, Citomegalovirus; CNAP, compound nerve action potential; DMEM, Dulbecco's modified Eagle medium; ELISA, enzyme-linked immunosorbent assay; FBS,

fetal bovine serum; hBDNF, human brain-derived neurotrophic factor; HCV, Hepatitis C virus; HEK293, Human embryonic kidney 293 cell line; huPA, human urokinase plasminogen activator; IRES, Internal Ribosome Entry Site; MEM NEAA, Minimal essential medium Non-essential amino acids; muPA, murine urokinase plasminogen activator; pNCure, a plasmid for nerve cure; pNCure-f/rIntr, a plasmid for nerve cure with a synthetic intron (in forward or reverse orientation) before hBDNF start codon; PSGs, Penicillin-streptomycin-glutamine solution; RT-PCR, reverse transcription polymerase chain reaction; uPAR, urokinase plasminogen activator receptor; WPRE, Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element.

Recommended section assignment: Chemotherapy, Antibiotics, and Gene Therapy

Abstract

Timely nerve restoration is an important factor for the successful regeneration of tissues and organs. It is known that axon regeneration following nerve injury is a multifactorial process that depends on the local expression of neurotrophins, including brain-derived neurotrophic factor (BDNF). Along with the survival of neurons, the active reorganization of the extracellular matrix is an important step for the growth of axons to their targets. Urokinase serine protease is part of the plasminogen activator system, which provides the vectoriality of the process of fibrinolysis and matrix reorganization, facilitating the growth of nerves to their targets. Based on this and in view of the results of our previous studies, we suggest that a combined bicistronic plasmid encoding the complementary proteins BDNF and urokinase may be beneficial in nerve regeneration. The ability of this bicistronic plasmid to stimulate nerve restoration was confirmed by *in vitro* stimulation of Neuro2a neurite growth and *in vivo* nerve conductivity and histology studies. To our knowledge, this is the first article that demonstrates the effectiveness of a bicistronic plasmid containing the human genes BDNF and uPA in the regeneration of the injured peripheral nerve. The results obtained demonstrate that plasmid vectors encoding several complementary-active therapeutic proteins may serve as a basis for developing prospective treatments for a wide range of multicomponent neural system disorders, such as nerve trauma.

Significance Statement

This study is the first to show the effectiveness of using a bicistronic plasmid encoding complementary-active human proteins brain-derived neurotrophic factor (hBDNF) and urokinase plasminogen activator (huPA) in the regeneration of the crushed peripheral nerve in a murine model.

Introduction

Blood vessels and nerve fibers are essential components of any tissue and provide for the normal development, functioning, and regeneration of tissues. However, tissue injuries sometimes lead to damage to blood vessels and nerve fibers. Following injury, both the regenerating blood vessels and nerve fibers begin to grow from the remaining stump. Unlike the regeneration of blood vessels, the regeneration of nerve fibers has rather restricted potential, which is caused by remote neuron cell bodies localization predominantly within the central nervous system, and the short-time profile of neurotrophic factor production (Gordon B.J and Gordon T., 2003). Contemporary research on tissue regeneration describes it as a complex multi-step process that requires the participation of a variety of cellular and signaling components, which interchange and/or cooperate with each other during the regeneration progress (Li et al., 2016; Wang et al., 2017; Nimiritsky et al., 2019). This may explain the relatively moderate effects of monogenic gene therapy in clinical studies, despite the promising results that were obtained in preclinical studies (Gupta et al., 2009; Makarevich et al., 2015). Despite advances in medicine, there are no effective therapeutic approaches to, or treatments for nerve injury, and persistent disability is reported in 75% of all cases following severe nerve injury (Odinak and Zhivolupov, 2009; Shevelev, 2005).

The modern concept of successful regeneration highlights the necessity of multimodal and complex interventions and stimulation of the regenerating tissue (Li et al., 2016; Wang et al., 2017; Nimiritsky et al., 2019). With this in mind, the

activation of several signaling cascades that drive different modalities and complement each other seems feasible. Among the therapeutic compounds that complement each other during injured nerve regeneration, neurotrophic factors and proteases are of particular interest. In this combination, a neurotrophic factor would be used to stimulate the survival and growth of nerve fibers and a protease would be used to promote neurite growth cone migration via cleavage of the endoneurial channels from inhibitory molecules and fibrin deposits, as well as to activate endogenous growth factors by proteolysis (Benga et al., 2017; Karagyaur et al., 2017).

It is known that the expression of neurotrophins in an injured nerve is not sustained for longer than a month. Short-term neurotrophin expression and a lack of integrity and passableness of the nerve trunk are the main challenges for successful nerve regeneration (Jia et al., 2014; Gordon T. and English, 2016) due to increased injured neuron death and a lack of neurite outgrowth. In contrast, prolonged neurotrophin secretion (e.g. via neurotrophin gene delivery) and adequate neurite channels clearance helps the regenerating neurites to reach their target organs. Brain-derived neurotrophic factor (BDNF) can be used as a stimulator for motor neuron regeneration, according to previous research (Carlone et al., 1987; Karagyaur et al., 2015). A wide range of proteins are suitable complements to BDNF. However, proteases, including plasminogen activators that stimulate fibrinolysis and angiogenesis, take part in neurite channels clearance and activate BDNF, are among the most attractive. Among these,

urokinase plasminogen activator (uPA) particularly stands out. It was recently shown that uPA is necessary for the directed growth of axons via binding to its receptor (uPAR) on the surface of growth cones (Semina et al., 2016; Merino et al., 2017).

In our previous work, we showed that single-gene plasmids encoding hBDNF or muPA stimulated the recovery of an injured nerve (Karagyaur et al., 2015). Alongside existing research, this finding confirms the important role of BDNF and uPA in the process of neuron survival and neurite regeneration. Taking into account the complexity and multiple stages of the regenerative process, we surmised that a bicistronic genetic construct encoding both hBDNF and huPA could be beneficial for enhancing nerve regeneration. We decided to create the desired bicistronic construct in the form of a plasmid, because plasmids are still the most safe and convenient genetic instruments for heterologous protein expression in medicine and basic science. Here, we show that a bicistronic plasmid construction encoding growth factors of complementary modalities (BDNF and uPA) stimulates nerve fiber restoration after nerve crush in a clinically relevant model of nerve injury.

Materials and Methods

Plasmids

To stimulate nerve regeneration, a bicistronic vector encoding hBDNF and huPA cDNAs was created. As a backbone, a pVax1m plasmid was used, which was obtained by introducing a SV40 enhancer sequence between a CMV promoter and a CMV enhancer in the pVax1 vector (ThermoFisher Scientific, #V26020) to increase plasmid uptake by the cell nucleus (Dean et al., 2005). hBDNF and huPA cDNAs were obtained by RT-PCR of total RNA extracted from human adipose-derived mesenchymal stromal cells, which were obtained from the biomaterial collection within the frame of Moscow University Project "Noah's Ark" (primer sequences are listed in Suppl. Table 1). hBDNF cDNA was introduced into the pVax1m vector subsequent to the CMV promoter and the Kozak consensus sequence (GCCACC) using NheI and HindIII restriction sites (Kozak, 1984). huPA cDNA (cloned using Acc65I and EagI restriction sites) was separated from hBDNF by HCV IRES (Lukavsky, 2009), assembled from synthetic oligonucleotides (PCR-amplified and cloned using HindIII and Acc65I restriction sites). According to the study of R. Jubin (Jubin et al., 2000) HCV IRES provides for the production of the second protein, huPA, at a 10-fold lower molar rate than the first protein, hBDNF. Each cDNA carries its own start and stop codons (the start codon for huPA is built in the HCV IRES). The plasmid pVax1m-hBDNF-IRES-huPA was named "a plasmid for nerve cure" (pNCure for short). To

increase the productivity of the plasmid, four additional variants of the plasmid were constructed with a synthetic intron and WPRE sequence (Fig. 1). The synthetic intron was designed de novo, assembled from synthetic oligonucleotides and introduced between the CMV promoter and Kozak consensus sequence (NheI restriction site). Intron's sequence is 5'-**GTGAGTAACAGAGATCTAATACGCTTCTCTTGCTAAGCCTAACCTTT TCTGTCTTGTTTCTGCTTTCCCCCCCCCACAG**-3' (bold letters mark intron key motifs). It was not obvious which intron orientation (forward vs reverse) would be advantageous in protein production, so both plasmid variants were stored (pNCure-f/rIntr). The WPRE sequence was cloned from pAAV-EF1a-tdTomato-WPRE-pGHpA (a gift from Botond Roska, Addgene plasmid #67527) (Wertz et al., 2015) and introduced into pNCure-f/rIntr variants between the huPA stop codon and bovine growth hormone polyadenylation signal (XbaI restriction site). The obtained plasmid variants were tested in parallel with the previously described pVax1-hBDNF plasmid (Karagyaur et al., 2015).

Treatment

Animals

The study was performed on mature male F1 CBA/C57Bl mice aged 8-10 weeks, with standard weight characteristics (20–23 g), obtained from SPF vivarium of Faculty of Medicine (Lomonosov Moscow State University). Animal

housing and research procedures were conducted in compliance with Directive 201/63/EU.

Intramuscular plasmid injection

In our previous work on the intramuscular delivery of plasmids, we used a quick injection of a plasmid solution at a single muscle point, followed by electroporation. However, for therapeutic use, multiple points for plasmid injection are used (muscle infiltration) with no electroporation, as this causes massive necrosis of muscle tissue, of which approximately 10–30% depends on electroporation voltage (Manthorpe et al., 1993; Karagyaur et al., 2015). Therefore, here we used a more clinically relevant plasmid delivery protocol, by substituting the single injection electroporation regime with multiple intramuscular injections and no electroporation. To test the efficacy of using multiple injections with no electroporation, a solution containing pVax1-b-Gal or pNCure-fIntr (as the most productive) plasmid (60 ul, 1ug/ul) was injected at a single point or at six distinct points in the anterior tibial muscles of experimental mice (three muscles per each group). Further, 72 h after the injections were administered, the mice were sacrificed and their anterior tibial muscles were isolated. To assess b-Gal expression, the appropriate muscles were preserved in Tissue-Tek® O.C.T. Compound (Sakura) and sliced. Slices were fixed and stained for b-Galactosidase expression (X-Gal Staining Protocol) and b-Gal positive muscle fibers were counted. To assess how the injection of a plasmid into

multiple sites of a muscle changed recombinant protein production, an explant culture (Jang et al., 2004) was prepared from the muscles in which pNCure-fIntr had been injected. These muscle explants were conditioned in DMEM High Glucose (HyClone, #SH30022FS) with 2% fetal bovine serum (HyClone, #SH3007103) and 1% Penicillin-Streptomycin-Glutamine solution (ThermoFisher Scientific, #10378016). The conditioned medium was concentrated 20x using a combination of desalting-lyophilization-reconstitution and analyzed for recombinant hBDNF and huPA concentration as it was described above. The concentration procedure was necessary to adjust for the rather low productivity of the muscle explants and moderate sensitivity of the hBDNF and huPA ELISA systems.

Nerve injury model

To assess the ability of the plasmid to stimulate peripheral nerve regeneration following nerve crush, a mouse model was used (Karagyaur et al., 2015). Forty mice were randomized into four groups (n = 10): a positive and a negative control groups and two experimental groups (Suppl. Fig. 1). This number of experimental animals is the minimum needed to assess the pharmacological effects and conduct a statistical analysis of the results. Mice were anesthetized with avertin injections and the common peroneal nerve (n. peroneus communis) was isolated from the surrounding tissue under sterile conditions and crushed using 1 mm wide forceps with silicone coated branches two times for 20 seconds. After the wound site was

sewn, the plasmids (pVax1m 1 ug/ul (as control), pNCure-fIntr 1 or 2 ug/ul) were injected at six distinct sites in the anterior tibial muscles of mice from the Control, pNCure1, and pNCure2 groups. Plasmid injections were repeated at 4, 9 and 13 days after nerve crush (Suppl. Fig. 1).

Mice from the positive control group (B12) were given subcutaneous injections of a vitamin B12 solution (65 ug/kg) on the day of the trauma and then every 48 h until the end of the experiment.

Measurements

HEK293 transfection and ELISA of conditioned media

To determine which plasmid variant provided maximal recombinant protein expression, HEK293 cultures were transfected with pNCure plasmid variants, pVax1-hBDNF and the control plasmid pVax1m (in duplicates) using Lipofectamine2000, according to the manufacturer's instructions. Transfected cells were cultivated in DMEM High Glucose (HyClone, #SH30022FS) supplemented with 2% fetal bovine serum (FBS) (HyClone, #SH3007103) and 1% Penicillin-Streptomycin-Glutamine (PSG) solution (ThermoFisher Scientific, #10378016). 72 h after transfection, the conditioned media was collected and analyzed for hBDNF and huPA concentrations using the Human Free BDNF Quantikine ELISA Kit (R&D, #DBD00) and the uPA (URK) Human ELISA Kit (Abcam, #ab119611), respectively.

In vitro testing of the ability of hBDNF and huPA to stimulate neurite outgrowth in mouse Neuro2a cells

The murine neuroblastoma cell line Neuro2a was used as a neurite outgrowth *in vitro* model. This cell line is known to differentiate (i.e. enable neurite outgrowth) in serum deprivation conditions (Semina et al. 2016; Namsi et al., 2018). The Neuro2a cell line was cultured in DMEM High Glucose (HyClone, #SH30022FS) supplemented with 10% FBS (HyClone, #SH3007103), 1% PSG solution (ThermoFisher Scientific, #10378016) and 1% MEM Non-Essential Amino Acids (MEM NEAA) (ThermoFisher Scientific, #11140050).

HEK293 cultures were transfected with pVax1m, pVax1-hBDNF, and pNCure-fIntr plasmids. Neuro2a cells were seeded in a 24-well plate in low confluent (10–15%). After 24 h of cultivation of the cells in DMEM High Glucose with 10% FBS, the media was replaced with DMEM High Glucose, containing 1% FBS to stimulate neurite outgrowth. Conditioned media from HEK293 cells transfected with pNCure-fIntr, pVax1-hBDNF or control plasmid pVax1m was added to the Neuro2a cells (with final hBDNF concentration in wells “pNCure-fIntr”, “pVax1-hBDNF” about 50 ng/ml).

Some wells were used as positive (1% FBS) or negative (10% FBS) control groups to assess neurite outgrowth. Each group existed in triplicates. Cells were incubated for at least 72 h. Pictures were taken with a bright light microscope equipped with a CCD camera. Neurites were counted in each group using ClickCounter2 software and normalized per number of cells.

Nerve regeneration assessment

On the 18th day after injury, the effect of the multiple plasmid injections on functional and structural nerve recovery was assessed. Nerve conductivity and the number of axons were evaluated. Nerve conductivity measurements were performed as previously described (Karagyaur et al., 2015). The animals were sacrificed and the injured nerves were isolated and placed in a Laily solution (135.0 mM NaCl, 4.0 mM KCl, 2.2 mM CaCl₂, 1.0 mM MgCl₂, 0.9 mM NaH₂PO₄, 16.3 mM NaHCO₃, 11.0 mM D-glucose, pH 7.2-7.4). Three series of compound nerve action potentials (CNAP) were independently recorded for each nerve using a monopolar aspirating silver electrode, 10 mm distal from the injury site, connected to an AD-converter and output and were analyzed using PowerGraph Professional 3.3 software. The length of the latency period and the amplitude of the CNAPs were measured. The number of axons was evaluated on formalin-fixed frozen sections of a common peroneal nerve fragment, 3 mm distal to injury site, using immunofluorescent staining with rabbit anti-NF200 antibody («Sigma», cat # 4142) and morphometric analysis (Lopatina et al., 2011; Karagyaur et al., 2015).

Statistical Analysis

Statistical analysis was performed using SigmaPlot11.0 Software. Data were assessed for normality of distribution using the Kolmogorov–Smirnov test. Differences between the treatment and control groups were then analyzed using

the Student t-test or ANOVA on ranks Dunn's test, depending whether there was normal distribution or not. Data are expressed as mean \pm standard deviation or median (25%; 75%) depending on the test used. $P < 0.05$ was considered statistically significant.

Results

Plasmid variant pNCure-fIntr provides maximum expression of hBDNF and huPA in HEK293 cells

Analysis of the expression of recombinant hBDNF and huPA in HEK293 cells showed that the plasmid pNCure variant containing the synthetic intron sequence fIntr produced significantly higher hBDNF than the basic pNCure plasmid (69.3 ± 9.1 vs 53.3 ± 7.3 ng / ml [n = 3, *P < 0.05]). However, huPA production did not differ significantly between the pNCure-fIntr and pNCure plasmid variants: 9.2 ± 1.4 vs 9.0 ± 2.5 ng / ml (n = 3) (Fig. 2). The addition of the WPRE element pNCure-f/rIntr-WPRE did not significantly increase the production of recombinant proteins. However, it did increase the size of the plasmid by 10%. For this reason, we decided not to use variants of the pNCure plasmid containing WPRE in this study. The effect of the pNCure-fIntr plasmid on hBDNF expression was almost equivalent to that of pVax1-hBDNF, as previously described (Karagyaur et al., 2015).

pNCure-fIntr plasmid stimulates neurite outgrowth in a Neuro2a cell model

The conditioned media stimulated neurite outgrowth in all groups compared to negative control group (DMEM supplemented with 10% FBS). However, the most prominent effect was observed in the pVax1-hBDNF and pNCure-fIntr groups, in which 29.4 ± 2.5 % and 34.3 ± 3.5 % of cells had neurites, respectively.

The difference was significant compared to the negative control, positive control and pVax1m groups, in which 8.2 ± 1.4 , 22.9 ± 2.7 and 18.3 ± 2.1 % of cells had neurites, respectively ($n = 20$, $*P < 0.05$) (Fig. 3). The number of neurites in the positive control and pVax1m groups differed significantly from the negative control group ($n = 20$, $**P < 0.05$). No significant difference was observed between the pVax1-hBDNF and pNCure-fIntr groups, but a tendency toward better neuritogenesis was noted in the pNCure-fIntr group.

Multiple intramuscular injections of a plasmid are as efficient as a single-point injection

In our previous studies, we used a single intramuscular injection for therapeutic gene delivery to stimulate repair of the damaged nerve (Karagyaur et al., 2015). However, the approach used was not relevant in terms of clinical application. Hydrodynamic shock and electroporation are not used in clinical settings as they cause pain and significant traumatic effects. Multiple intramuscular injections of small amounts of a plasmid solution (known as muscle chipping) are more clinically relevant, but it was unclear whether they would provide the acceptable level of recombinant protein expression. Using the marker plasmid (pVax1-b-Gal), we showed that muscle chipping provides plasmid delivery to muscle fibers of equal efficacy to a per saltum injection accompanied by hydrodynamic shock. Injections at six different points of the anterior tibial muscle provided transfection of 68.3 ± 12.7 muscle fibers (β -galactosidase

positive), while a per saltum injection of a pVax1-b-Gal plasmid solution accompanied by hydrodynamic shock resulted in transfection of only 52.8 ± 11.5 muscle fibers (the difference is not significant, $n = 3$) (Fig. 4).

The analysis of hBDNF and huPA in a concentrated conditioned culture medium from muscle explants revealed that muscle chipping with small amounts of the plasmid pNCure-fIntr provided the same level of recombinant protein production. Six injections of pNCure-fIntr (muscle chipping) resulted in 136.5 ± 18.4 pg/ml of hBDNF and 12.3 ± 1.7 pg/ml of huPA, while a single intramuscular injection of pNCure-fIntr accompanied by hydrodynamic shock resulted in 107.3 ± 12.5 pg/ml of hBDNF and 9.6 ± 2.0 pg/ml of huPA in a culture medium. The difference between two groups was not significant, but an increase in recombinant protein expression in the muscle chipping group was noted ($n = 3$) (Fig. 4).

Since the effectiveness of the clinically relevant method of muscle chipping with the plasmid solution was confirmed experimentally, this delivery method was used for evaluating of the therapeutic efficacy of a constructed bicistronic plasmid vector.

A bicistronic plasmid encoding hBDNF and huPA stimulates post-traumatic functional and structural nerve recovery

Crushing of the common peroneal nerve caused nerve dysfunction, as evidenced by a decrease in CNAP amplitude and CNAP latent period extension (Suppl. Fig. 2). Intramuscular injections of pNCure-fIntr stimulated the functional

and structural recovery of the crushed nerve. On the 18th day of the experiment, CNAP amplitude was 1.7- and 2.2-fold higher in animals from the pNCure1 and pNCure2 groups compared to the Control group animals, which received injections of an empty pVax1m vector : 0.475 (0.338; 0.612) and 0.627 (0.506; 0.832) mV vs 0.285 (0.245 ; 0.353) mV, respectively (n = 10, ***P < 0.001 vs control) (Fig.5). A comparable increase in CNAP amplitude was also observed in the B12 control group – 0.457 (0.329; 0.573) mV (n = 10; *P < 0.05 vs control). CNAP amplitudes between the pNCure1 and pNCure2 groups did not differ significantly (P = 0.174).

The latency period of the CNAP of the damaged nerve (conductivity indicator) did not differ significantly in all groups, but a tendency to its diminishing diminish with dose increments was observed in the pNCure1 and pNCure2 groups (Fig. 5). The latency in the pNCure1 group was 0.715 (0.622; 0.893) ms, and was 0.507 (0.45; 0.887) ms in the pNCure2 group, while in Control and B12 groups, the latency was 0.861 (0.68; 0.983) and 0.771 (0.7; 0.94) ms, respectively.

An analysis of nerve cryosections stained for markers of mature axons NF200 revealed that the pNCure-fIntr plasmid stimulated the recovery of the nerve structure. Both the number and size of the positively stained structures in nerves from the pNCure1 and pNCure2 groups was larger compared to nerves from the Control group animals. On the 18th day of the experiment, nerve sections from the pNCure1 and pNCure2 groups contained 573 (489; 619) and 554 (518;

633) positively stained structures (nerve fibers), respectively ($n > 5$, $*P < 0.05$ vs Control), while in Control and B12 groups, the number of positively stained structures was 276 (234; 311) and 461 (407; 545) ($n > 5$, $*P < 0.05$ vs Control) (Fig.6). No significant difference between the pNCure1, pNCure2, and B12 groups was observed, though a tendency toward a higher number of restored nerve fibers was noted in the pNCure1 and pNCure2 groups.

Discussion

Successful tissue regeneration requires restoration of its blood and nerve supply. While a blood vessel can be regrown from a number of endothelial cell islets in a multifocal mode, a damaged nerve fiber regenerates only from its proximal stump along the nerve bed and is highly sensitive to neurotrophin expression (Benga et al., 2017; Karagyaur et al., 2017). The lack of effective specific drugs and the abundance of disabilities caused by nerve injury means that new therapeutic approaches should be sought. Regeneration is known to be a complex and multi-stage process, and this changes the paradigm of the current pro-regenerative therapy (Li et al., 2016; Wang et al., 2017; Nimiritsky et al., 2019). It is becoming increasingly clear that it is necessary to influence the different stages of the regenerative process in order to obtain the desired outcome. This can be achieved by combining medicines (e.g. gene therapy molecules) with different modalities.

In this paper, we studied the efficacy of a combined gene therapy with drug encoding brain-derived neurotrophic factor and urokinase plasminogen activator on the stimulation of damaged nerve fiber recovery in a model of peripheral nerve injury. The combination of a growth factor and a protease appeared to be beneficial: BDNF stimulated neuron survival and neurite growth (Ebendal, 1992), while uPA is a key molecule in fibrin degradation (the main inhibitor of neurite growth after trauma), neurite navigation, matrix remodeling and growth factor maturation (Siconolfi and Seeds, 2001). For proper folding, most proteins are

synthesized in preforms and then mature via protease-mediated cleavage (matrix metalloproteinases, plasmin, plasminogen activators). BDNF is one such molecule. Moreover, its proform, proBDNF, has greater affinity to the alternative neurotrophin receptor p75 that activates neuron death program (McPhee & Barker, 1997). Mature BDNF stimulates neuron survival and neurite elongation via TrkB receptor activation (Ebendal, 1992) and earlier research has shown that plasmin, activated by uPA or tPA, causes maturation of BDNF and other growth factors (Rifkin et al., 1997).

As the aim of this study was to create a gene therapy drug for treating nerve injury in humans, human orthologues of BDNF and uPA were encoded in pNCure plasmid. The BDNF sequence is highly conservative among the species and its human orthologue can be successfully used and tested in murine models (Lucini et al., 2018). However, uPA functions by interacting with the receptor uPAR, which has strong species-specificity (Lin et al., 2010) and this makes the usage and testing of the human uPA orthologue in murine models impossible. Despite this, the ability of hBDNF and muPA to stimulate the regeneration of the crushed murine nerve was demonstrated in our previous studies (Karagyaur et al., 2015). Moreover, here we showed that a bicistronic plasmid encoding hBDNF and huPA expressed in human cell line HEK293 has a tendency to promote better neuritogenesis in Neuro2a cells *in vitro*. One possible explanation is that the expression of huPA in human cell lines allows the activation of huPA via its binding to uPAR on the HEK293 cell surface, which leads to the conversion of

plasminogen to plasmin and further growth factor activation. Previously, it was shown that the plasminogen/plasmin system plays an important role in the activation of many proteins, including growth factors (Rifkin et al., 1997). Unexpectedly, the pVax1m conditioned media also provided significant neurite outgrowth. We propose that this could be caused by the basic productivity of cytokines and growth factors by the HEK293 cell line.

To increase the productivity of the basic pNCure plasmid, some short regulatory elements were introduced. A short synthetic intron, inserted before the Kozak consensus sequence and start codon, significantly increased protein expression, while the addition of the WPRE element after the stop codon did not. As they increased the size of the plasmid by 10% and had an insignificant influence on protein expression, the variants of pNCure containing WPRE were not used in this study. The results of the study demonstrated that the effect of the pNCure plasmid variants constructed here on hBDNF production did not differ dramatically from the previously studied pVax1-hBDNF plasmid, despite the addition of the second gene urokinase plasminogen activator and an increase in plasmid size of 20%.

In our previous study (Karagyaur et al., 2015), we showed that a single intramuscular injection of hBDNF encoding plasmid, followed by an electroporation procedure, provided hBDNF expression in murine tissues for 7–10 days and stimulated the repair of a damaged nerve. However, the approach used was not relevant to clinical applications; hydrodynamic shock and

electroporation are not used in clinical settings because they cause pain and traumatic effects. Furthermore, it should be kept in mind that recombinant proteins in human tissues have to be expressed for weeks or even months to complete the regeneration process. The small size and high regenerative potential ensure effective recovery of a crushed nerve, even with weak expression of neurotrophic factors in mice but not in humans. The clinical use of gene-therapeutic drugs requires repeated injections and prolonged protein expression. Here, we showed that the clinically relevant method of multiple intramuscular plasmid injections is as efficient as the previously used single site injection. Additional tests showed that repeated multiple intramuscular injections of a bicistronic plasmid still provided local protein expression and did not increase their ectopic expression (in press). Since the effectiveness of this delivery method was confirmed experimentally, it was used in this study for bicistronic pNCure-fIntr plasmid intramuscular delivery in a murine nerve injury model.

Based on our previous experience of *in vivo* testing of hBDNF and muPA encoding plasmids, the results of previous *in vitro* testing and the absence of relevant human nerve injury models, we decided to test the activity of a pNCure-fIntr bicistronic plasmid in the familiar murine nerve crush model. We were not expecting expressed huPA to work as well in mice as it did in humans.

In vivo testing in mice showed that the bicistronic pNCure-fIntr plasmid stimulated the conductivity restoration of a damaged nerve. The amplitude of CNAP in mice treated with the pNCure-fIntr plasmid was greater than in the

control group. Because CNAP's amplitude in direct ratio characterizes the number of nerve fibers involved in nerve impulse conduction (Fugleholm et al., 2000; Vleggeert-Lankamp, 2007), we concluded that the pNCure-fIntr plasmid stimulated neurite post-traumatic survival and outgrowth *in vivo*. This data confirmed the results of *in vitro* testing and was supported by immunohistochemical studies of nerve structure (Fig.6). The latent period (i.e., the time interval from the moment of stimulation to the moment of CNAP registration) that characterizes the speed of nerve conductance in inverse ratio was not restored significantly in the pNCure-fIntr group, though some improvement was observed. This indicates that the pNCure-fIntr plasmid did not significantly increase the myelination of restored nerve fibers. The most prominent effect of the pNCure-fIntr plasmid was observed at the higher dose 120 ug (Fig. 5).

Because the concept of the experiment indicated multiple plasmid injections that could not last less than 18 days, relatively sparing method of damage to the peripheral nerve and high endogenous regenerative potential of the mice leveled the statistically significant difference during the latent period between the experimental and control groups. This was confirmed by the magnitude of the CNAP latent period in the experimental and control groups, in which the values were close to those for the intact nerve (0.6 ms).

Vitamin B12 was used as a positive control for *in vivo* studies as it is one of the few drugs registered in Russia for treating nerve injuries and it shares signal cascades with the neurotrophines ras/erk and Akt.

We found that the bicistronic plasmid pNCure-fIntr, encoding the human orthologues BDNF and uPA, was able to stimulate neuritogenesis *in vitro* and post-traumatic peripheral nerve conductivity and structure restoration *in vivo*. This apparently occurred due to the activation of neuron survival and growth of the crushed axons.

To our knowledge, this is the first study that has tested a bicistronic plasmid construction encoding a neurotrophin and plasminogen activator for enhancement of the multicomponent and multi-stage process of nerve fiber regeneration.

As treatment outcomes for many neurological and neurosurgery disorders are poor, and as BDNF and uPA are important growth factors in nerve regeneration, this successfully tested bi-modal plasmid, in combination with an appropriate delivery system, could serve as the basis for developing new prospective treatments for a wide range of multicomponent neural system disorders, such as strokes and neurodegenerative disorders.

Authorship Contributions

Participated in research design: Karagyaur, Semina, Balabanyan, Stambolsky, Tkachuk

Conducted experiments: Karagyaur, Rostovtseva, Semina, Klimovich, Popov

Contributed new reagents or analytic tools: Karagyaur, Semina, Balabanyan

Performed data analysis: Karagyaur, Rostovtseva, Semina, Klimovich, Balabanyan, Makarevich, Stambolsky, Tkachuk

Wrote or contributed to the writing of the manuscript: Karagyaur, Semina, Balabanyan, Makarevich, Stambolsky, Tkachuk

References

Benga A, Zor F, Korkmaz A, Marinescu B, and Gorantla V (2017) The neurochemistry of peripheral nerve regeneration. *Indian J Plast Surg* **50(1)**:5-15.

Carlone R, Kim JK, and Rathbone M (1987) Purification of a chick brain-derived growth factor by reversed-phase high-performance liquid chromatography, *J Neurosci* **7(7)**:2163-2167. doi: 10.1523/JNEUROSCI.07-07-02163.1987.

Dean DA, Strong DD, and Zimmer WE (2005) Nuclear entry of nonviral vectors. *Gene Ther* **12(11)**:881-890.

Ebendal T (1992) Function and evolution in the NGF family and its receptors. *J Neurosci Res* **32(4)**:461-470.

Fugleholm K, Schmalbruch H, and Krarup C (2000) Post reinnervation maturation of myelinated nerve fibers in the cat tibial nerve: chronic electrophysiological and morphometric studies. *J Peripher Nerv Syst* **5(2)**:82-95.

Gordon BJ and Gordon T (2003) Neurotrophic factors and their receptors in axonal regeneration and functional recovery after peripheral nerve injury. *Mol Neurobiol* **27(3)**:277-323.

Gordon T and English AW (2016) Strategies to promote peripheral nerve regeneration: electrical stimulation and/or exercise. *Eur J Neurosci* **43(3)**:336–350. doi: 10.1111/ejn.13005.

Gupta R, Tongers J, and Losordo DW (2009) Human studies of angiogenic gene therapy. *Circ Res* **105(8)**:724-736.

Jang HS, Kim HJ, Kim JM, Lee YS, Kim KL, Kim JA, Lee JY, Suh W, Choi JH, Jeon ES, Byun J, and Kim DK (2004) A Novel ex Vivo Angiogenesis Assay Based on Electroporation-Mediated Delivery of Naked Plasmid DNA to Skeletal Muscle. *Mol Ther* **9**:464–474.

Jia X, Romero-Ortega MI, Teng YD (2014) Peripheral nerve regeneration: mechanism, cell biology, and therapies. *Biomed Res Int* **145304**. doi: 10.1155/2014/145304.

Jubin R, Vantuno NE, Kieft JS, Murray MG, Doudna JA, Lau JY, and Baroudy BM (2000) Hepatitis C virus internal ribosome entry site (IRES) stem loop IIIId contains a phylogenetically conserved GGG triplet essential for translation and IRES folding. *J Virol* **74(22)**:10430-10437.

Karagyaur M, Dyikanov D, Makarevich P, Semina E, Stambolsky D, Plekhanova O, Kalinina N, and Tkachuk V (2015) Non-viral transfer of BDNF and uPA stimulates peripheral nerve regeneration. *Biomed Pharmacother* **74**:63-70. doi: 10.1016/j.biopha.2015.07.002.

Karagyaur MN, Makarevich PI, Shevchenko EK, Stambolsky DV, Kalinina NI, and Parfyonova YeV (2017) Modern approaches to peripheral nerve regeneration after injury: The prospects of gene and cell therapy. *Genes & cells* **12(1)**. doi: 10.23868/201703001.

Kozak M (1984) Point mutations close to the AUG initiator codon affect the efficiency of translation of rat preproinsulin in vivo. *Nature* **308(5956)**:241–246. doi: 10.1038/308241a0.

Li J, Zhang S and Amaya E (2016) The cellular and molecular mechanisms of tissue repair and regeneration as revealed by studies in *Xenopus*. *Regeneration (Oxf)* **3(4)**:198–208. doi: 10.1002/reg2.69.

Lin L, Gårdsvoll H, Huai Q, Huang M, and Ploug M (2010) Structure-based Engineering of Species Selectivity in the Interaction between Urokinase and Its Receptor. *J Biol Chem* **285**:10982-10992. doi: 10.1074/jbc.M109.093492.

Lopatina T, Kalinina N, Karagyaur M, Stambolsky D, Rubina K, Revischin A, Pavlova G, Parfyonova Y, Tkachuk V (2011) Adipose-derived stem cells stimulate regeneration of peripheral nerves: BDNF secreted by these cells promotes nerve healing and axon growth de novo. *PLoS One* **6(3)**: e17899. doi: 10.1371/journal.pone.0017899.

Lucini C, D'Angelo L, Cacialli P, Palladino A, and de Girolamo P (2018) BDNF, Brain, and Regeneration: Insights from Zebrafish. *Int J Mol Sci* **19(10)**:e3155. doi: 10.3390/ijms19103155.

Lukavsky PJ (2009) Structure and function of HCV IRES domains. *Virus Res* **139(2)**:166-171.

Makarevich PI, Rubina KA, Diykanov DT, Tkachuk VA, and Parfyonova YV (2015) Therapeutic Angiogenesis Using Growth Factors: Current State and Prospects for Development. *Kardiologiya* **55(9)**:59-71.

Manthorpe M, Cornefert-Jensen F, Hartikka J, Felgner J, Rundell A, Margalith M, and Dwarki V (1993) Gene therapy by intramuscular injection of

plasmid DNA: studies on firefly luciferase gene expression in mice. *Hum Gene Ther* **4(4)**:419-431. doi: 10.1089/hum.1993.4.4-419.

McPhee IJ and Barker PA (1997) Brain-derived Neurotrophic Factor Binding to the p75 Neurotrophin Receptor Reduces TrkA Signaling While Increasing Serine Phosphorylation in the TrkA Intracellular Domain. *J Biol Chem* **272(38)**:23547–23551.

Merino P, Diaz A, and Yepes M (2017) Urokinase-type plasminogen activator (uPA) and its receptor (uPAR) promote neurorepair in the ischemic brain. *Receptors Clin Investig* **4(2)**:e1552.

Namsi A, Nury T, Hamdouni H, Yammine A, Vejux A, Vervandier-Fasseur D, Latruffe N, Masmoudi-Kouki O, and Lizard G (2018) Induction of Neuronal Differentiation of Murine N2a Cells by Two Polyphenols Present in the Mediterranean Diet Mimicking Neurotrophins Activities: Resveratrol and Apigenin. *Diseases (Basel, Switzerland)* **6(3)**:e67. doi: 10.3390/diseases6030067.

Nimiritsky PP, Eremichev RY, Alexandrushkina NA, Efimenko AY, Tkachuk VA, and Makarevich PI (2019) Unveiling Mesenchymal Stromal Cells' Organizing Function in Regeneration. *Int J Mol Sci* **20(4)**:e823. doi: 10.3390/ijms20040823.

Odinak MM and Zhivolupov SA (2009) Diseases and injuries of the peripheral nervous system [russian]. 1st ed. SpetsLit, St. Petersburg.

Rifkin DB, Gleizes PE, Harpel J, Nunes I, Munger J, Mazzieri R, and Noguera I (1997) Plasminogen/plasminogen activator and growth factor activation. *Ciba Found Symp* **212**:105-115. discussion 116-118.

Semina E, Rubina K, Sysoeva V, Rysenkova K, Klimovich P, Plekhanova O, and Tkachuk V (2016) Urokinase and urokinase receptor participate in regulation of neuronal migration, axon growth and branching. *Eur J Cell Biol*, **95**:295-310.

Shevelev IN (2005) Traumatic lesions of the brachial plexus [russian]. 1st ed. Moscow, Moscow.

Siconolfi LB and Seeds NW (2001) Induction of the Plasminogen Activator System Accompanies Peripheral Nerve Regeneration after Sciatic Nerve Crush. *J Neurosci* **21(12)**:4336–4347.

Vleggeert-Lankamp CLAM (2007) The role of evaluation methods in the assessment of peripheral nerve regeneration through synthetic conduits: a systematic review laboratory investigation. *J Neurosurg* **107(6)**:1168-1189.

Wang H, Zhu H, Guo Q, Qian T, Zhang P, Li S, Xue C, and Gu X (2017) Overlapping Mechanisms of Peripheral Nerve Regeneration and Angiogenesis Following Sciatic Nerve Transection. *Front Cell Neurosci* **11**:323. doi: 10.3389/fncel.2017.00323.

Wertz A, Trenholm S, Yonehara K, Hillier D, Raics Z, Leinweber M, Szalay G, Ghanem A, Keller G, Rozsa B, Conzelmann KK, and Roska B (2015) Presynaptic Networks. Single-cell-initiated monosynaptic tracing reveals layer-

specific cortical network modules. *Science* **349(6243)**:70-74. doi:
10.1126/science.aab1687.

X-Gal Staining Protocol for beta-Galactosidase.
http://www.ihcworld.com/_protocols/special_stains/x_gal.htm.

Footnote

The study was supported by the Russian fund for basic research - project #18-015-00535. This study was conducted using biomaterial collected within the frame of Moscow University Project "Noah's Ark" and equipment purchased as a part of Moscow State University Program of Development.

Figure Legend

Fig. 1. Used plasmid variants encoding hBDNF and huPA.

Fig. 2. pNCure plasmid variants provide an acceptable level of hBDNF (A) and huPA (B) expression in culture of HEK293. Data are represented as mean \pm st. deviation. (n = 3; *P < 0.05).

Fig. 3. A conditioned media of HEK293 cells transfected with pVax1m-hBDNF and pNCure-fIntr plasmids stimulates neurite outgrowth in a Neuro2a cell model. On the pictures - Neuro2a cell cultures 72 h following neuritogenesis induction. A- negative control (DMEM 10% FBS), B- positive control (DMEM 1% FBS), C, D & E- conditioned media from HEK293 cells, transfected with pVax1m, pVax1m-hBDNF or pNCure-fIntr plasmids (100x). The graph plot demonstrates the percentage of cells with neurites in each group. Data are represented as mean \pm st. deviation. (n = 20; *P < 0.05 vs negative control, positive control and pVax1m group; **P < 0.05 vs negative control group).

Fig. 4. Multiple intramuscular injections of small volumes of plasmid solution (muscle chipping) provide the same amount of plasmid DNA delivery as a single injection accompanied by hydrodynamic shock (b-galactosidase staining) (100x, n = 3). Data are represented as mean \pm st. deviation.

Fig. 5. The pNCure-fIntr plasmid stimulates nerve conduction recovery: CNAP Amplitude (A), CNAP Latency (B). The maximum stimulating effect was observed after pNCure-fIntr injection at a dosage 120 ug/mouse. The average CNAP amplitude and latency of an intact nerve corresponds to the dashed lines.

Data are represented as median with 25% and 75% (percentile) (n = 10, *P < 0.05 and ***P < 0.001 vs control).

Fig. 6. The pNCure-fIntr plasmid stimulates nerve structure recovery. On the pictures – nerve stump cryosections 3 mm distally from the crush site (on the 18th day after crush) immunostained with axon-specific anti-NF200 antibody.

(50x). The graph plot demonstrates the number of NF200-positive structures per cryosection in each group. The average number of axons of an intact nerve corresponds to the dashed line. Data are represented as mean \pm st. deviation. (n > 5, *P < 0.05 vs control).

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5





FIGURE 6



A bicistronic plasmid encoding brain-derived neurotrophic factor and urokinase plasminogen activator stimulates peripheral nerve regeneration after injury


Maxim Karagyaur, Alexandra Rostovtseva, Ekaterina Semina, Polina Klimovich, Vadim Balabanyan, Pavel Makarevich, Vladimir Popov, Dmitry Stambolsky, Vsevolod Tkachuk

Table 1. Oligonucleotides used for pNCure variants plasmid construction


Name	Sequence
hBDNF-f	5'- GGAGGCTAGCCACCATGGCGACCATCCTTTTCCTTACTATGGTTATTTTCATACTTTGG
hBDNF-r	5'- GGAGAAGCTTTCATCTTCCCCTTTTAATGGTCAATGTACATACACAAGAAGTGTCTATC
huPA-f	5'- GGAGGGTACCAGAGCCCTGCTGGCGCGCCTGCTTCTCTGC
huPA-r	5'- GGAGACGGCCGCTCAGAGGGGCCAGGCCATTCTCTTCCTTGGTGTGAC
Intr-f1	5'- CTAGACTGGAGGTGAGTAACAGAGATCTAATACG
Intr-f2	5'- CTTCTCTTGCTAAGCCTAACCTTTTCTGTCTTGTTTCTGCTTTCCCCCCCCCACAGTTCG
Intr-r1	5'- CTAGCGAACTGTGGGGGGGGGAAAGCAGAAACAA
Intr-r2	5'- GACAGAAAAGGTTAGGCTTAGCAAGAGAAGCGTATTAGATCTCTGTTACTCACCTCCAGT
WPRE-f	5'- GGAGTCTAGACATCAACCTCTGGATTACAAAATTTGTGAAAGATTG
WPRE-r	5'- GGAGTCTAGAACAGGCGGGGAGGCGGCCCAAAGGGAG
IRES-f1	5'- GGAGAAGCTTGCCAGCCCCCGATTGGGGGCGCACTCCACCATAGATCACTCCCCTGTGAGGA ACTACTGTCTTCACGCAGAAAGCG
IRES-f2	5'- TCTAGCCATGGCGTTAGTATGAGTGTCGTGCAGCCTCCAGGACCCCCCTCCCGGGAGAGCCAT AGTGGTCTGCGGAACCGGTG

IRES-f3	5'- AGTACACCGGAATTGCCAGGACGACCGGGTCCTTTCTTGGATTAACCCGCTCAATGCCTGGAGA TTTGGGCGTGCCCCCGCGAGAC
IRES-f4	5'- TACTAGCCGAGTAGTGTTGGGTCGCGAAAGGCCTTGTGGTACTGCCTGATAGGGTGCTTGCGAG TGCCCCGGGAGGTCTCGTAGACCG
IRES-f5	5'- TGCATCATGAGCACAAATCCTAAACCTCAAGAAAAAGGTACCCTCC
IRES-r1	5'- GGAGGGTACCTTTTTCTTGAGGTTTAGGATTTGTGCTCATGATGCACGGTCTACGAGACCTCCC GGGGCACTCGCAAGCACCC
IRES-r2	5'- TATCAGGCAGTACCACAAGGCCTTTCGCGACCCAACACTACTCGGCTAGTAGTCTCGCGGGGGC ACGCCCAAATCTCCAGGCATTGAG
IRES-r3	5'- CGGGTTAATCCAAGAAAGGACCCGGTCGTCCTGGCAATTCCGGTGTACTCACCGGTTCCGCAGA CCACTATGGCTCTCCCGGGAGG
IRES-r4	5'- GGGGGTCCTGGAGGCTGCACGACACTCATACTAACGCCATGGCTAGACGCTTTCTGCGTGAAG ACAGTAGTTCCTCACAGGGGAGTG

IRES-r5	5' - ATCTATGGTGGAGTGTCGCCCAATCGGGGGCTGGCAAGCTTCTCC
---------	--



Suppl. Fig.1. Nerve injury model with a complex of function and structure assessing methods.



Suppl. Fig.2. Representative samples of CNAPs registered distally from the injury site in control and experimental groups on the 18th day after peripheral nerve injury.