

**Role of Efficacy as a Determinant of Locomotor Activation
by Mu Opioid Receptor Ligands in Female and Male Mice**

Edna J. Santos¹, Matthew L. Banks¹, and S. Stevens Negus¹

¹Department of Pharmacology and Toxicology, Virginia Commonwealth University,
Richmond VA 23298

RUNNING TITLE PAGE

Running title: Efficacy as a determinant of hyperlocomotion by MOR ligands

Corresponding Author: S. Stevens Negus
Department of Pharmacology and Toxicology
Virginia Commonwealth University
410 N. 12th Street
Richmond, Virginia. 23298
Phone: 804-828-3158
Fax: 804-828-2117
sidney.negus@vcuhealth.org

Text Pages:	17
Tables:	2
Figures:	6
References:	48
Abstract:	250
Introduction	610 (excluding references)
Discussion:	1465 (excluding references)

Nonstandard abbreviations:

CHO	Chinese hamster ovary
DMSO	dimethylsulfoxide
E _{max}	maximum effect
EP ₅₀	agonist proportion in an agonist/antagonist mixture that produces 50% agonist alone E _{max}
FENT/NTX	fentanyl/naltrexone
MOR	mu opioid receptor
NAQ	17-Cyclopropyl-methyl-3,14β-dihydroxy-4,5α-epoxy-6α- [(3'-isoquinolyl)acetamido]-morphinan
pFent	proportion of fentanyl in a fentanyl/naltrexone mixture

Recommended Section Assignment: Behavioral Pharmacology

ABSTRACT

Mu opioid receptor (MOR) agonists produce locomotor hyperactivity in mice as one sign of opioid-induced motor disruption. The goal of this study was to evaluate the degree of MOR efficacy required to produce this hyperactivity. Full dose-effect curves were determined for locomotor activation produced in male and female ICR mice by (1) eight different single-molecule opioids with high to low MOR efficacy, and (2) a series of fixed-proportion fentanyl/naltrexone mixtures with high to low fentanyl proportions. Data from the mixtures were used to quantify the efficacy requirement for MOR agonist-induced hyperactivity relative to efficacy requirements determined previously for other MOR agonist effects. Specifically, efficacy requirement was quantified as the EP_{50} value, which is the "Effective Proportion" of fentanyl in a fentanyl/naltrexone mixture that produces a maximal effect equal to 50% of the maximal effect of fentanyl alone. Maximal hyperactivity produced by each drug and mixture in the present study correlated with previously published data for maximal stimulation of GTP γ S binding in MOR-expressing Chinese hamster ovary cells as an *in vitro* measure of relative efficacy. Additionally, the EP_{50} value for hyperactivity induced by fentanyl/naltrexone mixtures indicated that opioid-induced hyperactivity in mice has a relatively high efficacy requirement in comparison to some other MOR agonist effects, and in particular is higher than the efficacy requirement for thermal antinociception in mice or fentanyl discrimination in rats. Taken together, these data show that MOR agonist-induced hyperactivity in mice is efficacy dependent and requires relatively high levels of MOR agonist efficacy for its full expression.

Significance Statement: Mu opioid receptor (MOR) agonist-induced hyperlocomotion in mice is dependent on the MOR efficacy of the agonist and requires a relatively high degree of efficacy for its full expression.

INTRODUCTION

Morphine and other mu opioid receptor (MOR) agonists produce a wide range of physiological and behavioral effects that include both therapeutically useful effects like analgesia and undesirable effects that include disrupted motor function (Yaksh and Wallace, 2018). One key determinant of the effects produced by any given MOR ligand is the relationship between two factors: (a) drug efficacy to activate MOR-coupled signaling mechanisms, and (b) efficacy requirements for different MOR-mediated effects (Selley et al., 2021). This relationship is illustrated in Figure 1. For any in vitro or in vivo test system, increasing drug doses will produce increasing MOR occupation and increasing levels of MOR activation. Maximal receptor activation plateaus at high doses that saturate receptors, and high-efficacy MOR ligands will produce higher plateaus than lower efficacy ligands. Within this framework of dose- and efficacy-dependent MOR activation, different effects require different thresholds of receptor activation for their expression and are constrained by different ceilings imposed by either experimental or biological limits. Together, the threshold and ceiling for a given effect define its efficacy requirement, which specifies the range of receptor-activation levels across which increasing doses will produce increasing effect. Moreover, the degree to which different effects have different efficacy requirements can be exploited in drug development, because low-efficacy drugs may have sufficient efficacy to produce some therapeutic effects but lack sufficient efficacy to produce some undesirable effects.

Mice are commonly used in preclinical drug development studies, and one MOR agonist effect in mice is a stimulation of horizontal locomotor activity (Frischknecht et al., 1983; Michael-Titus et al., 1989; Narita et al., 1993; Osborn et al., 2010; Raehal et al., 2005; Varshneya et al., 2019). Opioid-induced locomotor activation in mice is one manifestation of MOR-mediated motor disruption, and as such, it can be considered an undesirable opioid effect. Additionally, opioid-induced locomotor stimulation involves activation of the mesolimbic dopamine system

and serves as one behavioral consequence of enhanced mesolimbic dopamine signaling (Botz-Zapp et al., 2021; Chefer et al., 2003; Funada et al., 1993; Severino et al., 2020; Urs and Caron, 2014; Walters et al., 2005). Lastly, locomotor activation is an unconditioned behavioral effect that requires no prior training for its expression, and it can be continuously and quantitatively measured in commercially available locomotor-activity chambers (Chakraborty et al., 2021; Raehal et al., 2005; Varshneya et al., 2019). These features make opioid-induced locomotor activation useful as an endpoint for early evaluation of the in vivo potency, effectiveness, and time course of novel opioids. The utility of opioid-induced locomotor activation as a preclinical endpoint in drug evaluation would be further enhanced by clarification of its efficacy requirement relative to other opioid effects in mice and in other in vitro and in vivo test systems.

Accordingly, the goal of the present study was to evaluate the efficacy requirements for opioid-induced locomotor activation in male and female mice. Two parallel sets of studies were conducted. First, dose-effect curves were determined for a panel of eight MOR ligands with a range of maximal effects to stimulate GTP γ S binding as an in vitro measure of relative MOR efficacy (Selley et al., 1998; Thompson et al., 2004; Thomsen et al., 2014; Yuan et al., 2013). Second, dose-effect curves were also determined for a panel of drug mixtures composed of the high-efficacy MOR agonist fentanyl and antagonist naltrexone. We have shown previously the fixed proportion of fentanyl to naltrexone in fentanyl/naltrexone mixtures can be precisely manipulated to yield mixtures with graded maximal effects in both in vitro assays of ligand-stimulated GTP γ S binding and in vivo assays across multiple endpoints in multiple species of test subject (Cornelissen et al., 2018; Schwienteck et al., 2019; Selley et al., 2021). Additionally, fentanyl/naltrexone mixtures can be used to identify the effective proportion of fentanyl sufficient to produce 50% of the maximum effect of fentanyl alone (defined as the EP₅₀ value) as a metric of the efficacy requirement for any in vitro and in vivo MOR-mediated effect (Cornelissen et al., 2018; Schwienteck et al., 2019; Selley et al., 2021). Prevailing evidence suggests that locomotor activation may have a higher efficacy requirement than some other opioid effects

such as thermal antinociception in mice (Chakraborty et al., 2021; Varshneya et al., 2021, 2019). As a result, we predicted that the EP₅₀ value for locomotor activation in mice would be high relative to EP₅₀ values we have determined previously for other opioid effects in mice, rats, and rhesus monkeys.

METHODS

Animals

Subjects were male and female ICR mice (Envigo, Frederick, MD) that were 6–8 weeks old upon arrival to the laboratory. Males weighed 27-50 g and females weighed 22–50 g throughout the study. Mice were housed in same-sex, littermate groups in cages with corncob bedding (Envigo), a “nestlet” composed of pressed cotton (Ancare, Bellmore, NY), a cardboard tube for enrichment, and ad libitum access to food (Teklad LM-485 Mouse/Rat Diet; Envigo). Cages were mounted in a RAIR HD Ventilated Rack (Laboratory Products, Seaford, DE) in a temperature-controlled room with a 12-hour light/dark cycle (lights on from 6:00 AM to 6:00 PM) in a facility approved by the American Association for Accreditation of Laboratory Animal Care. All experiments were performed during the light phase of the daily light/dark cycle beginning 1 week after arrival at the laboratory. Ethical animal-use protocols were approved by the Virginia Commonwealth University Institutional Animal Care and Use Committee (Protocol #AD10001093) and complied with the National Research Council Guide for the Care and Use of Laboratory Animals.

Apparatus

Horizontal locomotor activity was assessed during 60-minute sessions in test boxes (16.8 x 12.7 cm² floor area x 12.7 cm high) housed in sound-attenuating chambers (Med Associates, St. Albans, VT) and located in a procedure room separate from the housing room.

Each box had black plexiglass walls, a clear plexiglass ceiling equipped with a house light, bar floors, and six photobeams arranged at 3-cm intervals across the long wall and 1 cm above the floor. Beam breaks were monitored by a microprocessor operating Med Associates software. The primary dependent variable was the total number of beam breaks, excluding consecutive interruptions of the same beam, during the 60-minute session.

Pharmacological Procedure

The primary goal of the study was to test a range of MOR ligands that varied in their relative efficacy at the MOR as quantified by maximum agonist-stimulated GTP γ S binding in Chinese hamster ovary (CHO) cells expressing the mouse or human MOR in previously published studies (Obeng et al., 2018; Selley et al., 2021, 1998; Thompson et al., 2004). This was accomplished by testing two different categories of treatments. First, a range of eight different single-molecule MOR ligands was tested. These drugs and their associated dose ranges were as follows (listed from highest to lowest E_{max} in studies of agonist-stimulated GTP γ S binding): methadone, 0.32 – 32 mg/kg (Middaugh and Zemp, 1976), fentanyl, 0.032 – 3.2 mg/kg (Varshneya et al., 2019), morphine, 1.0 – 100 mg/kg (Loggi et al., 1991), hydrocodone, 1.0 – 100 mg/kg (Jacob et al., 2017), buprenorphine, 0.01 – 3.2 mg/kg (Cowan et al., 1977), nalbuphine, 0.32 – 32 mg/kg (Patrick et al., 1999), NAQ (17-Cyclopropyl-methyl-3,14 β - dihydroxy-4,5 α -epoxy-6 α -[(3'-isoquinolyl)acetamido]-morphinan), 1.0 – 100 mg/kg (Zhang et al., 2014), and naltrexone, 0.1 – 3.2 mg/kg (Castellano and Puglisi-Allegra, 1982). Nalbuphine, NAQ, and naltrexone produced little or no locomotor stimulation across the dose-range tested, so each of these drugs was further evaluated for effectiveness to block locomotor activation by 10 mg/kg morphine.

The second category of treatments consisted of a series of fixed-proportion fentanyl/naltrexone mixtures. In these mixtures, the proportion of fentanyl to naltrexone was

fixed at a constant value for a given mixture (e.g. 1:1 fentanyl/naltrexone), and changes in the dose of one drug of the mixture were matched by equivalent changes in the other drug. We have reported previously that the net MOR efficacy of fentanyl/naltrexone mixtures can be precisely calibrated both in vitro and in vivo by adjusting the fentanyl proportion in the mixture, such that increasing fentanyl proportions result in increasing levels of MOR efficacy for the mixture. The present study compared effects of five different fentanyl/naltrexone mixtures ranging from 100:1 to 3.2:1 fentanyl/naltrexone.

With two exceptions noted below, a different group of 12 mice (six females, six males) were used to test each drug or mixture, and we have previously presented a detailed rationale for this group size and sex allocation (Diester et al., 2019). For this study, cohorts of up to 36 mice were generally used at any one time to test three different drugs or mixtures, and mice in each cohort were randomly assigned to the different treatments. Within each group, test sessions were conducted twice a week with at least 48 hours between sessions. All mice received a vehicle control and all doses of the designated test drug or mixture, and dose order was randomized across mice using a Latin-square design. The experimenter was not blinded to treatment because data collection was automated by the Med Associates software. There were no exclusion criteria, and all data were included in final analysis. On test days, mice were brought to the procedure room at least 2 hours before session onset. After subcutaneous test-drug administration, mice were returned to their home cages for the 5-minute pretreatment interval and then placed into the locomotor activity boxes at session onset. Doses for each drug or mixture were varied in 0.5 or 1.0 log-unit increments across a ≥ 10 -fold dose range with the intent of progressing from low doses that produced little or no effect to high doses that produced maximal increases in locomotor activation for that drug. For nalbuphine, NAQ, and naltrexone, antagonism studies were conducted after completion of drug-alone studies in the same mice. Doses of the test drug were administered 10 min before 10 mg/kg morphine, and locomotor

sessions began 5 min after morphine administration. There were two exceptions to this general design. First, in the case of hydrocodone, only six of the original mice (three of each sex) were tested at the high dose of 100 mg/kg due to limited drug supply. Because a clear effect plateau had not been reached at this dose, more drug was acquired and a higher dose of 320 mg/kg hydrocodone was tested in four other mice (two of each sex); however, all mice died, and further studies with hydrocodone were not pursued. Second, in the case of buprenorphine, the initial group was tested only up to a dose of 1.0 mg/kg due again to limited drug supply. Because a clear effect plateau had not been reached at this dose, more drug was acquired, and a higher dose of 3.2 mg/kg was tested in six other mice (three of each sex).

Data and Statistical Analysis

The primary dependent variable was the total number of beam breaks, excluding consecutive interruptions of the same beam, during each 60-minute session. These data were first analyzed within each drug or mixture to assess dose-dependent effects. Initial within-drug analysis proceeded in four phases as described previously for studies that include both females and males but are not intended a priori to detect sex differences (Diester et al., 2019). First, because sex was not the primary variable of interest, pooled data from both females and males were analyzed by repeated-measures one-way ANOVA with dose as the single variable, and a significant ANOVA was followed by a Holm-Sidak post-hoc test to both (a) identify doses producing effects different from vehicle and (b) evaluate presence or absence of a significant difference between the highest doses to identify an effect plateau for E_{\max} determination. For this and all other analyses described below, the criterion for significance was $p < 0.05$. Data for the highest doses of hydrocodone and buprenorphine were not included in the one-way ANOVA for these drugs because of the lower number of mice tested; rather, effects of these doses were compared to the next lower dose by t-test (paired for hydrocodone, unpaired for buprenorphine). Second, data were segregated by sex and again submitted to repeated-measures one-way

ANOVA followed by Holm-Sidak post-hoc test to assess dose-dependent effects within each sex. Third, male and female data were directly compared by two-way ANOVA with sex as a between-subjects factor and drug dose as a within-subjects factor. A significant sex \times dose interaction was followed by a Holm-Sidak post-hoc test. These first three steps of data analysis were performed using GraphPad Prism 9.0 (La Jolla, CA). Lastly, the two-way ANOVA results were submitted to power analyses to calculate the Cohen's f effect size, achieved power ($1 - \beta$), and the total number of animals predicted as necessary to detect a significant effect of sex, dose, and the sex \times dose interaction given the effect size, $\alpha = 0.05$, and power ($1 - \beta$) = 0.8 using the free statistical analysis program G*Power (Faul et al., 2007). Regarding the antagonism studies, data analysis was performed as described above in steps 1-3 with the exception that test drugs were evaluated for their effectiveness to decrease locomotor stimulant effects of morphine. Taken together, this strategy for experimental design and data analysis is intended to treat sex as an important but secondary variable of interest and to provide exploratory power analysis that can guide future studies explicitly designed to explore sex as a biological variable (Diester et al., 2019).

Following this within-drug analysis, three additional types of analyses were conducted. First, the maximal effects of each drug or fentanyl/naltrexone mixture at any dose were compared, and these E_{\max} values were considered to be different if 95% confidence limits did not overlap. Second, the E_{\max} of each drug or mixture for locomotor stimulation was transformed to a percentage of the fentanyl-alone E_{\max} (% Fent Max) using the equation $(\text{Test Drug } E_{\max} - \text{Vehicle Baseline}) / (\text{Fentanyl } E_{\max} - \text{Vehicle Baseline}) * 100$, where " E_{\max} " was the maximum number of locomotor counts for a test drug or fentanyl at any dose, and "Baseline" was the number of counts after vehicle treatment in that group. Values for % Fent Max of each drug and mixture were then graphed as a function of previously published E_{\max} values of each drug or

mixture to stimulate GTP γ S binding in CHO cells expressing cloned MOR (Obeng et al., 2018; Selley et al., 2021, 1998; Thompson et al., 2004). Data for single-molecule ligands and fentanyl/naltrexone mixtures were submitted separately to linear regression analyses for the linear sections of their respective curves to identify the magnitudes of GTP γ S binding (95% CL) associated with 50% Fent Max. Values were considered to be statistically similar if 95% confidence limits overlapped, and we predicted that these values would be similar for both single-molecule MOR ligands and fentanyl/naltrexone mixtures. Lastly, data for the fentanyl/naltrexone mixtures were used to determine an EP₅₀ value, defined as the proportion of fentanyl in the fentanyl/naltrexone mixture that produces an E_{max} equal to 50% of the fentanyl-alone E_{max}. As we have described previously, the EP₅₀ value determined from a series of fentanyl/naltrexone mixtures can be used to quantify the efficacy requirement for a given endpoint of MOR agonist-induced effects, such that higher EP₅₀ values indicate higher efficacy requirements. To calculate the EP₅₀ value, the E_{max} of each mixture was again expressed as % Fent Max and graphed as a function of the fentanyl proportion (pFent) for each mixture. These pFent-E_{max} data were submitted to nonlinear regression to determine the EP₅₀ (95% CL). This EP₅₀ value for locomotor activity in mice determined in the present study was then compared to previously determined EP₅₀ values for fentanyl/naltrexone mixtures to produce a range of other effects in previously published studies (Cornelissen et al., 2018; Schwienteck et al., 2019; Selley et al., 2021). EP₅₀ values across endpoints were considered to be different if 95% confidence limits did not overlap.

Materials

(\pm)Methadone HCl, fentanyl HCl, morphine sulfate, hydrocodone bitartrate, buprenorphine HCl, nalbuphine HCl, and naltrexone HCl were all provided by the National Institute on Drug Abuse Drug Supply Program. NAQ (17-Cyclopropyl-methyl-3,14 β - dihydroxy-4,5 α -epoxy-6 α -[(3'-isoquinolyl)acetamido]-morphinan) was synthesized by Dr. Yan Zhang. In

addition to these single-molecule test drugs, five fentanyl/naltrexone (FENT/NTX) mixtures were tested with fentanyl-to-naltrexone proportions of 100:1, 56:1, 32:1, 10:1, and 3.2:1. All compounds were administered subcutaneously (SC) per body weight in volumes of 0.1 – 0.9 ml and dissolved in sterile saline, except for NAQ which was dissolved in 10% DMSO and 90% water.

RESULTS

Figure 2 shows pooled data from both sexes for locomotor effects of all single-molecule opioids. One-way ANOVA results and E_{\max} values for each drug are shown in Table 1, and Figure 3 shows the time course of effects produced by selected doses of each drug over the 60-minute session. Methadone, fentanyl, morphine, hydrocodone, buprenorphine, and nalbuphine produced dose-dependent and significant locomotor stimulation, whereas NAQ and naltrexone did not. Each drug was tested up to an effect plateau at which increasing doses failed to produce further significant increases in locomotion. Note that, for hydrocodone, a higher dose of 320 mg/kg was tested in a subset of four male and female mice, and all died in ≤ 30 min. No dose of any other drug produced lethality in any other mice. E_{\max} values for methadone, fentanyl, morphine, hydrocodone, and buprenorphine were similar to each other and higher than for nalbuphine. Nalbuphine, NAQ, and naltrexone all produced a dose-dependent blockade of morphine-induced locomotor stimulation.

Figure 4 shows pooled data from both sexes for locomotor effects of the fentanyl/naltrexone mixtures. One-way ANOVA results and E_{\max} values for each mixture are shown in Table 2. The 100:1, 56:1, 32:1, and 10:1 fentanyl/naltrexone mixtures produced dose-dependent and significant increases in locomotor activity, whereas the 3.2:1 mixture did not. The mixture with the highest fentanyl proportion (100:1) produced the highest E_{\max} value, which

was not significantly different from the E_{\max} for fentanyl alone (see Table 1), and mixtures with progressively lower fentanyl proportions produced progressively lower E_{\max} values.

Results in Figures 2-4 indicate that, within boundaries described below, increasing MOR efficacy is associated with increasing locomotor activation in mice. The efficacy requirements for locomotor activation were quantified in two ways. First, Figure 5 shows the relationship between (a) the E_{\max} value of each single-molecule opioid and fentanyl/naltrexone mixture in the present study of locomotor activation and (b) the E_{\max} value in prior studies of ligand-stimulated GTP γ S binding in CHO cells expressing cloned MOR. Drugs or mixtures with in vitro E_{\max} values from 0% to approximately 50% of the DAMGO E_{\max} produced graded increases in locomotor activity; however, further increases in the in vitro E_{\max} values (with morphine, fentanyl, and methadone) did not produce further increases in locomotor activity. The mean (95% CL) magnitude of ligand-stimulated GTP γ S binding associated with a locomotor E_{\max} equal to 50% of the fentanyl-alone E_{\max} was similar for both single-molecule opioids [30.8 (25.1-37.2)] and fentanyl/naltrexone mixtures [29.2 (10.2-41.9)]. The mean (95% CL) slopes of the regressions were also similar [2.43 (1.66-3.21) for single-molecule opioids; 1.91 (0.81-3.01) for fentanyl/naltrexone mixtures].

Second, Figure 6 shows determination of the locomotor EP_{50} value, with EP_{50} value defined as the fentanyl proportion of the fentanyl/naltrexone mixture sufficient to produce an E_{\max} equal to 50% of the fentanyl-alone E_{\max} . Insofar as the EP_{50} value serves as a metric of the efficacy requirement for a given MOR-mediated effect, these results indicate that the efficacy requirement determined in the present study for locomotor activation in mice [EP_{50} (95% CL) = 18.6 (11.4-38.0)] is higher than in assays of opioid discrimination in rats or thermal antinociception in mice, similar to thermal antinociception in rats and rhesus monkeys, and lower than for stimulation of GTP γ S binding in CHO cells expressing cloned MOR.

Although the present study was not intended to rigorously evaluate sex differences in opioid effects, it did include both male and female subjects and did permit two-way dose x sex ANOVAs and subsequent post hoc power analysis for preliminary evaluation of sex as determinant of opioid effects. Results of these analyses are shown in Supplementary Table 1 (main effects of dose), Table 2 (main effects of sex) and Table 3 (dose x sex interaction), which show two-way ANOVA results, Cohen's effect size, current power, and projected sample size to achieve power ≥ 0.8 for all treatments. These analyses confirmed a main effect of dose for most single-molecule opioids and fentanyl/naltrexone mixtures, but not for NAQ, naltrexone, or the 3.2:1 fentanyl/naltrexone mixture. Main effects of sex or dose x sex interactions were rare, and in general, post-hoc power analysis indicated that power and associated sample sizes were too low to detect sex differences. Nonetheless, there were main effects of sex for the lowest two fentanyl/naltrexone mixtures (10:1 and 3.2:1) as shown in Supplementary Figure 1, with males showing higher locomotion regardless of dose, including after vehicle treatment. There was also a significant dose \times sex interaction for both the 32:1 fentanyl/naltrexone mixture and for hydrocodone, but for both treatments, post-hoc analysis did not identify a significant effect of sex at any dose of the mixture as shown in Supplementary Figure 2. Thus, even these significant sex effects provided weak evidence for a role of sex as a determinant of opioid-induced hyperactivity.

DISCUSSION

This study evaluated locomotor activation produced in mice by a panel of single-molecule opioids and fentanyl/naltrexone mixtures. There were three main findings. First, these results provide evidence for efficacy-dependent MOR agonist effects on maximal locomotor activation. This finding suggests that in vivo assessment of locomotor activity can serve as an efficient tool for in vivo stratification of the MOR efficacies of opioid ligands. Second, the apparent efficacy

requirement for locomotor activation was relatively high in comparison to previously determined efficacy requirements for other *in vivo* opioid effects in mice, such as antinociception. To the degree that locomotor activation in mice is an undesirable sign of opioid-induced motor disruption, these findings suggest the potential for low-efficacy MOR ligands to produce effects of potential therapeutic benefit (e.g. thermal antinociception) with minimal motor disruption. Lastly, the present results provided weak evidence for sex differences in opioid-induced locomotor stimulation, but when differences were observed, locomotor activity was higher in males. These results could provide a foundation for future efforts to explore sex differences in opioid-induced locomotor activation.

Efficacy dependence of MOR agonist-induced hyperactivity in mice.

MOR agonists produce locomotor activation in several strains of mice, including ICR mice (Bailey et al., 2010; Brase et al., 1977; Rethy et al., 1971; Szumlinski et al., 2020). This hyperactivity is expressed as continuous, unidirectional, and thigmotactic rotation around the perimeter of available space with reduced vertical activity (i.e. rearing, climbing) (Marcais-Collado et al., 1983; Michael-Titus et al., 1989; Mickley et al., 1989), and it can be viewed as a sign of adverse MOR-agonist induced motor disruption relative to other effects such as antinociception associated with therapeutic benefit.

The present study extends on these previous findings in its explicit examination of MOR efficacy as a determinant of MOR agonist-induced hyperactivity. Most single-molecule opioids and fentanyl/naltrexone mixtures produced dose-dependent increases in locomotion, and peak levels of activity across different drugs and mixtures were associated with peak levels of MOR-coupled G-protein signaling as measured by *in vitro* assays of ligand-stimulated GTP γ S binding in CHO cells expressing cloned MORs. This relationship was well described by a linear function up to a point, suggesting that MOR agonist-induced locomotor activation is mediated by MOR-

coupled G-protein signaling; however, drugs or mixtures that exceeded an in vitro E_{max} value of ~50% of the reference agonist DAMGO all produced similar E_{max} values for locomotor activation. These findings suggest that biological or procedural constraints impose a ceiling on maximal locomotor activation by high-efficacy MOR agonists. Conversely, no significant locomotor activation was produced by the low-efficacy MOR agonist NAQ or by the low-proportion 3.2:1 fentanyl/naltrexone mixture, both of which produce low but detectable levels of ligand-stimulated GTP γ S binding (Selley et al., 2021; Yuan et al., 2013). Taken together, these results show that MOR agonist-induced hyperactivity in mice is efficacy dependent, with graded E_{max} values within a range of low- to intermediate-efficacy agonists and a plateau of peak hyperactivity for high-efficacy agonists. Additionally, these results indicate that in vivo hyperactivity had a slightly higher efficacy threshold to detect agonist activity, a substantially lower ceiling, and a lower overall efficacy requirement than in vitro stimulation of GTP γ S binding for detection of MOR agonist effects.

This study examined efficacy dependence of MOR agonist-induced hyperactivity in mice using both single-molecule opioids and fentanyl/naltrexone mixtures. Results with the single-molecule opioids were suggestive of efficacy dependence, but the low-efficacy agonists nalbuphine and NAQ in this series have relatively low MOR selectivity (Pick et al., 1992; Yuan et al., 2011), and nalbuphine in particular produces agonist effects mediated by kappa opioid receptors (KOR) in mice (Narver, 2015; Patrick et al., 1999). Because KOR agonists decrease locomotor activity in mice (Gwynn and Domino, 1984; Kuzmin et al., 2000), it is possible that low locomotor activity with these drugs in general and nalbuphine in particular resulted from low selectivity for MOR vs. KOR rather than from low MOR efficacy. However, fentanyl/naltrexone mixtures with low fentanyl proportions also produced low peak levels of hyperactivity. With the mixtures, all agonist effects are produced by the highly MOR-selective opioid fentanyl, and net efficacy is controlled by the inclusion of naltrexone to block MORs and limit the maximal number

of receptors that can be occupied by fentanyl. Moreover, linear regression indicated that the magnitude of GTP γ S binding associated with an intermediate level of hyperactivity (50% of the E_{max} for fentanyl alone) was the same for single-molecule opioids and fentanyl/naltrexone mixtures. This suggests that low MOR efficacy is sufficient to explain the low levels of hyperactivity produced by nalbuphine and NAQ in this study, although any additional KOR agonist effects may also have contributed. Overall, the inclusion of data with fentanyl/naltrexone mixtures strengthens the conclusion of efficacy dependence for MOR agonist-induced hyperactivity in mice.

Efficacy requirements for MOR agonist-induced hyperactivity in mice relative to other *in vivo* effects.

The determination of dose-effect curves and E_{max} values for a range of fixed-proportion fentanyl/naltrexone mixtures provides a strategy to quantify efficacy requirements across opioid endpoints as the EP_{50} value, or the "Effective Proportion" of fentanyl in a fentanyl/naltrexone mixture required to produce an E_{max} equal to 50% of the fentanyl-alone E_{max} (Cornelissen et al., 2018; Schwienteck et al., 2019; Selley et al., 2021). For example, evidence cited above indicates that MOR agonist-induced hyperactivity in mice has a lower efficacy requirement than ligand-stimulated GTP γ S binding in MOR CHO cells, and this conclusion is further supported and quantified by reference to the EP_{50} values, with the EP_{50} (95%CL) being significantly lower for hyperactivity in mice than for GTP γ S binding in MOR CHO cells.

Two other general conclusions are suggested by a comparison of the present results with our previously published results (Cornelissen et al., 2018; Schwienteck et al., 2019; Selley et al., 2021). First, the EP_{50} for hyperactivity in mice was high relative to other *in vivo* behavioral endpoints in mice, rats, and monkeys, and in particular, was significantly higher than the EP_{50} from an assay of thermal nociception in mice. Insofar as opioid antinociception is related to a

therapeutic opioid effect (analgesia) whereas hyperactivity is related to an adverse effect (motor disruption), these results provide evidence for the potential of low-efficacy MOR agonists to produce analgesic effects without producing at least some degree of motor impairment. It should be noted that the EP_{50} for hyperactivity in mice was not significantly lower than that for thermal antinociception in rats or monkeys, suggesting that the window of opportunity here is narrow; nonetheless, these findings agree with other evidence to suggest that low-efficacy opioids can produce thermal antinociception without hyperactivity in mice (Varshneya et al., 2021, 2019). Second, the EP_{50} for hyperactivity in mice was also significantly higher than for a fentanyl discrimination assay in rats. Drug discrimination procedures model drug-induced subjective effects that may contribute to abuse potential, and as such, this finding suggests that abuse-related of MOR agonist effects have very low efficacy requirements. The fentanyl/naltrexone-mixture approach has not yet been applied to other endpoints of abuse-related opioid effects; however, evidence using other approaches to assess efficacy requirements (e.g. comparing low- and high-efficacy agonists or evaluating abuse-related effects after MOR downregulation with irreversible antagonists or genetic receptor knockdown) has also suggested that abuse-related MOR effects have relatively low efficacy requirements (Negus and Moerke, 2019; Sora et al., 2001; Zernig et al., 1997). Thus, although both hyperactivity in mice and rewarding/reinforcing effects of opioids in multiple species all appear to be mediated at least in part by mesolimbic dopamine signaling as a common neural substrate, it appears that lower MOR efficacy is required for behavioral reward/reinforcement processes than for unconditioned hyperactivity. One implication of these findings is that low-efficacy MOR agonists may produce little or no evidence of hyperactivity in mice but nonetheless produce rewarding/reinforcing effects sufficient to underlie abuse potential.

Sex differences in MOR agonist-induced hyperactivity.

This study focused on MOR efficacy as the principal independent variable and was neither intended nor powered to detect sex differences in opioid effects; however, the study did include both male and female mice and permitted preliminary assessment of sex as a determinant of opioid effects (Diester et al., 2019). In general, evidence for sex differences was weak and did not vary systematically as a function of MOR efficacy. Studies with most drugs and mixtures found only small effect sizes for the main effect of sex or the sex \times dose interaction, and post hoc power analysis indicated that most group sizes were underpowered to detect significance of any sex differences that might actually exist. In the cases where the main effect of sex or sex \times dose interaction was significant, locomotion tended to be higher in males, but this could not be attributed to higher sensitivity to opioid-induced hyperactivity because the sex \times dose interaction either was not significant or was not followed by a significant post hoc effect of sex at any dose. Previous studies also have found little evidence for sex differences in opioid-induced hyperactivity in mice. Main effects of sex have been observed suggestive of different baseline levels of activity, but the sex showing higher activity has varied (Collins et al., 2016; Kavaliers and Innes, 1987; Szumlinski et al., 2020). We could find only one study to show a significant sex \times dose interaction with a significant post hoc sex difference, with male deer mice showing higher activity than females during the light phase after treatment with 1 mg/kg morphine (Kavaliers and Innes, 1987). The power analysis of the present results could provide an empirical foundation for future studies explicitly designed to investigate sex as a determinant of opioid-induced hyperactivity in mice.

AUTHORSHIP CONTRIBUTIONS

Participated in research design: Santos, Banks, Negus

Conducted experiments: Santos

Performed data analysis: Santos, Banks, Negus

Wrote or contributed to writing of the manuscript: Santos, Banks, Negus

REFERENCES

- Bailey A, Metaxas A, Al-Hasani R, Keyworth HL, Forster DM, Kitchen I (2010) Mouse strain differences in locomotor, sensitisation and rewarding effect of heroin; association with alterations in MOP-r activation and dopamine transporter binding. *European Journal of Neuroscience* 31, 742–753. <https://doi.org/10.1111/j.1460-9568.2010.07104.x>
- Botz-Zapp CA, Foster SL, Pulley DM, Hempel B, Bi GH, Xi ZX, Newman AH, Weinschenker D, Manvich DF (2021) Effects of the selective dopamine D3 receptor antagonist PG01037 on morphine-induced hyperactivity and antinociception in mice. <https://doi.org/10.1101/2020.04.07.029918>
- Brase DA, Loh HH, Way EL (1977) Comparison of the effects of morphine on locomotor activity, analgesia and primary and protracted physical dependence in six mouse strains. *J Pharmacol Exp Ther* 201, 368–374.
- Castellano C, Puglisi-Allegra S (1982) Effects of naloxone and naltrexone on locomotor activity in C57BL/6 and DBA/2 mice. *Pharmacology Biochemistry and Behavior* 16, 561–563. [https://doi.org/10.1016/0091-3057\(82\)90415-4](https://doi.org/10.1016/0091-3057(82)90415-4)
- Chakraborty S, DiBerto JF, Faouzi A, Bernhard, SM, Gutridge AM, Ramsey S, Zhou Y, Provasi D, Nuthikattu N, Jilakara R, Nelson MNF, Asher WB, Eans SO, Wilson LL, Chintala SM, Filizola M, van Rijn RM, Margolis EB, Roth BL, McLaughlin JP, Che T, Sames, D, Javitch JA, Majumdar S (2021) A Novel Mitragynine Analog with Low-Efficacy Mu Opioid Receptor Agonism Displays Antinociception with Attenuated Adverse Effects. *J Med Chem* 64, 13873–13892. <https://doi.org/10.1021/acs.jmedchem.1c01273>

- Chefer VI, Kieffer BL, Shippenberg TS (2003) Basal and morphine-evoked dopaminergic neurotransmission in the nucleus accumbens of MOR- and DOR-knockout mice. *Eur J Neurosci* 18, 1915–1922. <https://doi.org/10.1046/j.1460-9568.2003.02912.x>
- Collins D, Reed B, Zhang Y, Kreek MJ (2016) Sex differences in responsiveness to the prescription opioid oxycodone in mice. *Pharmacology Biochemistry and Behavior* 148, 99–105. <https://doi.org/10.1016/j.pbb.2016.06.006>
- Cornelissen JC, Obeng S, Rice KC, Zhang Y, Negus SS, Banks ML (2018) Application of Receptor Theory to the Design and Use of Fixed-Proportion Mu-Opioid Agonist and Antagonist Mixtures in Rhesus Monkeys. *J Pharmacol Exp Ther* 365, 37–47. <https://doi.org/10.1124/jpet.117.246439>
- Cowan A, Lewis JW, Macfarlane IR (1977) Agonist and Antagonist Properties of Buprenorphine, a New Antinociceptive Agent. *British Journal of Pharmacology* 60, 537–545. <https://doi.org/10.1111/j.1476-5381.1977.tb07532.x>
- Diester CM, Banks ML, Neigh GN, Negus SS (2019) Experimental design and analysis for consideration of sex as a biological variable. *Neuropsychopharmacol.* 44, 2159–2162. <https://doi.org/10.1038/s41386-019-0458-9>
- Faul F, Erdfelder E, Lang, AG, Buchner A, (2007) G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods* 39, 175–191. <https://doi.org/10.3758/BF03193146>
- Frischknecht HR, Siegfried B, Riggio G, Waser PG, (1983) Inhibition of morphine-induced analgesia and locomotor activity in strains of mice: A comparison of long-acting opiate

antagonists. *Pharmacology Biochemistry and Behavior* 19, 939–944.

[https://doi.org/10.1016/0091-3057\(83\)90395-7](https://doi.org/10.1016/0091-3057(83)90395-7)

Funada M, Suzuki T, Narita M, Misawa M, Nagase H, (1993) Modification of morphine-induced locomotor activity by pertussis toxin: biochemical and behavioral studies in mice. *Brain Res* 619, 163–172. [https://doi.org/10.1016/0006-8993\(93\)91608-u](https://doi.org/10.1016/0006-8993(93)91608-u)

Gwynn J Domino F (1984) Genotype-Dependent Behavioral Sensitivity to Mu vs. Kappa Opiate Agonists. I. Acute and Chronic Effects on Mouse Locomotor Activity 231, 6.

Jacob JC, Poklis JL, Akbarali HI, Henderson G, Dewey WL (2017) Ethanol Reversal of Tolerance to the Antinociceptive Effects of Oxycodone and Hydrocodone. *J Pharmacol Exp Ther* 362, 45–52. <https://doi.org/10.1124/jpet.117.241083>

Kavaliers M, Innes DGL (1987) Sex and day-night differences in opiate-induced responses of insular wild deer mice, *Peromyscus maniculatus triangularis*. *Pharmacology Biochemistry and Behavior* 27, 477–482. [https://doi.org/10.1016/0091-3057\(87\)90351-0](https://doi.org/10.1016/0091-3057(87)90351-0)

Kuzmin A, Sandin J, Terenius L, Gren SOO, (2000) Dose- and Time-Dependent Bimodal Effects of μ -Opioid Agonists on Locomotor Activity in Mice 295, 12.

Loggi G, Laviola G, Alleva E, Chiarotti F, (1991) Morphine effects on mouse locomotor/exploratory activity: Test dependency, test reliability, uni- and multi-variate analyses. *Pharmacology Biochemistry and Behavior* 38, 817–822.
[https://doi.org/10.1016/0091-3057\(91\)90248-Z](https://doi.org/10.1016/0091-3057(91)90248-Z)

Marcas-Collado H, Chaillet P, Costentin J (1983) Inhibition of the spontaneous climbing behavior elicited in mice by opiates. *J Pharmacol Exp Ther* 227, 466–471.

- Michael-Titus A, Dourmap N, Costentin J (1989) MU and delta opioid receptors control differently the horizontal and vertical components of locomotor activity in mice. *Neuropeptides* 13, 235–242. [https://doi.org/10.1016/0143-4179\(89\)90076-0](https://doi.org/10.1016/0143-4179(89)90076-0)
- Mickley GA, Mulvihill MA, Postler MA (1989) Brain μ and δ opioid receptors mediate different locomotor hyperactivity responses of the C57BL/6J mouse. *6*.
- Middaugh LD, Zemp JW (1976) Effects of Methadone on Activity and on Brain Monoamines in Two Strains of Mice I. *Pharmacology Biochemistry & Behavior* 5, 367–370.
- Narita M, Takahashi Y, Takamori K, Funada M, Suzuki T, Misawa M, Nagase H (1993) Effects of κ -Agonist on the Antinociception and Locomotor Enhancing Action Induced by Morphine in Mice. *The Japanese Journal of Pharmacology* 62, 15–24.
<https://doi.org/10.1254/jjp.62.15>
- Narver HL (2015) Nalbuphine, a non-controlled opioid analgesic, and its potential use in research mice. *Lab Animal* 44, 106–110. <https://doi.org/10.1038/lab.701>
- Negus SS, Moerke MJ, 2019. Determinants of opioid abuse potential: Insights using intracranial self-stimulation. *Peptides* 112, 23–31.
- Obeng S, Yuan Y, Jali A, Selley DE, Zhang Y (2018) In vitro and in vivo functional profile characterization of 17-cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(isoquinoline-3-carboxamido)morphinan (NAQ) as a low efficacy μ opioid receptor modulator. *Eur J Pharmacol* 827, 32–40. <https://doi.org/10.1016/j.ejphar.2018.03.013>
- Osborn MD, Lowery JJ, Skorput AGJ, Giuvelis D, Bilsky EJ (2010) In vivo characterization of the opioid antagonist nalmefene in mice. *Life Sci* 86, 624–630.
<https://doi.org/10.1016/j.lfs.2010.02.013>

Patrick CA, Ko MCH, Woods JH (1999) Comparison of Antinociceptive Effects Induced by Kappa Opioid Agonists in Male and Female Mice. *analgesia (elmsford n y)* 4, 397–404.

<https://doi.org/10.3727/107156999819565766>

Pick C, Paul D, Pasternak GW (1992) Nalbuphine, a mixed kappa 1 and kappa 3 analgesic in Mice. *The Journal of Pharmacology and Experimental Therapeutics* 262, 1044–1050.

Raehal KM, Lowery JJ, Bhamidipati CM, Paolino RM, Blair JR, Wang D, Sadée W, Bilsky EJ (2005)

In vivo characterization of 6beta-naltrexol, an opioid ligand with less inverse agonist activity compared with naltrexone and naloxone in opioid-dependent mice. *J Pharmacol Exp Ther* 313, 1150–1162.

<https://doi.org/10.1124/jpet.104.082966>

Rethy CR, Smith CB, Villarreal JE (1971) Effects of narcotic analgesics upon the loco motor activity and brain catecholamine content of the mouse. *Journal of Pharmacology and Experimental Therapeutics* 176, 472–479.

Schwienteck KL, Faunce KE, Rice KC, Obeng S, Zhang Y, Blough BE, Grim TW, Negus SS, Banks ML (2019) Effectiveness comparisons of G-protein biased and unbiased mu opioid receptor ligands in warm water tail-withdrawal and drug discrimination in male and female rats.

Neuropharmacology 150, 200–209. <https://doi.org/10.1016/j.neuropharm.2019.01.020>

Selley DE, Banks ML, Diester CM, Jali AM, Legakis LP, Santos EJ, Negus SS (2021) Manipulating Pharmacodynamic Efficacy with Agonist + Antagonist Mixtures: In Vitro and In Vivo Studies with Opioids and Cannabinoids. *J Pharmacol Exp Ther* 376, 374–384.

<https://doi.org/10.1124/jpet.120.000349>

- Selley DE, Liu Q, Childers SR (1998) Signal Transduction Correlates of Mu Opioid Agonist Intrinsic Efficacy: Receptor-Stimulated [35S]GTP γ S Binding in mMOR-CHO Cells and Rat Thalamus 285, 10.
- Severino AL, Mittal N, Hakimian JK, Velarde N, Minasyan A, Albert R, Torres C, Romaneschi N, Johnston C, Tiwari S, Lee AS, Taylor AM, Gavériaux-Ruff C, Kieffer BL, Evans CJ, Cahill CM, Walwyn WM (2020) μ -Opioid Receptors on Distinct Neuronal Populations Mediate Different Aspects of Opioid Reward-Related Behaviors. eNeuro 7, ENEURO.0146-20.2020. <https://doi.org/10.1523/ENEURO.0146-20.2020>
- Sora I, Elmer G, Funada M, Pieper J, Li XF, Hall S, Uhl GR (2001) μ Opiate Receptor Gene Dose Effects on Different Morphine Actions: Evidence for Differential in vivo μ Receptor Reserve. Neuropsychopharmacology 25, 14.
- Szumliński KK, Coelho MA, Tran T, Stailey N, Lieberman D, Gabriella I, Swauncy I, Brewin LW, Ferdousian S (2020) Who is HOT and who is LOT? Detailed characterization of prescription opioid-induced changes in behavior between 129P3/J and 129S1/SvImJ mouse substrains. Genes, Brain and Behavior 19, e12609. <https://doi.org/10.1111/gbb.12609>
- Thompson CM, Wojno H, Greiner E, May EL, Rice KC, Selley DE (2004) Activation of G-Proteins by Morphine and Codeine Congeners: Insights to the Relevance of *O* - and *N* - Demethylated Metabolites at μ - and δ -Opioid Receptors. J Pharmacol Exp Ther 308, 547–554. <https://doi.org/10.1124/jpet.103.058602>

- Thomsen M, Fulton BS, Caine SB (2014) Acute and chronic effects of the M1/M4-preferring muscarinic agonist xanomeline on cocaine vs. food choice in rats. *Psychopharmacology (Berl)* 231, 469–479. <https://doi.org/10.1007/s00213-013-3256-9>
- Urs NM, Caron MG (2014) The physiological relevance of functional selectivity in dopamine signalling. *Int J Obes Suppl* 4, S5-8. <https://doi.org/10.1038/ijosup.2014.3>
- Varshneya NB, Walentiny DM, Moisa LT, Walker TD, Akinfiresoye LR, Beardsley PM (2021) Fentanyl-related substances elicit antinociception and hyperlocomotion in mice via opioid receptors. *Pharmacol Biochem Behav* 208, 173242. <https://doi.org/10.1016/j.pbb.2021.173242>
- Varshneya NB, Walentiny DM, Moisa LT, Walker TD, Akinfiresoye LR, Beardsley PM (2019) Opioid-like antinociceptive and locomotor effects of emerging fentanyl-related substances. *Neuropharmacology* 151, 171–179. <https://doi.org/10.1016/j.neuropharm.2019.03.023>
- Walters CL, Godfrey M, Li X, Blendy JA (2005) Alterations in morphine-induced reward, locomotor activity, and thermoregulation in CREB-deficient mice. *Brain Res* 1032, 193–199. <https://doi.org/10.1016/j.brainres.2004.11.013>
- Yaksh and Wallace (2018) Opioids, analgesia, and pain management, in Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 13e (Brunton LL, Hilal-Dandan R and Knollmann BC eds) pp 355-386, McGraw-Hill, New York.
- Yuan Y, Elbegdorj O, Beletskaya IO, Selley DE, Zhang Y (2013) Structure activity relationship studies of 17-cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(isoquinoline-3'-carboxamido)morphinan (NAQ) analogues as potent opioid receptor ligands: Preliminary

results on the role of electronic characteristics for affinity and function. *Bioorganic & Medicinal Chemistry Letters* 23, 5045–5048.

<https://doi.org/10.1016/j.bmcl.2013.07.043>

Yuan Y, Li G, He H, Stevens DL, Kozak P, Scoggins KL, Mitra P, Gerk PM, Selley DE, Dewey WL, Zhang Y (2011) Characterization of 6 α - and 6 β - *N*-Heterocyclic Substituted Naltrexamine Derivatives as Novel Leads to Development of Mu Opioid Receptor Selective Antagonists. *ACS Chem. Neurosci.* 2, 346–351. <https://doi.org/10.1021/cn2000348>

Zernig G, Lewis JW, Woods JH (1997) Cloccinamox inhibits the intravenous self-administration of opioid agonists in rhesus monkeys: comparison with effects on opioid agonist-mediated antinociception. *Psychopharmacology* 129, 233–242.
<https://doi.org/10.1007/s002130050185>

Zhang Y, Braithwaite A, Yuan Y, Streicher JM, Bilsky EJ, (2014) Behavioral and cellular pharmacology characterization of 17-cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(isoquinoline-3'-carboxamido)morphinan (NAQ) as a mu opioid receptor selective ligand. *European Journal of Pharmacology* 736, 124–130.
<https://doi.org/10.1016/j.ejphar.2014.04.041>

FOOTNOTE

Supported by grants P30DA033934, R25GM090084, and T32DA007027 from the National Institutes of Health. No author has an actual or perceived conflict of interest with the contents of this article.

FIGURE LEGENDS

Figure 1. Dose and efficacy of the drug and efficacy requirement of the effect as determinants of drug effects. Drugs produce their effects by acting on a population of target receptors in the biological system to which they are administered. For drugs with both affinity and efficacy at the target receptor, receptor theory predicts that increasing drug doses will produce increasing levels of receptor occupancy and activation. (A) Peak levels of total receptor activation are determined by drug efficacy, such that drugs with higher efficacy will produce higher plateau levels of receptor activation than drugs with lower efficacy. Abscissa: Drug dose in arbitrary units. Ordinate: % Total Possible Activation of all available receptors. (B) Different effects mediated by the receptor require different levels of receptor activation for their expression. Effects with a low efficacy requirement (Effect 1) require low levels of receptor activation to surpass the threshold and reach the ceiling for their expression. Even low-efficacy drugs have sufficient efficacy to produce these effects. Effects with a high efficacy requirement (Effect 2) require higher levels of receptor activation to surpass the threshold and reach the ceiling for their expression. High-efficacy drugs can produce sufficient receptor activation to produce these effects, although this will require higher doses than for effects with low efficacy requirements. Lower efficacy drugs may have sufficient efficacy to surpass the threshold but not reach the ceiling and may therefore function as partial agonists for these effects. Even lower efficacy drugs may lack sufficient efficacy even to reach the threshold, in which case they will function as antagonists.

Figure 2. Locomotor activating effects of opioids with differing MOR efficacy. (A) Effects of opioids administered alone. (B) Effects of nalbuphine, NAQ and naltrexone administered as a pretreatment to 10 mg/kg morphine. Abscissae: Dose in mg/kg. Ordinates: Locomotor counts per 60 minutes. In general, all points show mean \pm SEM for N=12 mice, and filled symbols indicate a significant difference compared to vehicle within each drug as determined by repeated-measures one-way ANOVA followed by the Holm-Sidak post hoc test, $p < 0.05$. There were two exceptions. The dashed line to 3.2 mg/kg buprenorphine indicates low sample size

(N=6) and a different cohort of mice for this dose, and the filled point indicates different from vehicle by unpaired t-test. The dashed line at 100 mg/kg hydrocodone indicates low sample size (N=6) but in the same cohort of mice for this dose, and the filled symbol indicates different from vehicle by paired t-test. A different group of four male and female mice tested with a higher hydrocodone dose (320 mg/kg) all died, so further studies at this dose were not conducted and these data are not included in the graph. Statistical results for Panel A are shown in Table 1. For Panel B one-way ANOVA results were as follows. Nalbuphine: $F(1.64, 18.02) = 14.42$; $p=0.0003$; NAQ: $F(4.15, 45.68) = 8.67$; $p<0.0001$; Naltrexone: $F(2.59, 28.45) = 27.35$; $p<0.0001$.

Figure 3. Time course of locomotor activating effects produced by opioids with differing MOR efficacy. Each panel shows time course data over a 60-minute session for a different drug. For most drugs, data are shown for vehicle, the lowest dose to significantly increase locomotion, and the E_{max} dose producing the highest level of locomotor activation. Nalbuphine, NAQ, and naltrexone show only data for vehicle and the highest dose tested. Abscissae: Time in min for the 60-min session. Ordinates: movement counts over the 60-minute session. Each point shows mean \pm SEM from 12 mice except the high dose for hydrocodone, which show N=6.

Figure 4. Locomotor activating effects of fentanyl/naltrexone mixtures. Abscissa: Dose fentanyl in mg/kg. The naltrexone dose was proportional to the fentanyl dose as indicated by fixed fentanyl/naltrexone (FENT/NTX) proportions for each mixture. Ordinate: Locomotor counts per 60 minutes. All points show mean \pm SEM for N=12 mice, and filled symbols indicate a significant difference compared to vehicle within each mixture as determined by repeated-measures one-way ANOVA followed by the Holm-Sidak post hoc test, $p<0.05$. Statistical results are shown in Table 2.

Figure 5. Relationship between MOR agonist and fentanyl/naltrexone mixture effects on in vitro activation of GTP γ S binding and in vivo locomotor stimulation. Abscissa: E_{\max} for each drug or mixture to stimulate GTP γ S binding in CHO cells expressing cloned MOR from previously published studies (see text for citations). Data are expressed as a percentage of the maximum effect of the high-efficacy MOR agonist DAMGO, which was included as a standard in each study. Ordinate: E_{\max} for each drug or mixture to stimulate locomotor activity in the present study. Data are expressed as a percentage of the maximum effect produced by fentanyl. The blue dotted line shows linear regression for single-molecule opioids on the linear portion of the curve (morphine, fentanyl, and methadone excluded). The gray solid line shows linear regression for the fentanyl/naltrexone mixtures. Abbreviations: METHD, methadone; FENT, fentanyl; MORPH, morphine; HYD, hydrocodone; BUP, buprenorphine; NALB, nalbuphine; NAQ, NAQ; NTX, naltrexone.

Figure 6. EP_{50} values as a metric of efficacy requirement for different effects produced by fentanyl/naltrexone mixtures. (A) Abscissa: Fentanyl proportion in different fentanyl/naltrexone mixtures. Ordinate: Maximum locomotor activating effects of each mixture expressed as a percentage of the fentanyl-alone maximum. Each point shows mean \pm SEM for 12 mice, and nonlinear regression was used to calculate the EP_{50} , which is defined as the fentanyl proportion in a fentanyl/naltrexone mixture that would produce a maximum effect equal to 50% of the fentanyl-alone maximum effect. (B) EP_{50} value (95% CL) for locomotor activation in the present study relative to EP_{50} values for fentanyl/naltrexone mixtures determined in previous studies using various behavioral endpoints in mice, rats, and rhesus monkeys or in the in vitro assay of ligand-stimulated GTP γ S binding in CHO cells expressing cloned MOR. Assays with EP_{50} values to the left of the shaded box have lower efficacy requirements than locomotor activation, whereas points to the right of the shaded box have higher efficacy

requirements than locomotor activation. Abbreviations: Drug Discrim, drug discrimination; TW, warm-water tail-withdrawal with water temperature specified in °C.

TABLES

Table 1. One-way ANOVA results and E_{max} values for data shown in Figure 2A.

Opioid	One-way ANOVA	E _{max} (95% CI)
Methadone	F(2.89, 31.87) = 38.81; p<0.0001	7500 (6505, 8495)
Fentanyl	F(2.97, 32.63) = 26.94; p<0.0001	7393 (6346, 8440)
Morphine	F(2.76, 30.40) = 31.75; p<0.0001	6925 (5662, 8188)
Hydrocodone	F(2.08, 22.86) = 13.81; p=0.0001	6153 (5025, 7280)
Buprenorphine	F(2.58, 28.34) = 39.39; p<0.0001	6867 (5802, 7933)
Nalbuphine	F(3.45, 37.90) = 3.29; p=0.0254	2639 (1954, 3324)
NAQ	F(2.52, 27.67) = 2.06; p=0.1376	-
Naltrexone	F(2.41, 26.46) = 1.69; p=0.2001	-


*Hydrocodone and buprenorphine data in this table does not include high doses because different N

Table 2. One-way ANOVA results and E_{max} values for fentanyl/naltrexone mixtures in Figure 4.

Fentanyl/naltrexone mixture	One-way ANOVA	E_{max} (95% CI)
100:1	F(2.39, 26.37) = 40.84; p<0.0001	8615 (7386, 9843)
56:1	F(2.16, 23.79) = 36.28; p<0.0001	6777 (5805, 7748)
32:1	F(3.27, 35.94) = 14.06; p<0.0001	5458 (4854, 6061)
10:1	F(2.68, 29.43) = 6.60; p=0.0021	4338 (3033, 5643)
3.2:1	F(3.82, 41.99) = 0.76; p=0.5506	-

Figure 1

(A) Theoretical Curves Relating Dose & Efficacy to Total Receptor Activation



(B) Endpoints with Different Efficacy Requirements






Figure 2

Figure 3




Figure 4




Figure 5





Figure 6



**Role of Efficacy as a Determinant of Locomotor Activation
by Mu Opioid Receptor Ligands in Female and Male Mice**

Edna J. Santos¹, Matthew L. Banks¹, and S. Stevens Negus¹

¹Department of Pharmacology and Toxicology, Virginia Commonwealth University,
Richmond VA 23298

Journal title: Journal of Pharmacology and Experimental Therapeutics
Manuscript number: JPET-AR-2021-001045

SUPPORTING INFORMATION

Supplementary Table 1. Two-way ANOVA results with power analysis of main effect of dose.

Opioid/Mixture	F statistic; p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power \geq 0.8
Methadone	F(2.820, 28.20) = 38.75; p<0.0001	1.965	1.0	10
Fentanyl	F(3.337, 33.37) = 29.25; p<0.0001	1.710	0.975	10
Morphine	F(2.715, 27.15) = 29.89; p<0.0001	1.728	0.953	10
Hydrocodone	F(2.757, 27.57) = 18.95; p<0.0001	1.376	0.896	10
Buprenorphine	F(2.627, 26.27) = 38.49; p<0.0001	1.962	0.984	10
Nalbuphine	F(3.358, 33.58) = 3.18; p=0.032	0.564	0.205	55
NAQ	F(2.489, 24.89) = 1.92; p=0.159	0.439	0.144	90
Naltrexone	F(2.356, 23.56) = 1.56; p=0.229	0.396	0.122	>100
100:1	F(2.614, 26.14) = 44.61; p<0.0001	2.112	0.993	10
56:1	F(2.154, 21.54) = 33.20; p<0.0001	1.822	0.936	10
32:1	F(3.036, 30.36) = 16.39; p,0.0001	1.280	0.781	15
10:1	F(2.821, 28.21) = 6.856 p=0.002	0.828	0.373	30
3.2:1	F(3.515, 35.15) = 0.75; p=0.551	0.273	0.081	>100

Supplementary Table 2. Two-way ANOVA results with power analysis of main effect of sex.

Opioid/Mixture	F statistic; p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power≥0.8
Methadone	F(1, 10) = 0.42; p=0.534	0.124	0.068	>100
Fentanyl	F(1, 10) = 0.00; p=0.952	0.010	0.050	>100
Morphine	F(1, 10) = 1.77; p=0.213	0.299	0.156	90
Hydrocodone	F(1, 10) = 0.20; p=0.665	0.084	0.058	>100
Buprenorphine	F(1, 10) = 0.85; p=0.379	0.212	0.102	>100
Nalbuphine	F(1, 10) = 0.48; p=0.506	0.212	0.102	>100
NAQ	F(1, 10) = 0.04; p=0.845	0.055	0.053	>100
Naltrexone	F(1, 10) = 0.02; p=0.897	0.027	0.051	>100
100:1	F(1, 10) = 1.41; p=0.263	0.320	0.172	80
56:1	F(1, 10) = 0.00; p=0.977	0.007	0.050	>100
32:1	F(1, 10) = 0.96; p=0.351	0.359	0.203	65
10:1	F(1, 10) = 7.15; p=0.023	1.198	0.962	10
3.2:1	F(1, 10) = 12.66; p=0.005	1.073	0.917	10

Supplementary Table 3. Two-way ANOVA results with power analysis of dose x sex interaction.

Opioid/Mixture	F statistic, p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power≥0.8
Methadone	F(4, 40) = 0.98; p=0.428	0.313	0.282	35
Fentanyl	F(5, 50) = 1.94; p=0.104	0.441	0.607	20
Morphine	F(5, 50) = 0.36; p=0.875	0.189	0.133	75
Hydrocodone	F(4, 40) = 5.09; p=0.002	0.713	0.944	10
Buprenorphine	F(5, 50) = 0.75; p=0.589	0.274	0.248	40
Nalbuphine	F(5, 50) = 0.62; p=0.689	0.248	0.206	45
NAQ	F(4, 40) = 0.30; p=0.877	0.173	0.110	>100
Naltrexone	F(4, 40) = 0.18; p=0.947	0.135	0.085	>100
100:1	F(5, 50) = 2.02; p=0.093	0.449	0.626	20
56:1	F(5, 50) = 0.07; p=0.997	0.081	0.063	>100
32:1	F(5, 50) = 2.82; p=0.025	0.531	0.794	15
10:1	F(5, 50) = 1.43; p=0.231	0.378	0.460	25
3.2:1	F(5, 50) = 0.79; p=0.564	0.280	0.259	40

Supplementary Figure 1. Significant main effect of sex.

Figure 1a





Figure 1b



Supplementary Figure 1. Treatments for which there was a main effect of sex in statistical analysis. Abscissae: Dose fentanyl in mg/kg (VEH=vehicle). The naltrexone dose = fentanyl dose \div fentanyl proportion (10 in 1a, 3.2 in 1b). Ordinates: Locomotor counts per 60 minutes. Male and female data are shown in green and orange bars, respectively, and individual data are shown by points. Males had higher locomotor scores than females for these two treatments.

Supplementary Figure 2. Significant dose x sex interaction.

Figure 2a





Figure 2b



Supplemental Figure 2. Treatments for which there was a sex x dose interaction. Abscissae: Dose in mg/kg (VEH=vehicle). For panel 2a, the abscissa shows fentanyl dose, and the naltrexone dose = fentanyl dose ÷ fentanyl proportion (32). Ordinates: Locomotor counts per 60 minutes. Male and female data are shown in green and orange bars, respectively, and individual data are shown by points. Post hoc analysis did not indicate a significant difference between males and females at any dose of either treatment.

**Role of Efficacy as a Determinant of Locomotor Activation
by Mu Opioid Receptor Ligands in Female and Male Mice**

Edna J. Santos¹, Matthew L. Banks¹, and S. Stevens Negus¹

¹Department of Pharmacology and Toxicology, Virginia Commonwealth University,
Richmond VA 23298

Journal title: Journal of Pharmacology and Experimental Therapeutics
Manuscript number: JPET-AR-2021-001045

SUPPORTING INFORMATION

Supplementary Table 1. Two-way ANOVA results with power analysis of main effect of dose.

Opioid/Mixture	F statistic; p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power \geq 0.8
Methadone	F(2.820, 28.20) = 38.75; p<0.0001	1.965	1.0	10
Fentanyl	F(3.337, 33.37) = 29.25; p<0.0001	1.710	0.975	10
Morphine	F(2.715, 27.15) = 29.89; p<0.0001	1.728	0.953	10
Hydrocodone	F(2.757, 27.57) = 18.95; p<0.0001	1.376	0.896	10
Buprenorphine	F(2.627, 26.27) = 38.49; p<0.0001	1.962	0.984	10
Nalbuphine	F(3.358, 33.58) = 3.18; p=0.032	0.564	0.205	55
NAQ	F(2.489, 24.89) = 1.92; p=0.159	0.439	0.144	90
Naltrexone	F(2.356, 23.56) = 1.56; p=0.229	0.396	0.122	>100
100:1	F(2.614, 26.14) = 44.61; p<0.0001	2.112	0.993	10
56:1	F(2.154, 21.54) = 33.20; p<0.0001	1.822	0.936	10
32:1	F(3.036, 30.36) = 16.39; p,0.0001	1.280	0.781	15
10:1	F(2.821, 28.21) = 6.856 p=0.002	0.828	0.373	30
3.2:1	F(3.515, 35.15) = 0.75; p=0.551	0.273	0.081	>100

Supplementary Table 2. Two-way ANOVA results with power analysis of main effect of sex.

Opioid/Mixture	F statistic; p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power≥0.8
Methadone	F(1, 10) = 0.42; p=0.534	0.124	0.068	>100
Fentanyl	F(1, 10) = 0.00; p=0.952	0.010	0.050	>100
Morphine	F(1, 10) = 1.77; p=0.213	0.299	0.156	90
Hydrocodone	F(1, 10) = 0.20; p=0.665	0.084	0.058	>100
Buprenorphine	F(1, 10) = 0.85; p=0.379	0.212	0.102	>100
Nalbuphine	F(1, 10) = 0.48; p=0.506	0.212	0.102	>100
NAQ	F(1, 10) = 0.04; p=0.845	0.055	0.053	>100
Naltrexone	F(1, 10) = 0.02; p=0.897	0.027	0.051	>100
100:1	F(1, 10) = 1.41; p=0.263	0.320	0.172	80
56:1	F(1, 10) = 0.00; p=0.977	0.007	0.050	>100
32:1	F(1, 10) = 0.96; p=0.351	0.359	0.203	65
10:1	F(1, 10) = 7.15; p=0.023	1.198	0.962	10
3.2:1	F(1, 10) = 12.66; p=0.005	1.073	0.917	10

Supplementary Table 3. Two-way ANOVA results with power analysis of dose x sex interaction.

Opioid/Mixture	F statistic, p value	Cohen's Effect Size (Cohen's F)	Current Power	Sample Size: Power≥0.8
Methadone	F(4, 40) = 0.98; p=0.428	0.313	0.282	35
Fentanyl	F(5, 50) = 1.94; p=0.104	0.441	0.607	20
Morphine	F(5, 50) = 0.36; p=0.875	0.189	0.133	75
Hydrocodone	F(4, 40) = 5.09; p=0.002	0.713	0.944	10
Buprenorphine	F(5, 50) = 0.75; p=0.589	0.274	0.248	40
Nalbuphine	F(5, 50) = 0.62; p=0.689	0.248	0.206	45
NAQ	F(4, 40) = 0.30; p=0.877	0.173	0.110	>100
Naltrexone	F(4, 40) = 0.18; p=0.947	0.135	0.085	>100
100:1	F(5, 50) = 2.02; p=0.093	0.449	0.626	20
56:1	F(5, 50) = 0.07; p=0.997	0.081	0.063	>100
32:1	F(5, 50) = 2.82; p=0.025	0.531	0.794	15
10:1	F(5, 50) = 1.43; p=0.231	0.378	0.460	25
3.2:1	F(5, 50) = 0.79; p=0.564	0.280	0.259	40

Supplementary Figure 1. Significant main effect of sex.

Figure 1a





Figure 1b



Supplementary Figure 1. Treatments for which there was a main effect of sex in statistical analysis. Abscissae: Dose fentanyl in mg/kg (VEH=vehicle). The naltrexone dose = fentanyl dose \div fentanyl proportion (10 in 1a, 3.2 in 1b). Ordinates: Locomotor counts per 60 minutes. Male and female data are shown in green and orange bars, respectively, and individual data are shown by points. Males had higher locomotor scores than females for these two treatments.

Supplementary Figure 2. Significant dose x sex interaction.

Figure 2a





Figure 2b



Supplemental Figure 2. Treatments for which there was a sex x dose interaction. Abscissae: Dose in mg/kg (VEH=vehicle). For panel 2a, the abscissa shows fentanyl dose, and the naltrexone dose = fentanyl dose ÷ fentanyl proportion (32). Ordinates: Locomotor counts per 60 minutes. Male and female data are shown in green and orange bars, respectively, and individual data are shown by points. Post hoc analysis did not indicate a significant difference between males and females at any dose of either treatment.