

Assessing the Value of the Zebrafish Conditioned Place Preference Model for Predicting Human Abuse Potential¹

A. J. Brock, S. M. G. Goody, A. N. Mead,¹ A. Sudwarts, M.O. Parker, and C.H. Brennan

School of Biological and Chemical Sciences, Queen Mary University of London, London, United Kingdom (A.J.B., A.S., C.H.B.); Global Safety Pharmacology, Drug Safety Research and Development, Pfizer Worldwide Research and Development, Groton, Connecticut (S.M.G.G., A.N.M.); and School of Health Sciences and Social Work, University of Portsmouth, Portsmouth, United Kingdom (M.O.P.)

Received May 16, 2017; accepted August 2, 2017

ABSTRACT

Regulatory agencies recommend that centrally active drugs are tested for abuse potential before approval. Standard pre-clinical assessments are conducted in rats or non-human primates (NHPs). This study evaluated the ability of the zebrafish conditioned place preference (CPP) model to predict human abuse outcomes. Twenty-seven compounds from a variety of pharmacological classes were tested in zebrafish CPP, categorized as positive or negative, and analyzed using standard diagnostic tests of binary classification to determine the likelihood that zebrafish correctly predict robust positive signals in human subjective effects studies (+HSE) and/or Drug Enforcement Administration drug scheduling. Results were then compared with those generated for rat self-administration and CPP, as well as NHP self-administration, using this same set of compounds. The findings reveal that zebrafish concordance

and sensitivity values were not significantly different from chance for both +HSE and scheduling. Although significant improvements in specificity and negative predictive values were observed for zebrafish relative to +HSE, specificity without sensitivity provides limited predictive value. Moreover, assessments in zebrafish provided no added value for predicting scheduling. By contrast, rat and NHP models generally possessed significantly improved concordance, sensitivity, and positive predictive values for both clinical measures. Although there may be predictive value with compounds from specific pharmacological classes (e.g., μ -opioid receptor agonists, psychostimulants) for zebrafish CPP, altogether these data highlight that using the current methodology, the zebrafish CPP model does not add value to the preclinical assessment of abuse potential.

Introduction

Zebrafish are one of the most widely used model systems in developmental biology. Owing to the numerous benefits of using this vertebrate system, including their small size, inexpensive housing costs, short life span, and unparalleled genetic tractability, they are also now becoming widely used in other areas of biological sciences, including pharmacological and behavioral screening (Guo, 2004; Stewart et al., 2015). The zebrafish system can provide a relatively high throughput in vivo option, bridging the gap between in vitro cell-based

models and in vivo rodent models; therefore, zebrafish may contribute to the refinement of animal use in research. Mounting evidence implicates zebrafish as a promising model species for reward and addiction research (Guo, 2004; Ninkovic et al., 2006; Mathur and Guo, 2010; Klee et al., 2012; Stewart et al., 2015). Thus, there may be a potential role for zebrafish abuse models in the drug discovery process, where centrally active compounds are typically assessed for their abuse potential in animal models before regulatory approval.


Two of the main animal models used to assess the rewarding properties of compounds are self-administration and conditioned place preference (CPP) (Panlilio and Goldberg, 2007; Tzschentke, 2007). Although self-administration has greater face and predictive validity to human drug-seeking behavior than CPP, the simplicity of CPP has led to its widespread use in the study of reward processes (Bardo and Bevins, 2000; Tzschentke, 2007). Based on the principles of Pavlovian conditioning, CPP is proposed to reflect the rewarding properties of drugs through examination of their association with contextual stimuli (Tzschentke, 1998, 2007). CPP protocols are typically conducted in an apparatus with a retractable partition that can divide the apparatus into two compartments, each side with distinct visual and/or tactile cues.

This work was supported by grants from the Medical Research Council UK (G1000403), National Committee for the Replacement Refinement and Reduction of Animals in Research (G1000053), BBSRC (BBM007863/1), EPSRC DTG (to A.S.), and a Royal Society Industry Fellowship to C.H.B.

S.M.G.G. is currently employed by Pfizer Drug Safety Research and Development, and A.N.M. was employed by Pfizer Drug Safety Research and Development during the inception of this work and the generation of data. A.N.M. is currently an employee of AstraZeneca. No information is presented that advocates for or promotes commercial products from any of our organizations. There are no other conflicts of interest.

¹Current affiliation: Drug Safety and Metabolism, AstraZeneca Innovative Medicines and Early Development, Cambridge, UK.

<https://doi.org/10.1124/jpet.117.242628>

 This article has supplemental material available at jpet.aspetjournals.org.

ABBREVIATIONS: CI, confidence interval; CNS, central nervous system; CPP, conditioned place preference; DMSO, dimethylsulfoxide; HSE, human subjective effects; NHP, nonhuman primate; PRU, proportionate reduction in uncertainty.

CPP usually comprises three phases: 1) the animal is allowed to explore the entire apparatus, and the time spent in each compartment is used as a measure of baseline preference; 2) the animal is sequentially restricted to each compartment for a period of time in which they are exposed to either drug or control, thus forming a Pavlovian association between the drug and the environment; and 3) the animal is once again allowed access to the entire apparatus in the absence of drug, and final place preference for the compartments is measured. If a significant change in preference toward the drug-paired compartment is observed, CPP is considered to have been established, reflecting rewarding properties of the drug.

Previous work has shown that cocaine (Darland and Dowling, 2001), amphetamine (Ninkovic et al., 2006), nicotine, ethanol (Kily et al., 2008; Brennan et al., 2011), and morphine (Lau et al., 2006) induce place preference in zebrafish; however, the number of compounds assessed in this model has been too limited to quantitatively analyze the predictive value of the model for effects in humans. The aim of the present study was to expand the number of compounds tested in zebrafish CPP using a standardized study design to determine the ability of the model to predict the outcome of robust positive human subjective effects (+HSE) and/or Drug Enforcement Administration (DEA) scheduling status in the United States. We selected scheduling status in addition to +HSE, because a number of factors other than the pharmacology of a drug can influence whether a drug is scheduled (e.g., medicinal uses, overall safety profile, as well as societal and economic factors). Compounds from a variety of pharmacological classes were tested, and the list included drugs that are considered to have positive outcomes in the clinic (including both strong and weak reinforcing drugs), as well as drugs that have negative clinical outcomes (i.e., do not exhibit +HSE or are unscheduled). The confidence of zebrafish CPP to predict positive clinical outcomes was determined using binary classification to categorize the selected drugs as true positives, true negatives, false positives, or false negatives in the model. Overall concordance and standard diagnostic tests of binary classifications were used to quantify objectively and statistically the predictive validity of zebrafish CPP. Similar analyses have been used to determine the predictive validity of other preclinical abuse assessments, including self-administration, drug discrimination, and locomotor activity in rats and non-human primates (NHPs) (O'Connor et al., 2011; Horton et al., 2013). As such, outcomes from zebrafish CPP were then compared with those reported in the literature with rat self-administration and CPP models, as well as NHP self-administration, using the same set of compounds.

Methods

Subjects. Wild-type zebrafish (*Danio rerio*) were either bred in house or obtained from a commercial supplier (Wades Tropical Import Ltd, London) at 4 weeks of age (Tubingen wild type) or Wades Singapore strain, respectively) and raised in the Queen Mary University London Fish Facility according to standard protocols. Based on power analysis of previous studies using nicotine and ethanol, we aimed to use approximately 20 fish for each concentration of each drug (including vehicle) in CPP experiments and 10 fish for each concentration of each drug (including vehicle) for locomotion analysis. All analyses were performed using 3- to 4-month-old (0.4–0.8 g) male and female fish with as close to 50% distribution of sexes across the concentration groups as possible. All behavioral experiments were carried out in a 28°C room

with water changes between each test subject to ensure water temperature was as consistent as possible for each fish and was similar to standard housing conditions. During experimental days, fish were fed in the morning (8:00–9:00 AM) as well as in the evening but not during an experiment. Fish were tested over an entire day (~8 hours, 10:00 AM–6:00 PM) with an equal number of fish from each concentration tested at the same time to minimize any effect that time of day may have on experimental outcomes. All experiments were carried out in accordance with the Animals (Scientific Procedures) Act, 1986, under local ethical guidelines from the Queen Mary Animal Care and Use Committee and under a UK Home Office project license.

CPP Experimental Design. CPP was conducted as previously published (Parker et al., 2013, 2014). Fish were singly housed for 1 week before being habituated to the conditioning tank over 2 consecutive days. The conditioning tank consisted of an opaque tank measuring 20 cm (w) × 15 cm (h) × 30 cm (l) containing 2.5 l of aquarium water with distinct visual cues (spots or stripes) on walls at each end of the tank (Supplemental Fig. 1). A ceiling-mounted camera and Noldus Ethovision XT 9 software (TrackSys, Nottingham, UK) were used to track automatically the behavior of the fish. After 2 days of habituation sessions, each drug was tested over 5 days, consisting of 1 baseline day and 3 conditioning days followed by 1 day of probe trials. During habituation, each individual fish was placed in the conditioning apparatus for 20 minutes with free access to both compartments and then returned to its home tank. On the day after the final habituation session, baseline assessments were carried out as follows: fish were placed in the conditioning tank for 10 minutes. Basal preference was determined by recording time spent in either compartment of the apparatus during the second 5 minutes of this period (% time spent in each side). Any fish with basal preference greater than 75% was excluded from the study. The number of fish excluded from the study by this criterion ranged widely, with 5% to 40% of the habituated fish showing a basal preference greater than 75%.

Each fish with basal preference less than 75% was then conditioned to the test drug (or its vehicle) over 3 consecutive days. Specifically, fish were placed in the conditioning tank containing 2.5 l of water and were restricted first to their preferred side for 20 minutes in the absence of drug (i.e., conditioned to vehicle) and then to their least preferred side for 20 minutes in the presence of test drug (or its vehicle in the case of the vehicle controls). The test drug (or vehicle) was added to the tank as a concentrated stock in a volume of 50-ml aquarium water. Each fish was conditioned to the same concentration of drug for 3 consecutive days. A total of five concentrations were used for each drug (including vehicle) with ~20 fish at each dose (dependent on how many fish were excluded from baseline). No data were recorded during the conditioning sessions. On the day after the final conditioning day, a 10-minute probe trial was conducted in which the fish were allowed free access to both sides of the tank in the absence of drug to measure the time spent on both sides. Place preference was calculated taking basal preference and preference for the drug-paired side into account, again using only the second 5-minute period for data collection. The last 5 minutes were used for analysis, because fish often display an initial “freezing” behavior as a stress response when first added to the test tank, particularly during baseline measurements. Omission of the first 5 minutes allows for a habituation period in which the fish are able to acclimate to the tank and ensures that this potential freezing behavior does not contribute toward calculations in change of preference scores. This was repeated in the probe trial to be consistent with the measurements being used for calculations in probe and baseline comparisons.

Locomotor Activity. All drugs were tested for their effects on locomotion (distance traveled) independent of the place preference assessment. This was included as a proxy measure to give an indication of whether the 20-minute exposure time used for CPP was sufficient to allow uptake of the test drug into the central nervous system (CNS). Each drug was tested at the same five concentrations as in the CPP assessment (using approximately 10 fish per concentration

TABLE 1
List of compounds tested by class along with supplier name, catalog number, concentration range tested and references used to select the concentration range

Compound	Supplier	Catalogue Number	Concentrations [μ M (mg/l)]	Vehicle	References
Atomoxetine HCl	Sequoia	SRP07328a	2–8.6 (0.6–2.5)	water	Wee and Woolverson (2004), Cantilena et al. (2012)
Bupropion HCl	Sequoia	SRP03446b	5.4–45.5 (1.5–12.5)	water	O'Connor et al. (2011), Vermoesen et al. (2011)
Caffeine	Johnson Matthey	A10431	26–260 (5–50)	water	Wong et al. (2010), O'Connor et al. (2011), Richendrfer et al. (2012)
Chlorpheniramine maleate	Sequoia	SRP02462c	1.3–10.2 (0.5–4)	water	Beardsley and Balster (1992), Hasenohrl et al. (2001)
Citalopram HBr	Sequoia	SRP03585c	1.2–14.8 (0.5–6)	water	Howell et al. (2007), Hiranita et al. (2009), Sackerman et al. (2010)
Cocaine HCl	Sigma	c5776	5.9–29.4 (2–10)	water	Darland and Dowling (2001), O'Connor et al. (2011)
D-Amphetamine sulfate	Sigma	A5880	5–50 (0.92–7.36)	water	Ninkovic and Bally-Cuif (2006), Irons et al. (2010), O'Connor et al. (2011)
Diazepam	Sigma	D0899	1.76–14 (0.5–4)	DMSO	Zhdanova et al. (2001), O'Connor et al. (2011), Richendrfer et al. (2012)
Diphenhydramine	Sequoia	SRP04365d	17–137 (5–40)	water	O'Connor et al. (2011)
Ethanol	VWR		100–300 (4.6–13.8)	water	Kily et al. (2008), O'Connor et al. (2011)
Fentanyl citrate	Sigma	F3886	0.008–0.303 (0.004–0.16)	water	O'Connor et al. (2011)
Fluoxetine HCl	Sequoia	SRP01950f	7.2–28.9 (0.6–2.5)	water	Lyness and Smith (1992), Howell et al. (2007), Maximino et al. (2011)
Ketamine HCl	Sigma	K2753	18.2–72.9 (5–20)	water	O'Connor et al. (2011), Riehl et al. (2011)
Methohexital	Sequoia	SRP02643m	3.8–38.1 (1–10)	DMSO	O'Connor et al. (2011)
Morphine sulfate	Sigma	M8777	0.7–7.9 (0.5–6)	water	Lau et al. (2006), Bretaud et al. (2007), O'Connor et al. (2011)
MPEP	Sequoia	SRP04265m	5.2–31 (1–6)	DMSO	Platt et al. (2008), van der Kam et al. (2009)
Naloxone HCl	Sequoia	SRP00860n	3.13–25 (1.25–10)	water	O'Connor et al. (2011)
Nicotine hemisulfate	Sigma	N1019	0.5–10 (0.2–4.2)	water	Bencan and Levin (2008), Kily et al. (2008), O'Connor et al. (2011)
Oxycodone	Tocris	3958	0.28–1.7 (0.1–0.6)	water	O'Connor et al. (2011)
Pentobarbital sodium	Sigma	P3761	20–120 (5–30)	water	Zhdanova et al. (2001), Renier et al. (2007), O'Connor et al. (2011)
Phencyclidine HCl	Sigma	P3029	0.36–3.6 (0.1–1)	water	O'Connor et al. (2011), Kyzar et al. (2012)
Procaine HCl	Sigma	P9879	92–733 (25–200)	water	Woolverson and Balster (1979), Johanson and Aigner (1981)
Retigabine HCl	Sequoia	SRP01080r	5.3–40 (2–15)	DMSO	Chege et al. (2012)
Rimobant	Sequoia	SRP01287r	0.68–13.5 (0.3–6)	DMSO	Braida et al. (2007), Justinova et al. (2011)
Tetracaine HCl	Sigma	T7508	0.83–6.3 (0.25–1.9)	water	Woolverson and Balster (1979), Wilcox et al. (2000), Wilcox et al. (2005)
9 Δ -THC	Sigma	T2386	0.16–0.64 (0.05–0.2)	ethanol	O'Connor et al. (2011)
WIN 55 212–2 mesylate	Sequoia	SRP00600w	0.06–1.15 (0.03–0.6)	DMSO	Wiley et al. (1995), Lecca et al. (2006)

THC, Δ -9 tetrahydrocannabinol; MPEP, 2-methyl-6(phenylethynyl)pyridine.

per drug). The order of drug exposure was pseudo-randomized between subjects. Fish were pre-exposed for 20 minutes in 1 l of aquarium water plus test drug in a tank measuring 11 cm (w) × 10 cm (h) × 20 cm (l). After drug exposure, the fish were netted into locomotor activity assay tanks [22 cm (w) × 16 cm (h) × 27 cm (l)] containing fresh aquarium water. Locomotor activity was recorded for 20 minutes using a ceiling mounted camera and Noldus Ethovision XT 9 software (TrackSys). These data were sorted into 2-minute time bins to allow temporal as well as spatial analysis.

Drugs and Concentrations. Twenty-seven compounds were analyzed in this study; 26 drugs were tested at four different concentrations plus vehicle, and one drug (ethanol) was tested at three different concentrations plus vehicle. The compounds that were tested included drugs that had been previously used in zebrafish CPP studies, including nicotine, ethanol, cocaine, morphine, caffeine, and amphetamine (Darland and Dowling, 2001; Lau et al., 2006; Ninkovic and Bally-Cuif, 2006; Ninkovic et al., 2006; Kily et al., 2008; Brennan et al., 2011; Collier et al., 2014), and 21 previously untested compounds (see Table 1 for drugs tested, supplier, concentration range, and vehicle).

Concentration ranges were based on concentrations previously found to be rewarding in zebrafish in other studies or using a concentration that corresponded to doses that were positive in either rodent or NHP CPP or self-administration studies. We employed a maximum final tank concentration (mg/l, administered as μM or mM solution in the tank water) equivalent to $2\times$ the mammalian effective dose (mg/kg). Compounds with no known rewarding properties were tested at similar ranges (mg/l, administered as a μM or mM solution in the tank water, equivalent to $2\times$ the mg/kg dose used in mammals). All drugs were made up as stock solutions in either water or DMSO (see Table 1) and stored frozen where applicable. All drugs were used at a pH of between 6.95 and 7.5 and were assessed for toxic effects before CPP analysis; starting at the lowest concentration, 3 fish were placed individually in a volume of 200 ml of the drug solution in fish water for an hour. Fish were assessed for signs of toxicity (difficulty swimming, exaggerated breathing, hemorrhaging gills, internal bleeding) during this time and at regular intervals for the following 6 hours and the following morning. If no adverse signs were detected, the concentration was increased and the procedure was repeated until the maximal intended dose had been assessed. Concentrations that induced signs of toxicity were not used further. Drugs were diluted from frozen stocks 10 minutes before use. All drugs were added to the conditioning tanks in a volume of 50 ml aquarium water. When used, the final concentration of DMSO did not exceed 0.1%.

CPP Data Analysis. Concentration-response graphs with 95% confidence interval (CI) estimates are included for all drugs assessed in the zebrafish CPP (see Supplemental Fig. 2). We assessed each of the compounds tested under the null hypothesis to ensure they did not induce a change in preference for the drug-paired stimulus after conditioning. A change in preference was calculated as the proportion of time spent in the drug-paired side in the probe trial minus the proportion of time spent in the drug-paired side during the basal preference trial. To assess whether a drug induced a statistically significant change in preference, a two-stage approach was implemented. First, an overall test for evidence of nonmonotonicity was performed. If there was no evidence of nonmonotonicity (i.e., the response was monotonic or, in other words, followed an increasing or decreasing trend with increasing concentration of drug), a sequential trend test was used. If there was evidence of nonmonotonicity (i.e., the response did not follow an increasing or decreasing trend with increasing concentration of drug), a Dunnett's post hoc test was performed.

In cases where there was no evidence of nonmonotonicity and a sequential trend test was conducted, a trend test across all concentrations was first performed. If this test was not statistically significant, the analysis was stopped and no concentrations were declared statistically significantly different from vehicle. If this test was statistically significant, the highest concentration was declared

statistically significant, and the process was repeated for all but the highest concentration. This process continued for the remaining lower concentrations, until the point at which the test was no longer statistically significant. In cases where there was evidence of non-monotonicity, a Dunnett's post hoc test was conducted to compare each concentration to vehicle.

Only drugs that induced a statistically significant increase in preference for the drug-paired side at any concentration were considered a positive in zebrafish CPP. If a drug induced a statistically significant decrease in preference for the drug-paired side (i.e., the drug might be considered aversive) or CPP was not observed at any concentration, the drug was deemed a negative in zebrafish CPP. CPP data were analyzed in RStudio (version 0.99.489) (RStudio, Boston, MA). For all analyses, statistical significance was set at $\alpha = 0.05$.

Locomotor Behavior Data Analysis. Data were fitted to linear mixed effects models, with distance traveled as the dependent variable, and time and concentration as fixed effects (fish identification as a random effect). Time bins in which the fish moved less than 100 cm in 2 minutes were checked for tracking failure, and where tracking failed, they were removed from analysis. For each drug, any time bins in which the fish moved two standard deviations from the mean distance moved across all drug concentrations in a drug group including paired controls were regarded as outliers and were excluded from analysis. Furthermore, any fish that "froze" (failed to travel 100 cm in any 2-minute time bin over the entire assay) were removed (20 of 1300 fish randomly distributed across all drugs and concentrations).

Binary Classification, Concordance, and Diagnostic Tests. The ability of zebrafish CPP to predict clinical outcomes with respect to +HSE, and scheduling status was assessed and compared with outcomes calculated with self-administration and CPP models in rats, as well as self-administration in NHPs. Classification of positives and negatives in zebrafish CPP are described above. For all other endpoints (+HSE, scheduling status, rat self-administration, rat CPP, and NHP self-administration), classification was conducted as described in Horton et al. (2013) and Supplemental Table 3. In brief, PubMed was the primary tool for locating peer-reviewed source documents. Google Scholar search engine was used as a follow up to obtain additional resources, but only peer-reviewed data or government documents were used for classifications. Many of the classifications were captured originally in Horton et al. (2013). A positive in rat/NHP self-administration or rat CPP was defined as a drug maintaining a higher level of responding under a fixed ratio schedule or inducing place preference for the drug-paired side, respectively, than the drug's vehicle. In cases where there were differences between data published in the literature, a drug was considered positive if any studies revealed a positive result. A drug was considered to induce +HSE if $>50\%$ of participants (volunteers with a history of drug use) reported positive "euphoric," "drug-liking," or "high" effects after drug administration using scales such as the Visual Analog Scale of global drug effects, the Addiction Research Center Inventory, the Drug Class Questionnaire, or the Profile of Mood States. If a drug has been classified with a scheduling status of I-V by the Controlled Substances Act, it was considered a positive for scheduling status. Drugs were considered negative in the clinical outcomes if they did not induce +HSE or were unscheduled.

For each preclinical model (zebrafish CPP, rat self-administration, rat CPP, NHP self-administration), drugs were classified relative to +HSE and scheduling status as either a true positive (positive in both the preclinical model and clinical measure), true negative (negative in both the preclinical model and clinical measure), false positive (positive in the preclinical model but negative in the clinical measure), or false negative (negative in the preclinical model but positive in the clinical measure). The overall concordance of each preclinical model was calculated by dividing the number of compounds for which the model correctly predicted the clinical outcome (total of true positives and true negatives) divided by the total number of compounds. Standard diagnostic tests of binary classifications (e.g., sensitivity

and specificity) and estimation of diagnostic value and level of uncertainty (e.g., positive and negative predictive value (positive predictive value and negative predictive value, respectively), value added positive predictive value and negative predictive value, and proportionate reduction in uncertainty [PRU] for positive and negative findings) were calculated as described in Coulthard (2007) and Horton et al. (2013) (Table 2). Pretest prevalence, or the overall probability of a drug being abused, was set as described previously in Horton et al. (2013). Briefly, the pretest prevalence was set at ~0.3, or 30%. This estimate was determined by dividing the number of scheduled drugs (II–V) in the United States in 2013 (total of 217) by the total number of approved drugs in the United States as described in the <http://www.fda.gov/cder/orange/default.htm>. U.S. Food and Drug Administration. Center for Drug Evaluation and Research Approved Drug Products with Therapeutic Equivalence Evaluations Orange Book (total of 1468), which gave a ratio of 0.15, or 15%. Given that many of the approved drugs are not centrally active and would therefore inherently lack abuse liability, this ratio was doubled, giving a final pretest prevalence of 0.3, or 30%, to estimate the likely prevalence in a data set with CNS-active compounds. For concordance, sensitivity, and specificity, pre-test probability was set at 50%, or chance. To determine the confidence or reliability of the statistical estimates used in this study, 95% CIs were calculated for each measure using R statistical software (version 3.3.0) and, in particular, the *PropCIs* and *pairwiseCI* libraries, as well as StatXact (Cytel Studio version 10; Cytel Inc. Cambridge, MA). In cases where CIs did not overlap with pretest probabilities (adjusted for prevalence where appropriate), it was considered that the model (e.g., zebrafish CPP, rat self-administration, rat CPP, or NHP self-administration) provided a statistically significant improvement in predictive value.

Results

Zebrafish CPP Analysis: Positives. As seen in previous studies, nicotine, ethanol, amphetamine, cocaine, and morphine induced statistically significant, concentration-dependent increases in preference for the drug-paired side (Fig. 1, A–E), with the largest magnitude changes occurring at concentrations of at 5 μ M ($25.6 \pm 7.8\%$), 171 mM ($21.5 \pm 3.7\%$), 50 μ M ($15.0 \pm 2.9\%$), 29.4 μ M ($26.9 \pm 4.3\%$), and 7.9 μ M ($19.8 \pm 7.8\%$), respectively (Table 3; Supplemental Table 1).

Of the previously untested compounds, the μ -opioid agonists fentanyl (Fig. 2A) and oxycodone (Fig. 2B) resulted in statistically significant, concentration-dependent increases in preference for the drug-paired side. The general anesthetics tetracaine (Fig. 2C) and phencyclidine (Fig. 2D), as well as the antihistamine chlorpheniramine (Fig. 2E), also significantly increased preference for the drug-paired side. The largest magnitude changes were observed for these drugs at the following concentrations: fentanyl, 0.076 μ M ($18.0 \pm 4.3\%$); oxycodone, 1.14 μ M ($25.0 \pm 5.0\%$); tetracaine, 6.3 μ M ($17.0 \pm 7.0\%$); phencyclidine, 3.57 μ M ($24.3 \pm 5.7\%$); chlorpheniramine, 5.1 μ M ($22.7 \pm 5.9\%$) (Tables 3 and 4; Fig. 2, A–E). All of these drugs were considered positives in the zebrafish model (Table 3; Supplemental Table 1).

Zebrafish CPP Analysis: Negatives. The remaining previously untested compounds did not induce a change in place preference in zebrafish at the concentrations evaluated in this study, including the general anesthetics ketamine (Fig. 3A) and procaine (Fig. 3B), the antidepressants atomoxetine (Fig. 3C), bupropion (Fig. 3D), citalopram (Fig. 3E), and fluoxetine (Fig. 3F), the antihistamine diphenhydramine (Fig. 3G), the benzodiazepine diazepam (Fig. 3H), the cannabinoid 1 receptor antagonist rimonabant (Fig. 3I), the cannabinoid

TABLE 2
Descriptions of and calculations for diagnostic tests conducted

Endpoint	Description	Calculation
Concordance	Proportion of compounds where the model accurately predicted clinical measure	$(TP + TN) / \text{Total \# of compounds}$
Sensitivity	Proportion of +HSE or scheduled drugs accurately identified by the model	$TP / (TP + FN)$
Specificity	Proportion of drugs that do not induce +HSE or are unscheduled accurately identified by the model	$TN / (TN + FP)$
Positive Predictive Value (PPV)	Proportion of drugs that are TPs relative to all drugs with a positive result	$TP / (TP + FP)$
Negative Predictive Value (NPV)	Proportion of drugs that are TNs relative to all drugs with a negative result	$TN / (TN + FN)$
Adjusted PPV	PPV adjusted for a pretest probability (prevalence) of 0.3	$(\text{sensitivity} \times \text{prevalence}) / [(\text{sensitivity} \times \text{prevalence}) + (1 - \text{specificity}) \times (1 - \text{prevalence})]$
Adjusted NPV	NPV adjusted for a pretest probability (prevalence) of 0.7	$\text{specificity} \times (1 - \text{prevalence}) / [\text{specificity} \times (1 - \text{prevalence}) + (1 - \text{sensitivity}) \times \text{prevalence}]$
Value added PPV (VaPPV)	Quantitative measurement of value added for positive results relative to pretest probability	Adjusted PPV – prevalence
Value added NPV (VaNPV)	Quantitative measurement of value added for negative results relative to pretest probability	Adjusted NPV – (1 – prevalence)
Proportionate reduction in uncertainty (PRU) positive (+)	Proportion by which model reduces uncertainty in predicting drugs with +HSE or are scheduled	$VaPPV / (1 - \text{prevalence})$
Proportionate reduction in uncertainty (PRU) negative (–)	Proportion by which model reduces uncertainty in predicting drugs without +HSE or are unscheduled	$VaNPV / \text{prevalence}$

FN, false negative; FP, false positive; TN, true negative; TP, true positive.




Fig. 1. Confirmation of change in preference (\pm S.E.M.) after conditioned place preference training in adult zebrafish in drugs previously assessed in this species are included to confirm reliability of the procedure. There were statistically significant concentration-dependent changes observed for nicotine (A) and ethanol (B), the stimulants amphetamine (C) and cocaine (D), and the opioid agonist morphine (E). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

receptor agonists Δ -9 tetrahydrocannabinol (Fig. 3J), and WIN-55,212 (Fig. 3K), the potassium channel opener retigabine (ezogabine) (Fig. 3L), the metabotropic glutamate receptor 5 (mGluR5) antagonist 2-methyl-6(phenylethynyl)pyridine (MPEP) (Fig. 3M), and the opioid receptor inverse agonist naloxone (Fig. 3N). The barbiturates methohexital and pentobarbital resulted in statistically significant decreases in preference for the drug-paired side, suggesting place aversion (Fig. 3, O and P). In agreement with previously published data (Collier et al., 2014), caffeine did not induce a statistically significant change in place preference (Fig. 3Q). All of these drugs were considered negatives in the zebrafish CPP model (Table 3; Supplemental Table 1).

Locomotor Activity Analysis. Locomotor activity was included as a proxy measure to give an indication of whether the 20-minute exposure time used for CPP was sufficient to allow uptake of the test drug into the CNS. All of the compounds tested in this study resulted in a significant concentration-dependent change in distance traveled over the 20-minute testing period, with the exception of procaine, phencyclidine, and fentanyl (Supplementary Fig. 3; Supplementary Table 2). Because phencyclidine and fentanyl resulted in a significant increase in place preference for the drug-paired side (Fig. 2; Table 3; Supplemental Table 1), they were assumed to be CNS active.

Predictive Value for Human Subjective Effects Outcome. Out of the 27 drugs evaluated in zebrafish CPP, HSE data were only available for 20 drugs, so only those 20 were used to determine the predictive value of zebrafish CPP for +HSE (Table 3; Supplemental Table 3). Data for rat CPP, rat self-administration, and NHP self-administration were only

available for 19, 18, and 19, respectively, of the 20 drugs with HSE data; therefore, only those drugs with data were used for calculating the predictive value of the models for +HSE using this data set. Table 4 and Supplemental Table 4 provide a statistical summary of the diagnostic tests of the models to +HSE. The overall concordance of zebrafish CPP relative to +HSE was 65%, compared with 84%, 83%, and 89% for rat CPP, rat self-administration, and NHP self-administration, respectively. Sensitivity values were generally high for +HSE with rat CPP and self-administration (0.93 for both models) and NHP self-administration (1.0), whereas sensitivity for zebrafish CPP was lower (0.53). For both concordance and sensitivity, only the rat and NHP models had values that were significantly greater than chance (or 50%). On the other hand, specificity for zebrafish CPP relative to +HSE was 1.0, whereas the values were lower for both rat and NHP models (ranging from 0.5 to 0.6). In this instance, only zebrafish had a specificity value that was significantly greater than chance. It should, however, be noted that the data set for specificity (i.e., clinical negatives) was relatively small in each of these cases ($n = 5$), which limits the ability to draw definitive conclusions. When estimated pretest prevalence (0.3) was taken into account, rat CPP and NHP self-administration possessed positive predictive values (0.5 and 0.52, respectively) that were significantly greater than the pretest prevalence. Although zebrafish CPP appeared to have the highest positive predictive value (1.0) compared with other models, there was overlap of 95% CIs with the pretest prevalence value of 0.3; therefore, this did not reach statistical significance. With regard to negative predictive value (i.e., the ability to correctly predict when a drug does not exhibit +HSE), zebrafish and rat CPP, as well as NHP self-administration, models offered

TABLE 3
Summary of binary classification for zebrafish and rat to human subjective effects and scheduling status.

Compound	Class	HSE	Scheduling	ZP CPP	Rat CPP	Self-Administration	Rat	References
Ketamine	Anesthetic	1	1	0	1		1	Horton et al. (2013), Guo et al. (2016)
Phencyclidine	Anesthetic	1	1	1	1		1	Marglin et al. (1989), Horton et al. (2013)
Procaine	Anesthetic	0	0	0	1		1	Spyraki et al. (1982), Fischman et al. (1983), Kiyatkin and Stein (1994)
Tetracaine	Anesthetic	X	0	1	X		X	dela Pena et al. (2011), Horton et al. (2013)
Atomoxetine	Antidepressant	0	0	0	0		0	Horton et al. (2013), Mori et al. (2013)
Bupropion	Antidepressant	0	0	0	1		1	Horton et al. (2013)
Citalopram	Antidepressant	X	0	0	X		0	Horton et al. (2013), Faillace et al. (2015)
Fluoxetine	Antidepressant	X	0	0	1		X	Horton et al. (2013), Horton et al. (2013)
Chlorpheniramine	Antihistamine	X	0	1	1		X	Suzuki et al. (1999), Horton et al. (2013)
Diphenhydramine	Antihistamine	1	0	0	1		1	Halpert et al. (2003), Horton et al. (2013)
Methohexital	Barbiturate	X	1	2	0		1	Pain et al. (1996), Horton et al. (2013)
Pentobarbital	Barbiturate	1	1	2	1		1	Bossert and Franklin (2003), Horton et al. (2013)
Diazepam	Benzodiazepine	1	1	0	0		1	Leri and Franklin (2000), Horton et al. (2013)
Rimonabant	Cannabinoid 1 receptor antagonist	0	X	0	0		X	Huestis et al. (2007), Li et al. (2008)
Δ^9 -THC	Cannabinoid receptor agonist	1	1	0	1		1	Braida et al. (2004), Horton et al. (2013)
WIN-55 212	Cannabinoid receptor agonist	X	X	0	2		1	Chaperon et al. (1998), Fattore et al. (2001)
Retigabine (Ezogabine)	K ⁺ channel opener	1	1	0	X		X	DEA (2011)
MPEP	mGluR5 receptor antagonist	X	X	0	1		1	van der Kam et al. (2009)
Fentanyl	μ -Opioid receptor agonist	1	1	1	1		1	Miller and Nation (1997), Vitale et al. (2003), Horton et al. (2013)
Morphine	μ Opioid receptor agonist	1	1	1	1		1	Zhang et al. (2012), Horton et al. (2013)
Oxycodone	μ -Opioid receptor agonist	1	1	1	1		1	Rutten et al. (2011), Horton et al. (2013)
Naloxone	Opioid receptor antagonist	0	0	0	0		0	Bardo and Neisewander (1986), Braida et al. (2005), Horton et al. (2013)
Amphetamine	Stimulant	1	1	1	1		1	Spyraki et al. (1982), Horton et al. (2013)
Caffeine	Stimulant	1	0	0	1		0	Bedingfield et al. (1998), Horton et al. (2013)
Cocaine	Stimulant	1	1	1	1		1	Bedingfield et al. (1998), Horton et al. (2013)
Ethanol	Stimulant	1	0	1	1		1	Morales et al. (2012), Horton et al. (2013)
Nicotine	Stimulant	1	0	1	1		1	Pascual et al. (2009), Horton et al. (2013)

1, positive finding; 2, aversive in CPP; 0, no effect; X, no data available




Fig. 2. Change in preference (\pm S.E.M.) after conditioned place preference training in adult zebrafish for opioid agonists fentanyl (A) and oxycodone (B), the general anesthetics tetracaine (C) and phencyclidine (D), and the antihistamine chlorpheniramine (E). * $P < 0.05$; ** $P < 0.01$.

negative predictive values that were significantly greater than pretest prevalence (0.83, 0.95, and 1.0, respectively). For this data set, although rat self-administration yielded a negative predictive value that was 0.94, which was within the same range as rat CPP and NHP self-administration; however, 95% CIs overlapped with pretest prevalence values for self-administration, therefore it did not reach statistical significance. Additionally, analysis of PRU values indicated that positive findings in rat CPP and NHP self-administration (PRU+) and negative findings in zebrafish and rat CPP, as well as NHP self-administration (PRU-), significantly reduced the uncertainty of predicting drugs correctly with regard to +HSE. These findings suggest higher confidence for NHP self-administration compared with other preclinical models that a positive result would correctly predict a drug with +HSE; however, classification of one drug, caffeine, as a positive in NHP self-administration appears to be what may be differentiating NHP self-administration from rat self-administration. Furthermore, there is lower confidence for zebrafish CPP compared with rat CPP and NHP self-administration that a negative result would result in a drug without +HSE.

Predictive Value for Scheduling Status. Twenty-four of the drugs evaluated in zebrafish CPP in this study have a scheduling designation by the Drug Enforcement Agency, so only these 24 were used to determine the predictive value (Table 4). Data for rat CPP, rat self-administration, and NHP self-administration were available for 24, 22, and 21, respectively, of the 24 drugs with scheduling status; therefore, only those drugs with scheduling data were used for calculating the predictive value of the rat and NHP models using this data set. Table 5 and Supplemental Table 4 provide a statistical summary of the diagnostic tests of the models to scheduling

status. With regard to scheduling, overall concordance was relatively similar for zebrafish and rat CPP (58% and 52%, respectively) and NHP self-administration (62%), whereas rat self-administration was higher (75%) and was the only model to have a value that was significantly greater than chance (50%). Similar to the data generated for +HSE, sensitivity values were generally high for scheduling status with rat CPP, rat self-administration, and NHP self-administration (0.82, 1.0, and 1.0, respectively), whereas sensitivity for zebrafish CPP was lower (0.5). With respect to scheduling status, specificity values were lower than the values generated for +HSE for all four models, ranging from 0.2 (rat CPP and NHP self-administration) to 0.44 (rat self-administration) to 0.67 (zebrafish CPP). Taking prevalence into account, rat self-administration provided the highest predictive values for both positive (0.44) and negative (1.0) findings, demonstrating increased absolute added value over pretest prevalence (0.3). This contrasts with results obtained for the other models, where adjusted predictive values were at or near pretest prevalence values. As a result, these findings reveal no or limited absolute added value for either positive or negative findings with regard to scheduling status for zebrafish and rat CPP, as well as NHP self-administration, for this data set. Similar to rat self-administration, NHP self-administration yielded a high adjusted negative predictive value (1.0); however 95% CIs overlapped with pretest prevalence. Examination of PRU+ and PRU- values shows that with respect to scheduling status, positive and negative results in the rat self-administration model significantly reduce the uncertainty of predicting scheduling by the highest proportions, at 0.19 and 1.0 for PRU+ and PRU-, respectively. Zebrafish CPP models offered lower reductions in uncertainty for positive and negative findings (proportions of 0.13

TABLE 4

Statistical summary of diagnostic tests for the prediction of human subjective effects

Summary for prediction of human subjective effects, demonstrating the number of false positives, true positives, true negatives, false negatives, and outcome ratios for various statistical outputs with regard to diagnostic value. Values in parentheses indicate 95% confidence intervals. Values indicate a significant difference from pretest probability [concordance, PPV, NPV, PPV (30% prev), NHP (30% prev)] or from zero predictive value (VaPPV, VaNPV). PPV and NPV represent observed predictive values. PPV (30%) and NPV (30% prev) represent predictive values corrected for an estimated prevalence of 0.3.

Model	False Positives	True Positives	True Negatives	False Negatives	Concordance	Sensitivity	Specificity	PPV	Adjusted PPV (30% prev)	Value Added PPV	NPV	Adjusted NPV (30% prev)	Value Added NPV	PRU Positive	PRU Negative
Zebrafish CPP	0	8	5	7	0.65 (0.43, 0.82)	0.53 (0.30, 0.75)	1.0 (0.57, 1.0)	1.0 (0.68, 1.0)	1.0 (0.29, 1.0)	0.7 (-0.01, 0.70)	0.42 (0.19, 0.68)	0.83 (0.70, 0.92)	0.13 (0, 0.22)	1.0 (-0.01, 1.0)	0.44 (0, 0.72)
Rat CPP	2	13	3	1	0.84 (0.62, 0.94)	0.83 (0.69, 0.99)	0.6 (0.23, 0.88)	0.87 (0.62, 0.96)	0.5 (0.33, 0.87)	0.2 (0, 0.57)	0.75 (0.3, 0.95)	0.95 (0.7, 1.0)	0.25 (0, 0.43)	0.28 (0, 0.82)	0.84 (0.01, 0.99)
Rat self-administration	2	13	2	1	0.83 (0.61, 0.94)	0.83 (0.69, 0.99)	0.5 (0.15, 0.85)	0.87 (0.62, 0.96)	0.44 (0.29, 0.84)	0.14 (-0.01, 0.54)	0.67 (0.21, 0.94)	0.94 (0.62, 1.0)	0.24 (-0.05, 0.3)	0.20 (-0.02, 0.77)	0.81 (-0.23, 0.99)

NPV, negative predictive value, PRU positive, proportionate reduction in uncertainty for positive findings; PPV, positive predictive value; PRU negative, proportionate reduction in uncertainty for negative findings.

and 0.19, respectively) that were not statistically significant, and rat CPP models yielded no reductions in uncertainty (proportions of 0.01 and 0.07 for PRU+ and PRU-, respectively). NHP self-administration did not yield any statistically significant reductions in uncertainty for positive (0.07) or negative (1.0) findings.

Discussion

In this work, 27 compounds were evaluated in zebrafish CPP to determine how the model predicts human abuse outcomes. At the concentrations tested, 10 induced CPP and were considered positives, and 17 did not induce CPP and deemed negatives. With the exception of procaine, each of the negatives affected locomotion, suggesting brain penetration. As procaine had no effect on locomotion or CPP, one possibility is that its brain penetration may have been impaired. Results from zebrafish CPP were then used to objectively and quantitatively analyze the model's predictive value for +HSE or scheduling. These results were compared with outcomes generated with rat self-administration and CPP, as well as NHP self-administration. The findings show that zebrafish CPP specificity, but not sensitivity, is significantly greater than pretest probability with respect to HSE; therefore, there is potential value in predicting negative clinical outcomes (drugs without +HSE). However, there is limited value for zebrafish to correctly predict drugs with +HSE. Because zebrafish values were not greater than pretest prevalence for scheduling status, the data indicate no added value for predicting scheduling. With few exceptions, rat and NHP models generally possessed concordance, sensitivity, and predictive values that were significantly greater than pretest probabilities for both clinical measures. Overall, although there may be predictive value for zebrafish for some pharmacological classes, such as μ -opioid receptor agonists and psychostimulants, these data highlight that rat and NHP models possess greater predictive value than zebrafish for this limited set of compounds.

This is the first study to demonstrate that fentanyl, oxycodone, tetracaine, phencyclidine, and chlorpheniramine are rewarding in zebrafish CPP. Given that all μ -opioid receptor agonists induced CPP in zebrafish, the data suggests conserved opioid reward pathways with mammals. In contrast, only 2 of the 4 anesthetics tested induced CPP, with no consistency in the mechanism of action and the observed CPP result. Although both ketamine and phencyclidine act as *N*-methyl-D-aspartate receptor antagonists, phencyclidine, but not ketamine, induced CPP. Tetracaine and procaine are local anesthetics that inhibit monoamine uptake transporters, but only tetracaine induced CPP. Similarly, both diphenhydramine and chlorpheniramine are histamine receptor H1 antagonists; however, CPP was only produced with chlorpheniramine. The basis for these differences has yet to be determined; however, it is possible that the concentrations tested, species-specific drug sensitivities or target expression play a role.

Data from zebrafish CPP were analyzed using standard diagnostic tests of binary classification to determine the model's predictive value relative to +HSE and scheduling. Other abuse-related preclinical models, such as in vitro binding and functional activity, locomotor activity, drug discrimination, and self-administration have been evaluated similarly




Fig. 3. Change in preference (\pm S.E.M.) after conditioned place preference training in adult zebrafish for the general anesthetics ketamine (A) and procaine (B); antidepressants atomoxetine (C), bupropion (D), citalopram (E), and fluoxetine (F); antihistamine diphenhydramine (G); benzodiazepine diazepam (H); CB1 receptor antagonist rimonabant (I); CB receptor agonists Δ -9 tetrahydrocannabinol (THC) (J) and WIN-55,212 (K); potassium channel opener retigabine (L); mGluR5 antagonist 2-methyl-6(phenylethynl)pyridine (MPEP; M); opioid receptor inverse agonist naloxone (N); barbiturates methohexital (O) and pentobarbital (P); caffeine (Q). * $P < 0.05$.

(Horton et al., 2013). To compare the ability of zebrafish CPP to predict clinical abuse outcomes with other preclinical models, binary classifications of rat and NHP self-administration and rat

CPP were generated and analyzed using published literature sources. Because not all drugs assessed in zebrafish have reported HSE data or are approved for medicinal use (and

TABLE 5

Statistical summary of diagnostic tests for the prediction of scheduling status

Summary for prediction of scheduling status, demonstrating the number of false positives, true positives, false negatives, and outcome ratios for various statistical outputs with regards to diagnostic value. Values in parentheses indicate 95% confidence intervals. Values indicate a significant difference from pretest probability [concordance, PPV, NPV, PPV (30% prev), NHP (30% prev), NPV (30% prev), PPV and NPV represent observed predictive values. PPV (30%) and NPV (30% prev) represent predictive values corrected for an estimated prevalence of 0.3.

Model	False positives	True positives	True negatives	False negatives	Concordance	Sensitivity	Specificity	PPV	Adjusted PPV (30% prev)	Value added PPV	NPV	Adjusted NPV (30% prev)	Value added NPV	PRU positive	PRU negative
Zebrafish CPP	4	6	8	6	0.58 (0.33, 0.76)	0.5 (0.25, 0.75)	0.67 (0.39, 0.86)	0.6 (0.31, 0.83)	0.39 (0.19, 0.7)	0.09 (-0.11, 0.4)	0.57 (0.33, 0.79)	0.76 (0.59, 0.88)	0.06 (-0.11, 0.18)	0.13 (-0.16, 0.57)	0.19 (-0.37, 0.59)
Rat CPP	8	9	2	2	0.52 (0.32, 0.72)	0.82 (0.52, 0.95)	0.20 (0.06, 0.51)	0.53 (0.31, 0.74)	0.30 (0.20, 0.44)	0 (-0.10, 0.14)	0.50 (0.15, 0.85)	0.72 (0.15, 0.97)	0.02 (-0.55, 0.27)	0.01 (-0.14, 0.20)	0.07 (-1.83, 0.91)
Rat self-administration	5	11	4	0	0.75 (0.53, 0.89)	1.0 (0.74, 1.0)	0.44 (0.19, 0.73)	0.69 (0.44, 0.86)	0.44 (0.32, 0.67)	0.14 (0.02, 0.37)	1.0 (0.51, 1.0)	1.0 (0.77, 1.0)	0.3 (0.07, 0.3)	0.19 (0.03, 0.53)	1.0 (0.22, 1.0)

NPV, negative predictive value; PPV, positive predictive value; PRU positive, proportionate reduction in uncertainty for positive findings; PRU negative, proportionate reduction in uncertainty for negative findings.

consequently do not have scheduling data), the sample sizes for our analyses were generally small (~18–24 drugs, depending on the model). Therefore, one limitation is that the estimates of variance are relatively high for some comparisons, and data should be interpreted with this in mind.

For both +HSE and scheduling, six drugs were considered true positives in zebrafish: phencyclidine, fentanyl, morphine, oxycodone, amphetamine, and cocaine. Four drugs (ethanol, nicotine, tetracaine, and chlorpheniramine) were considered true positives with regard to +HSE, but were deemed false positives with regard to scheduling (Tables 6 and 7). Classification of ethanol and nicotine as false positives with respect to scheduling is because both are unscheduled drugs due to their historically accepted societal use. Neither tetracaine nor chlorpheniramine are scheduled, which may reflect their medicinal use (e.g., tetracaine is typically used as ophthalmic drops) or lack of real-world human abuse. True positives included psychostimulants, μ -opioid receptor agonists, general anesthetics, and antihistamines. CPP was not induced by drugs from the barbiturate or benzodiazepine classes at the concentrations used here (Tables 6 and 7). Given that varying results have been reported with the barbiturates methohexital and pentobarbital in other abuse-related pre-clinical models depending on the route of administration and doses evaluated (Pickens et al., 1981; Mucha and Iversen, 1984; Lew and Parker, 1998; O'Connor et al., 2011), a negative result in a zebrafish model may reflect an inappropriate concentration range or may be due to differences in the procedures used (CPP versus self-administration).

The lack of false positives with regard to +HSE revealed that zebrafish CPP yielded the highest adjusted positive predictive value compared with rats and NHPs; however, because CIs for zebrafish overlapped with pretest prevalence, the data did not reach statistical significance. Inclusion of more compounds for this analysis could potentially alter the results to suggest significant added value. Although zebrafish CPP possessed significant value added for correctly predicting when a drug would not exhibit +HSE (i.e., adjusted negative predictive value), rat and NHP models were generally higher. Importantly, without confidence in the ability to predict correctly +HSE (i.e., positive clinical outcome), there is minimal value added overall for zebrafish CPP. Additionally, given that none of the diagnostic tests for correctly predicting scheduling with zebrafish CPP were significantly greater than chance or pretest prevalence, it can be concluded that zebrafish do not provide added value for predicting scheduling.

A few of the drugs that were false negatives with regard to +HSE and scheduling had limited solubility, requiring DMSO or ethanol as a solvent. Thus poor solubility may have prevented efficient uptake into the fish. However, methohexital induced significant conditioned place aversion, and all of the false negatives induced effects on locomotion. Diazepam also reportedly induces anxiolytic effects in zebrafish using similar concentrations (Bencan et al., 2009; Maximino et al., 2010, 011). Thus failure of these drugs to induce CPP does not appear to be due to inefficient uptake. DMSO could also influence visual acuity (Hull et al., 1969; Silverman, 1972), anxiety status (Hallare et al., 2004, 2005), or locomotion (Chen et al., 2011). Indeed, it is notable that “control” group data with drugs formulated in DMSO were more variable than when water was used.

TABLE 6

Summary of false positives and false negatives for each model with respect to positive human subjective effects

Zebrafish CPP		Rat CPP		Rat Self-Administration		NHP Self-Administration	
False positive	False negative	False positive	False negative	False positive	False negative	False positive	False negative
	Ketamine Diphenhydramine Pentobarbital Diazepam Δ 9-THC Retigabine Caffeine	Procaine Bupropion	Diazepam	Procaine Bupropion	Caffeine	Procaine Bupropion	

The finding that rat self-administration offers limited added value over pretest prevalence for HSE is consistent with previous reports using an even larger data set (56 drugs) (Horton et al., 2013). Interestingly, NHP self-administration does offer significantly greater predictive value over pretest for +HSE; however, this appears to be driven by one drug, caffeine, which was deemed as a positive here due to the criteria used, but variable results are reported in the literature. Subjective effects are generally assessed preclinically using drug discrimination procedures and are not always linked to the reinforcing effects of a drug (Ator, 2002; Martelle and Nader, 2009). Thus, it is no surprise that significant predictive value is added for rat drug discrimination compared with +HSE (Horton et al., 2013). Furthermore, given that both rat and zebrafish CPP yielded limited value with regard to scheduling, the data suggest a limitation of the model itself and not the species. CPP and self-administration measure fundamentally different behaviors. It is possible that these differences are important in terms of correctly predicting scheduling or +HSE.

Significant drawbacks exist for using zebrafish CPP as performed here in the pharmaceutical industry. First, the attrition rate can be high, given that 5–40% of zebrafish fail basal preference requirements and some fish tend to “freeze.” Second, immersing zebrafish in the tank water containing drug during conditioning can be cost prohibitive. The average drug requirement was 7 g (ranging from 0.04–68 g) for the concentrations tested here. Alternatives to immersing zebrafish in the tank water may exist, such as pretreating zebrafish in a smaller volume containing drug before immersing them in the conditioning tank or direct injections into zebrafish (Ninkovic et al., 2006; Cadet, 2009). For the latter, both ethical and practical limitations exist, including repeat injections that can cause local trauma at the injection site. Third, water-insoluble compounds that require solvents may

prove problematic. Administering compounds to zebrafish in food pellets (Zang et al., 2011) is an option that may be more cost effective and avoid solubility issues. Automated systems to deliver such pellets also exist (Brock et al., 2017), which may allow for the development of self-administration paradigms.

The intention of this work was not to propose replacing mammalian abuse potential assessments in drug development before approval. Rather, the goal was to determine whether there is potential value of zebrafish CPP earlier in drug discovery, when screening compounds might allow for redirection of resources away from targets or chemical series with high risk. Also, the compounds selected for this assessment were not all inclusive; therefore, it is possible that additional data may alter the interpretation of zebrafish CPP predictive value. Even so, by evaluating drugs from a variety of pharmacological classes that are both positive and negative clinically for abuse, this work expands our understanding of the translation of zebrafish CPP and offers some insight about how it might be applied in drug discovery, particularly with drugs from certain classes.

Acknowledgments

The authors thank Dr. Helen Moore for involvement with experimental design and conduct and Dr. David Potter and Dr. Dingzhou Li from the Pfizer Drug Safety Research and Development Statistics for advice on the statistical analyses conducted in the study.

Authorship Contributions

Participated in research design: Goody, Mead, Parker, Brennan.
Conducted experiments: Brock, Sudwarts.
Performed data analysis: Goody, Parker, Sudwarts.
Wrote or contributed to the writing of the manuscript: Brock, Goody, Mead, Parker, Brennan.

TABLE 7

Summary of false positives and false negatives for each model with respect to scheduling status

Zebrafish CPP		Rat CPP		Rat Self-Administration		NHP Self-Administration	
False positive	False negative	False positive	False negative	False positive	False negative	False Positive	False negative
Tetracaine Chlorpheniramine Ethanol Nicotine	Ketamine Methohexital Pentobarbital Diazepam Δ 9-THC Retigabine	Procaine Bupropion Fluoxetine Chlorpheniramine Diphenhydramine Caffeine Ethanol Nicotine	Methohexital Diazepam	Procaine Bupropion Diphenhydramine Ethanol Nicotine		Procaine Tetracaine Bupropion Chlorpheniramine Diphenhydramine Caffeine Ethanol Nicotine	

References

- Ator NA (2002) Relation between discriminative and reinforcing effects of midazolam, pentobarbital, chlordiazepoxide, zolpidem, and imidazenil in baboons. *Psychopharmacology (Berl)* **163**:477–487.
- Bardo MT and Bevins RA (2000) Conditioned place preference: what does it add to our preclinical understanding of drug reward? *Psychopharmacology (Berl)* **153**:31–43.
- Bardo MT and Neisewander JL (1986) Single-trial conditioned place preference using intravenous morphine. *Pharmacol Biochem Behav* **25**:1101–1105.
- Beardsley PM and Balster RL (1992) The intravenous self-administration of anti-histamines by rhesus monkeys. *Drug Alcohol Depend* **30**:117–126.
- Bedingfield JB, King DA, and Holloway FA (1998) Cocaine and caffeine: conditioned place preference, locomotor activity, and additivity. *Pharmacol Biochem Behav* **61**:291–296.
- Bencan Z and Levin ED (2008) The role of alpha7 and alpha4beta2 nicotinic receptors in the nicotine-induced anxiolytic effect in zebrafish. *Physiol Behav* **95**:408–412.
- Bencan Z, Sledge D, and Levin ED (2009) Buspirone, chlordiazepoxide and diazepam effects in a zebrafish model of anxiety. *Pharmacol Biochem Behav* **94**:75–80.
- Bossert JM and Franklin KB (2003) Reinforcing versus anticonvulsant drugs: effects on intracranial self-stimulation rate-frequency M50 indices. *Behav Brain Res* **144**:243–247.
- Braida D, Iosùè S, Pegorini S, and Sala M (2004) Delta9-tetrahydrocannabinol-induced conditioned place preference and intracerebroventricular self-administration in rats. *Eur J Pharmacol* **506**:63–69.
- Braida D, Iosùè S, Pegorini S, and Sala M (2005) 3,4-Methylenedioxymethamphetamine-induced conditioned place preference (CPP) is mediated by endocannabinoid system. *Pharmacol Res* **51**:177–182.
- Braida D, Limonta V, Pegorini S, Zani A, Guerini-Rocco C, Gori E, and Sala M (2007) Hallucinatory and rewarding effect of salvinorin A in zebrafish: kappa-opioid and CB1-cannabinoid receptor involvement. *Psychopharmacology (Berl)* **190**:441–448.
- Brennan CH, Parmar A, Kily LK, Ananthathavan A, Doshi A, and Patel S (2011) Conditioned place preference models of drug dependence and relapse to drug seeking: studies with nicotine and ethanol, in *Zebrafish Models in Neurobehavioral Research*, pp 163–179. Humana Press, Totowa, NJ.
- Bretau S, Li Q, Lockwood BL, Kobayashi K, Lin E, and Guo S (2007) A choice behavior for morphine reveals experience-dependent drug preference and underlying neural substrates in developing larval zebrafish. *Neuroscience* **146**:1109–1116.
- Brock AJ, Sudwartz A, Daggett J, Parker MO, and Brennan CH (2017) A fully automated computer based Skinner box for testing learning and memory in zebrafish. *bioRxiv*. doi doi.org/10.1101/110478:
- Cadet JL (2009) Amphetamine recapitulates developmental programs in the zebrafish. *Genome Biol* **10**:231.
- Cantilena L, Kahn R, Duncan CC, Li S-H, Anderson A, and Elkashaf A (2012) Safety of atomoxetine in combination with intravenous cocaine in cocaine-experienced participants. *J Addict Med* **6**:265–273.
- Chaperon F, Soubrié P, Puech AJ, and Thiébot MH (1998) Involvement of central cannabinoid (CB1) receptors in the establishment of place conditioning in rats. *Psychopharmacology (Berl)* **135**:324–332.
- Chege SW, Hortopan GA, T Dinday M, and Baraban SC (2012) Expression and function of KCNQ channels in larval zebrafish. *Dev Neurobiol* **72**:186–198.
- Chen TH, Wang YH, and Wu YH (2011) Developmental exposures to ethanol or dimethylsulfoxide at low concentrations alter locomotor activity in larval zebrafish: implications for behavioral toxicity bioassays. *Aquat Toxicol* **102**:162–166.
- Collier AD, Khan KM, Caramillo EM, Mohn RS, and Echevarria DJ (2014) Zebrafish and conditioned place preference: a translational model of drug reward. *Prog Neuropsychopharmacol Biol Psychiatry* **55**:16–25.
- Coulthard MG (2007) Quantifying how tests reduce diagnostic uncertainty. *Arch Dis Child* **92**:404–408.
- Darland T and Dowling JE (2001) Behavioral screening for cocaine sensitivity in mutant zebrafish. *Proc Natl Acad Sci USA* **98**:11691–11696.
- DEA (2011) Schedules of controlled substances: placement of ezogabine into schedule V. *Fed Regist* **76**:77895–77899.
- dela Peña IC, Ahn HS, Ryu JH, Shin CY, Park IH, and Cheong JH (2011) Conditioned place preference studies with atomoxetine in an animal model of ADHD: effects of previous atomoxetine treatment. *Eur J Pharmacol* **667**:238–241.
- Faillace MP, Zwiller J, and Bernabeu RO (2015) Effects of combined nicotine and fluoxetine treatment on adult hippocampal neurogenesis and conditioned place preference. *Neuroscience* **300**:104–115.
- Fattore L, Cossu G, Martellotta CM, and Fratta W (2001) Intravenous self-administration of the cannabinoid CB1 receptor agonist WIN 55,212-2 in rats. *Psychopharmacology (Berl)* **156**:410–416.
- Fischman MW, Schuster CR, and Rajfer S (1983) A comparison of the subjective and cardiovascular effects of cocaine and procaine in humans. *Pharmacol Biochem Behav* **18**:711–716.
- Guo R, Tang Q, Ye Y, Lu X, Chen F, Dai X, Yan Y, and Liao L (2016) Effects of gender on ketamine-induced conditioned place preference and urine metabolomics. *Regul Toxicol Pharmacol* **77**:263–274.
- Guo S (2004) Linking genes to brain, behavior and neurological diseases: what can we learn from zebrafish? *Genes Brain Behav* **3**:63–74.
- Hallare AV, Köhler HR, and Triebkorn R (2004) Developmental toxicity and stress protein responses in zebrafish embryos after exposure to diclofenac and its solvent, DMSO. *Chemosphere* **56**:659–666.
- Hallare AV, Pagulayan R, Lacdan N, Köhler HR, and Triebkorn R (2005) Assessing water quality in a tropical lake using biomarkers in zebrafish embryos: developmental toxicity and stress protein responses. *Environ Monit Assess* **104**:171–187.
- Halpert AG, Olmstead MC, and Beninger RJ (2003) Dimethylhydrazine produces a conditioned place preference in rats. *Pharmacol Biochem Behav* **75**:173–179.
- Hasenöhrl RU, Kuhlen A, Frisch C, Galosi R, Brändao ML, and Huston JP (2001) Comparison of intra-accumbens injection of histamine with histamine H1-receptor antagonist chlorpheniramine in effects on reinforcement and memory parameters. *Behav Brain Res* **124**:203–211.
- Hiranita T, Soto PL, Newman AH, and Katz JL (2009) Assessment of reinforcing effects of benzotropine analogs and their effects on cocaine self-administration in rats: comparisons with monoamine uptake inhibitors. *J Pharmacol Exp Ther* **329**:677–686.
- Horton DB, Potter DM, and Mead AN (2013) A translational pharmacology approach to understanding the predictive value of abuse potential assessments. *Behav Pharmacol* **24**:410–436.
- Howell LL, Carroll FI, Votaw JR, Goodman MM, and Kimmel HL (2007) Effects of combined dopamine and serotonin transporter inhibitors on cocaine self-administration in rhesus monkeys. *J Pharmacol Exp Ther* **320**:757–765.
- Huestis MA, Boyd SJ, Heishman SJ, Preston KL, Bonnet D, Le Fur G, and Gorelick DA (2007) Single and multiple doses of rimonabant antagonize acute effects of smoked cannabis in male cannabis users. *Psychopharmacology (Berl)* **194**:505–515.
- Hull FW, Wood DC, and Brobyn RD (1969) Eye effects of DMSO. Report of negative results. *Northwest Med* **68**:39–41.
- Irons TD, MacPhail RC, Hunter DL, and Padilla S (2010) Acute neuroactive drug exposures alter locomotor activity in larval zebrafish. *Neurotoxicol Teratol* **32**:84–90.
- Johanson CE and Aigner T (1981) Comparison of the reinforcing properties of cocaine and procaine in rhesus monkeys. *Pharmacol Biochem Behav* **15**:49–53.
- Justinová Z, Yasar S, Redhi GH, and Goldberg SR (2011) The endogenous cannabinoid 2-arachidonoylglycerol is intravenously self-administered by squirrel monkeys. *J Neurosci* **31**:7043–7048.
- Kily LJ, Cowe YC, Hussain O, Patel S, McElwaine S, Cotter FE, and Brennan CH (2008) Gene expression changes in a zebrafish model of drug dependency suggest conservation of neuro-adaptation pathways. *J Exp Biol* **211**:1623–1634.
- Kiyatkin EA and Stein EA (1994) Biphasic changes in mesolimbic dopamine signal during cocaine self-administration. *Neuroreport* **5**:1005–1008.
- Klee EW, Schneider H, Clark KJ, Cousin MA, Ebbert JO, Hooten WM, Karpyak VM, Warner DO, and Ekker SC (2012) Zebrafish: a model for the study of addiction genetics. *Hum Genet* **131**:977–1008.
- Kyzar EJ, Collins C, Gaikwad S, Green J, Roth A, Monnig L, El-Ounsi M, Davis A, Freeman A, Capezio N, et al. (2012) Effects of hallucinogenic agents mescaline and phencyclidine on zebrafish behavior and physiology. *Prog Neuropsychopharmacol Biol Psychiatry* **37**:194–202.
- Lau B, Bretau S, Huang Y, Lin E, and Guo S (2006) Dissociation of food and opiate preference by a genetic mutation in zebrafish. *Genes Brain Behav* **5**:497–505.
- Lecca D, Cacciapaglia F, Valentini V, and Di Chiara G (2006) Monitoring extracellular dopamine in the rat nucleus accumbens shell and core during acquisition and maintenance of intravenous WIN 55,212-2 self-administration. *Psychopharmacology (Berl)* **188**:63–74.
- Leri F and Franklin KB (2000) Diazepam in the ventral striatum dissociates dopamine-dependent and dopamine-independent place conditioning. *Neuroreport* **11**:2553–2557.
- Lew G and Parker LA (1998) Pentobarbital-induced place aversion learning. *Anim Learn Behav* **26**:219–224.
- Li F, Fang Q, Liu Y, Zhao M, Li D, Wang J, and Lu L (2008) Cannabinoid CB1(1) receptor antagonist rimonabant attenuates reinstatement of ketamine conditioned place preference in rats. *Eur J Pharmacol* **589**:122–126.
- Lyness WH and Smith FL (1992) Influence of dopaminergic and serotonergic neurons on intravenous ethanol self-administration in the rat. *Pharmacol Biochem Behav* **42**:187–192.
- Marglin SH, Milano WC, Mattie ME, and Reid LD (1989) PCP and conditioned place preferences. *Pharmacol Biochem Behav* **33**:281–283.
- Martelle JL and Nader MA (2009) A within-subject assessment of the discriminative stimulus and reinforcing effects of self-administered cocaine in rhesus monkeys. *Psychopharmacology (Berl)* **203**:343–353.
- Mathur P and Guo S (2010) Use of zebrafish as a model to understand mechanisms of addiction and complex neurobehavioral phenotypes. *Neurobiol Dis* **40**:66–72.
- Maximino C, de Brito TM, da Silva Batista AW, Herculano AM, Morato S, and Gouveia A, Jr (2010) Measuring anxiety in zebrafish: a critical review. *Behav Brain Res* **214**:157–171.
- Maximino C, da Silva AWB, Gouveia A, Jr, and Herculano AM (2011) Pharmacological analysis of zebrafish (*Danio rerio*) scototaxis. *Prog Neuropsychopharmacol Biol Psychiatry* **35**:624–631.
- Miller DK and Nation JR (1997) Chronic cadmium exposure attenuates the conditioned reinforcing properties of morphine and fentanyl. *Brain Res* **776**:162–169.
- Morales M, Varlinskaya EI, and Spear LP (2012) Evidence for conditioned place preference to a moderate dose of ethanol in adult male Sprague-Dawley rats. *Alcohol* **46**:643–648.
- Mori T, Shibasaki M, Ogawa Y, Hokazono M, Wang TC, Rahmadi M, and Suzuki T (2013) Comparison of the behavioral effects of bupropion and psychostimulants. *Eur J Pharmacol* **718**:370–375.
- Mucha RF and Iversen SD (1984) Reinforcing properties of morphine and naloxone revealed by conditioned place preferences: a procedural examination. *Psychopharmacology (Berl)* **82**:241–247.
- Ninkovic J and Bally-Cuif L (2006) The zebrafish as a model system for assessing the reinforcing properties of drugs of abuse. *Methods* **39**:262–274.
- Ninkovic J, Folchert A, Makhankov YV, Neuhauss SC, Sillaber I, Straehle U, and Bally-Cuif L (2006) Genetic identification of AChE as a positive modulator of addiction to the psychostimulant D-amphetamine in zebrafish. *J Neurobiol* **66**:463–475.
- O'Connor EC, Chapman K, Butler P, and Mead AN (2011) The predictive validity of the rat self-administration model for abuse liability. *Neurosci Biobehav Rev* **35**:912–938.
- Pain L, Oberling P, Sandner G, and Di Scala G (1996) Effect of propofol on affective state as assessed by place conditioning paradigm in rats. *Anesthesiology* **85**:121–128.

- Panlilio LV and Goldberg SR (2007) Self-administration of drugs in animals and humans as a model and an investigative tool. *Addiction* **102**:1863–1870.
- Parker MO, Brock AJ, Millington ME, and Brennan CH (2013) Behavioural phenotyping of casper mutant and 1-phenyl-2-thiourea treated adult zebrafish. *Zebrafish* **10**:466–471.
- Parker MO, Evans AM, Brock AJ, Combe FJ, Teh MT, and Brennan CH (2016) Moderate alcohol exposure during early brain development increases stimulus-response habits in adulthood. *Addict Biol* **21**:49–60.
- Pascual MM, Pastor V, and Bernabeu RO (2009) Nicotine-conditioned place preference induced CREB phosphorylation and Fos expression in the adult rat brain. *Psychopharmacology (Berl)* **207**:57–71.
- Pickens R, Muchow D, and DeNoble V (1981) Methohexital-reinforced responding in rats: effects of fixed ratio size and injection dose. *J Pharmacol Exp Ther* **216**:205–209.
- Platt DM, Rowlett JK, and Spealman RD (2008) Attenuation of cocaine self-administration in squirrel monkeys following repeated administration of the mGluR5 antagonist MPEP: comparison with dizocilpine. *Psychopharmacology (Berl)* **200**:167–176.
- Renier C, Faraco JH, Bourgin P, Motley T, Bonaventure P, Rosa F, and Mignot E (2007) Genomic and functional conservation of sedative-hypnotic targets in the zebrafish. *Pharmacogenet Genomics* **17**:237–253.
- Richendrfer H, Pelkowski SD, Colwill RM, and Creton R (2012) On the edge: pharmacological evidence for anxiety-related behavior in zebrafish larvae. *Behav Brain Res* **228**:99–106.
- Riehl R, Kyzar E, Allain A, Green J, Hook M, Monnig L, Rhymes K, Roth A, Pham M, Razavi R, et al. (2011) Behavioral and physiological effects of acute ketamine exposure in adult zebrafish. *Neurotoxicol Teratol* **33**:658–667.
- Rutten K, van der Kam EL, De Vry J, and Tzschentke TM (2011) Critical evaluation of the use of extinction paradigms for the assessment of opioid-induced conditioned place preference in rats. *Pharmacology* **87**:286–296.
- Sackerman J, Donegan JJ, Cunningham CS, Nguyen NN, Lawless K, Long A, Benno RH, and Gould GG (2010) Zebrafish behavior in novel environments: effects of acute exposure to anxiolytic compounds and choice of Danio rerio Line. *Int J Comp Psychol* **23**:43–61.
- Silverman HI (1972) The adverse effects of commonly used systemic drugs on the human eye. II. *Am J Optom Arch Am Acad Optom* **49**:335–362.
- Spyraki C, Fibiger HC, and Phillips AG (1982) Cocaine-induced place preference conditioning: lack of effects of neuroleptics and 6-hydroxydopamine lesions. *Brain Res* **253**:195–203.
- Stewart AM, Ullmann JF, Norton WH, Parker MO, Brennan CH, Gerlai R, and Kalueff AV (2015) Molecular psychiatry of zebrafish. *Mol Psychiatry* **20**:2–17.
- Suzuki T, Mori T, Tsuji M, Nomura M, Misawa M, and Onodera K (1999) Evaluation of the histamine H1-antagonist-induced place preference in rats. *Jpn J Pharmacol* **81**:332–338.
- Tzschentke TM (1998) Measuring reward with the conditioned place preference paradigm: a comprehensive review of drug effects, recent progress and new issues. *Prog Neurobiol* **56**:613–672.
- Tzschentke TM (2007) Measuring reward with the conditioned place preference (CPP) paradigm: update of the last decade. *Addict Biol* **12**:227–462.
- van der Kam EL, De Vry J, and Tzschentke TM (2009) The mGlu5 receptor antagonist 2-methyl-6-(phenylethynyl)pyridine (MPEP) supports intravenous self-administration and induces conditioned place preference in the rat. *Eur J Pharmacol* **607**:114–120.
- Vermoesen K, Serruys A-SK, Loyens E, Afrikanova T, Massie A, Schallier A, Michotte Y, Crawford AD, Esguerra CV, de Witte PAM, et al. (2011) Assessment of the convulsant liability of antidepressants using zebrafish and mouse seizure models. *Epilepsy Behav* **22**:450–460.
- Vitale MA, Chen D, and Kanarek RB (2003) Chronic access to a sucrose solution enhances the development of conditioned place preferences for fentanyl and amphetamine in male Long-Evans rats. *Pharmacol Biochem Behav* **74**:529–539.
- Wee S and Woolverton WL (2004) Evaluation of the reinforcing effects of atomoxetine in monkeys: comparison to methylphenidate and desipramine. *Drug Alcohol Depend* **75**:271–276.
- Wilcox KM, Rowlett JK, Paul IA, Ordway GA, and Woolverton WL (2000) On the relationship between the dopamine transporter and the reinforcing effects of local anesthetics in rhesus monkeys: practical and theoretical concerns. *Psychopharmacology (Berl)* **153**:139–147.
- Wilcox KM, Kimmel HL, Lindsey KP, Votaw JR, Goodman MM, and Howell LL (2005) In vivo comparison of the reinforcing and dopamine transporter effects of local anesthetics in rhesus monkeys. *Synapse* **58**:220–228.
- Wiley JL, Huffman JW, Balster RL, and Martin BR (1995) Pharmacological specificity of the discriminative stimulus effects of delta 9-tetrahydrocannabinol in rhesus monkeys. *Drug Alcohol Depend* **40**:81–86.
- Wong K, Stewart A, Gilder T, Wu N, Frank K, Gaikwad S, Suci C, Dileo J, Utterback E, Chang K, et al. (2010) Modeling seizure-related behavioral and endocrine phenotypes in adult zebrafish. *Brain Res* **1348**:209–215.
- Woolverton WL and Balster RL (1979) Reinforcing properties of some local anesthetics in rhesus monkeys. *Pharmacol Biochem Behav* **11**:669–672.
- Zang L, Morikane D, Shimada Y, Tanaka T, and Nishimura N (2011) A novel protocol for the oral administration of test chemicals to adult zebrafish. *Zebrafish* **8**:203–210.
- Zhang JJ, Ma X, and Yu LC (2012) Repeated paired-testing impairs extinction of morphine-induced conditioned place preference dependent on the inter-test interval in rats. *Neurosci Lett* **516**:72–74.
- Zhdanova IV, Wang SY, Leclair OU, and Danilova NP (2001) Melatonin promotes sleep-like state in zebrafish. *Brain Res* **903**:263–268.

Address correspondence to: Caroline H. Brennan, School of Biological and Chemical Sciences, Queen Mary, University of London, Mile End Rd., London, W10 5UW. E-mail: c.h.brennan@qmul.ac.uk
