

Special Section on Sexual Dimorphism in Neuroimmune Cells — Minireview

Sex-Dependent Mechanisms of Chronic Pain: A Focus on Microglia and P2X4R

✉ Katherine Halievski, ✉ Shahrzad Ghazisaeidi, and ✉ Michael W. Salter

Neurosciences and Mental Health Program, The Hospital for Sick Children, Toronto, Ontario, Canada (K.H., S.G., M.W.S.); The University of Toronto Centre for the Study of Pain, Toronto, Ontario, Canada (K.H., S.G., M.W.S.); and The Department of Physiology, University of Toronto, Toronto, Ontario, Canada (S.G., M.W.S.)

Received January 7, 2020; accepted February 25, 2020

ABSTRACT

For over two decades, purinergic signaling in microglia has persisted in the spotlight as a major pathomechanism of chronic pain. Of the many purinoreceptors, the P2X4R of the ionotropic family, has a well-described causal role underlying chronic neuropathic pain. This review will briefly examine microglial P2X4R signaling in the spinal cord as it relates to chronic pain through a historical lens, followed by a more in-depth examination of recent work, which has revealed major sex differences. We also discuss the generalizability of sex differences in microglial and P2X4R signaling in other pain conditions as well as in nonspinal regions. Finally, we speculate on remaining gaps in the literature as well as what can be done to address them with the

ultimate goal of using our collective knowledge to treat chronic pain effectively and in both sexes.

SIGNIFICANCE STATEMENT

Effective treatments are lacking for chronic pain sufferers, and this may be explained by the vast sex differences underlying chronic pain mechanisms. In this minireview, we focus on the roles of microglia and P2X4R in chronic pain, with specific attention to the circumstances under which these pathomechanisms differ between males and females. By delineating the ways in which pain occurs differently between the sexes, we can start developing successful therapies for all.

A Brief History of Microglial P2X4R Signaling in Pain Hypersensitivity

Chronic pain is a debilitating condition that affects up to 20% of the population and has an economic burden of \$43 billion per year in Canada alone (Lynch, 2011; Schopflocher et al., 2011). Unfortunately, existing treatments for chronic pain are not wholly effective and have significant side effects (Fornasari, 2017). Chronic pain is not phenotypically or mechanistically homogenous and may arise from many causes. To determine the best ways in which to treat patients with chronic pain, we can start first by examining the recognized

pathomechanisms of chronic pain. One well-defined pathomechanism is via purinergic signaling (Bernier et al., 2018). Purinergic receptors are activated by purine nucleosides or nucleotides, such as ATP, and include ionotropic P2X receptors, of which there are seven types, and metabotropic P2Y receptors, of which there are eight (Burnstock, 2018). The P2X4 receptor (P2X4R) class of receptor was first linked to chronic-pain signaling by Tsuda et al. (2003) and has been, to date, one of the best-characterized signaling pathways for different types of chronic pain (Inoue, 2019).

The initial evidence of P2X4R underlying chronic pain was the demonstration of reversal of hypersensitivity caused by traumatic peripheral nerve injury (PNI) by inhibiting P2X4Rs and increased P2X4R expression on microglia (Tsuda et al., 2003), a prominent cell type in the central nervous system. A functional consequence of increasing P2X4R expression and activation in microglia is the phosphorylation of p38 mitogen-activated protein kinase (MAPK), which induces transcription, translation, and release of brain-derived neurotrophic

Work of the authors is supported by Canadian Institutes of Health Research (CIHR) foundation grant [FDN-154336] to M.W.S.; M.W.S. is the Northbridge Chair in Paediatric Research at the Hospital for Sick Children. K.H. is supported by a L'Oréal Canada For Women in Science Research Excellence Fellowship, with the support of the Canadian Commission for UNESCO, and by the The Ontario Women's Health Scholars Award through the Ontario Ministry of Health and Long-Term Care.
<https://doi.org/10.1124/jpet.120.265017>

ABBREVIATIONS: BDNF, brain-derived neurotrophic factor; CSF-1, colony-stimulating factor 1; IL-6, interleukin-6; IRF, interferon regulatory factor; KCC2, K⁺-Cl⁻ cotransporter; LPS, lipopolysaccharide; NMDAR, N-methyl-D-aspartate receptor; p38-MAPK, p38 mitogen-activated protein kinase; PAG, periaqueductal gray; PGE-2, prostaglandin E2; PNI, peripheral nerve injury; P2X4R, purinergic receptor 4; ionotropic TNP-ATP, 2',3'-O-(2,4,6-Trinitrophenyl) adenosine 5'-triphosphate; TrkB, tropomyosin receptor kinase B; TLR4, toll-like receptor 4; VNUT, vesicular nucleotide transporter.

factor (BDNF) from microglia (Trang et al., 2009). Then, microglial BDNF serves as the messenger molecule that drives disinhibition and hyperexcitation of dorsal horn neurons (Coull et al., 2005), ultimately resulting in pain hypersensitivity. That the requisite BDNF derives from microglia is shown by the lack of pain hypersensitivity in mice in which the *Bdnf* gene is disrupted selectively in microglia (Sorge et al., 2015). BDNF activation of the tropomyosin receptor kinase B (TrkB) receptor in neurons leads to downregulation of the K⁺-Cl⁻ cotransporter KCC2 expression (Coull et al., 2003) and N-methyl-D-aspartate receptor (NMDAR) phosphorylation (Hildebrand et al., 2016). The spinal cord microglia P2X4R-BDNF to neuronal KCC2/NMDAR pathway is not only engaged following traumatic peripheral nerve injury but also mediates pain hypersensitivity caused by morphine (Ferrini et al., 2013) and in a model of herpetic pain (Matsumura et al., 2016). For a more in-depth overview of this signaling pathway of chronic pain, please see reviews on this topic (Salter and Beggs, 2014; Mapplebeck et al., 2017; Inoue, 2018).

In parallel to studies focusing on the P2X4R pathway within microglia as it relates to pain hypersensitivity caused by PNI, researchers linked aberrant microglial function more generally to pathologic pain (Marchand et al., 2005). Because P2X4Rs are strongly expressed by microglia, the two fields could then inform each other about possible overlapping mechanisms. A critical discovery that prompted the two fields to merge was the finding of an apparent sex specificity in the use of microglia during chronic pain; in a report by Sorge et al. (2011), the toll-like receptor 4 (TLR4) agonist lipopolysaccharide (LPS), which is primarily expressed by microglia in the central nervous system, caused pain behaviors in mice. However, this only occurred in males. Female mice showed no response to intrathecal LPS administration. These findings led the authors to further characterize the sex dependency of microglial P2X4R and BDNF pathway as it relates to chronic pain and to address the question of how female biology results in the same degree of pain hypersensitivity in the absence of a contributory role from microglia. Nevertheless, conditions under which microglia may mediate pain in females remain to be determined. In the following section, we describe what we have learned to date regarding specific nodes of the P2X4R sex-dependent pathway in chronic pain.

What Do We Know about Sex Differences in Spinal Microglia and P2X4R Signaling?

Since the discovery that P2X4R signaling in microglia is not an underlying driver of chronic pain hypersensitivity in females, where the pathway diverges between sexes has been identified (Sorge et al., 2015; Mapplebeck et al., 2018). There is a male-specific upregulation of P2X4R after PNI, which induces pain hypersensitivity, and this phenomenon is consistent across species and pain models. Using quantitative reverse-transcriptase polymerase chain reaction of the injured side of the dorsal spinal cord, an analysis at the mRNA level revealed that *P2rx4* transcripts were upregulated in male, but not in female, mice after PNI (Sorge et al., 2015). In a separate study, microglial P2X4R protein levels were measured in rats after nerve injury; fluorescence-activated cell sorting-isolated CD11b⁺ cells from the spinal cord showed upregulation of P2X4R via Western blot in males

but not females (Mapplebeck et al., 2018). Furthermore, in primary cultures of rat microglia, ATP stimulation alone increased P2X4R levels but, again, only in male-derived cells (Mapplebeck et al., 2018). Together, these data demonstrate that the P2X4R is upregulated in males but not in females after a chronic pain-inducing PNI.

The upregulation of the *P2rx4* message is controlled by the transcription factors interferon regulatory factor-5 (IRF5) and IRF8. IRF8 is an essential modulator of microglial reactivity, and it directly regulates IRF5 expression (Masuda et al., 2012, 2014). IRF5 then binds directly to the *P2rx4* promoter to drive gene expression (Masuda et al., 2014; Mapplebeck et al., 2018). When IRF5 is depleted in mouse spinal cord, *P2rx4* levels are not upregulated after PNI, and, correspondingly, such mice have reduced pain (Masuda et al., 2014). To address whether there are any sex differences in the levels of *Irf8* and *Irf5* mRNA during chronic pain, the spinal cords of males and females were probed. Surprisingly, *Irf8* and *Irf5* transcripts were upregulated equally in both sexes (Sorge et al., 2015). This suggested that the divergence between sexes in *P2rx4* mRNA expression occurs only after *Irf8/5* regulation. Indeed, IRF5 binds to the *P2rx4* promoter region with a higher affinity in male- compared with female-derived primary microglia as well as in the whole spinal cord of male rats compared with females (Mapplebeck et al., 2018). Thus, differential binding of IRF5 at the *P2rx4* promoter region in males versus females appears to underlie the sexually dimorphic upregulation of microglial P2X4R after PNI. While the mechanisms regulating the differential binding of IRF5 to *P2rx4* remain unknown, one possibility may be due to epigenetic modifications of the *P2rx4* promoter.

It is possible that under certain circumstances the *P2rx4* promoter may become accessible even in females. For example, Matsumura et al. (2016) developed a female mouse model of postherpetic neuralgia, a painful condition associated with complications from shingles (Hadley et al., 2016). They showed that inhibiting P2X4R with NP-1815-PX blunted pain in females with herpetic pain hypersensitivity. In this model, there was microgliosis in the dorsal horn, and the BDNF sequestering agent, TrkB-Fc, also suppressed the pain hypersensitivity, implying that the microglial P2X4R-BDNF pathway was activated. Whether there are other circumstances in which P2X4R may be upregulated in microglia in females, such as opioid-induced hyperalgesia, remains to be determined.

The endogenous ligand for P2X4Rs expressed in spinal microglia is presumed to be ATP (Tsuda et al., 2005). The source of ATP has been determined to be neurons within the dorsal horn itself rather than primary sensory neurons, microglia, or astrocytes (Masuda et al., 2016). Levels of ATP are heightened in the spinal cord after PNI because of an increase of vesicular nucleotide transporter (VNUT) in dorsal horn neurons, at least in males; mice lacking VNUT do not show PNI-induced increases in ATP, and hypersensitivity is suppressed (Masuda et al., 2016). Conversely, viral-driven expression of VNUT in dorsal horn neurons in VNUT-deficient mice rescues the increase in ATP and the hypersensitivity induced by PNI. Currently, it is unknown whether the VNUT increase is sex-differentiated. Thus, it would be interesting to determine whether this is true in females and what biological consequences this might have.

Microglia from male, but not female, rats exhibit ATP-stimulated increases phosphorylation of p38-MAPK and an increase in the expression and release of BDNF (Mapplebeck

et al., 2018), consistent with male-specific expression of P2X4Rs. In a separate study, a male-specific increase in phosphorylated p38-MAPK was also observed in microglia, and inhibition of p38-MAPK via intrathecal drug administration reduced PNI- and formalin-induced pain in male, but not female, rodents (Taves et al., 2016). It was later demonstrated that the key isoform driving pain in a sex-dependent manner is p38 α (Luo et al., 2018). Together, these data suggest that male microglia are uniquely prepared to mediate pain-inducing signaling. Indeed, male microglia that have been pre-exposed to ATP and then applied to naïve rat spinal cord *in vivo* induce a pain-like response, but female microglia that have been pretreated with ATP do not have this effect (Mapplebeck et al., 2018). It is not clear whether and how female microglia respond to ATP stimulation at the level of molecular mediators. Nevertheless, female microglia may be involved in the maintenance of pain in females under certain conditions, as described below (Vacca et al., 2014; Yang et al., 2015; Peng et al., 2016).

Given the distinct microglial response to ATP in males and females, it was important to determine whether outcomes downstream of microglia signaling—i.e., the changes in neurons that underlie chronic pain—also occur in females, as the sexes exhibit indistinguishable behavioral sensitization after PNI. Earlier evidence implied that neurons underlie chronic pain in males and females based on the finding that inhibiting NMDARs via intrathecal administration of AP5 suppressed pain similarly in females and males (Sorge et al., 2015). However, whether the loss of KCC2 function is also involved in females was not known until recently. Increased BDNF from microglia culminates in the reduction of KCC2 expression with a consequent loss of chloride gradient regulation of dorsal horn neurons in males (Coull et al., 2005). Recently, it was shown that KCC2 is downregulated in females as it is in males (Mapplebeck et al., 2019) despite the lack of congruency in the steps immediately preceding the male-typical mechanism of chronic pain, such as the requirement for microglial BDNF. Thus, we now know that after peripheral injury, microgliosis occurs in males and females and that there is an upregulation of transcription factors IRF8 and IRF5, but the critical point of divergence between the sexes is the binding of IRF5 to the *P2rx4* promoter. In the framework of this canonical microglia-P2X4R pathway, typically seen in males, and the microglia-P2X4R independent pathway, typically seen in females, converge at the level of KCC2 downregulation. It remains unclear how females achieve the KCC2 downregulation that results in pain hypersensitivity, although the adaptive immune system is hypothesized to be involved (Sorge et al., 2015). For a pictorial representation of this pathway in males and females, see Mapplebeck et al. (2016, 2017).

Several questions remain about sex differences in the sequelae leading up to P2X4R expression and after BDNF binding to TrkB on neurons. For example, sensory neurons express colony-stimulating factor 1 (CSF-1) after nerve injury, and this induces microglial activation and proliferation as well as mechanical hypersensitivity in males (Guan et al., 2016). In a recent report, Yu et al. (2020) found that CSF-1 from primary sensory neurons also induces proliferation of macrophages in the dorsal root ganglion after peripheral nerve injury in males but not in females. To what extent microgliosis in the female spinal cord (Sorge et al., 2015; Mapplebeck et al., 2018) is mediated by CSF-1 after nerve injury, and what the functions of those microglia are, is still an open question.

The role of sex hormones is also important to address. Though there is evidence that high-circulating male sex hormones are not required to induce pain from male ATP-stimulated microglia (Mapplebeck et al., 2018), there is evidence that *in vivo* testosterone treatment to females does enable the P2X4R-mediated microglia response to occur, at least in a chronic inflammatory model of pain (Sorge et al., 2015). Differences in pain models may explain why sex hormones may be critical for a microglia-dependent phenotype in one case but not another—more work certainly needs to be done to clarify these processes.

Finally, it continues to be a critical point of interest regarding whether P2X4R sex differences in chronic pain also occur in humans experiencing pain and whether we can use this knowledge to develop treatments specific for each individual. It is possible that P2X4R signaling in the context of chronic pain is sexually dimorphic in humans, as surface P2X4R expression on blood eosinophils is higher in males compared with females during healthy conditions (Paalme et al., 2019). Moreover, *P2RX4* level is elevated in circulating leukocytes after moderate exercise in chronic fatigue syndrome patients with comorbid fibromyalgia syndrome (Light et al., 2009). Whether this finding is related to the pain symptoms *per se* remains to be determined. Given the lack of studies examining P2X4R in humans in the context of chronic pain, it is imperative for the field to fill this gap if we are to translate knowledge from rodent studies to the clinic. Particularly in the context of this review, it will be interesting to determine whether expression of P2X4R increases in a sexually dimorphic manner during chronic pain in humans as it does in rodent microglia (Mapplebeck et al., 2018).

Generalizability of Spinal Microglial and/or P2X4R Dependency in Different Pain Conditions

The evidence above is based on studies that detailed sex-specific behavioral sensitization following PNI; that is, the P2X4R-mediated pain mechanisms of pain hypersensitivity. In this section, we describe evidence as to whether sex differences in the P2X4R-mediated pathomechanism occur in pain states other than nerve injury-induced pain (also see Table 1) as well as more generally examine the evidence for microglia involvement in pain in females.

Hyperalgesic priming is a model of chronic pain in which an initial pain-inducing injury causes a heightened response to a subsequent injury, even after the first one has apparently healed. Specifically, once animals with the initial injury—the priming injury—no longer show a pain-like response, a second injury, which normally would not cause pain hypersensitivity, is applied. In “primed” animals, the normally subthreshold second stimulus results in hypersensitivity, indicating that there has been a change in the nociceptive circuitry to make the animal now sensitive to a normally nonpainful stimulus.

Using a hyperalgesic model with intraplantar soluble interleukin-6 (IL-6) receptor as the molecule to establish priming and prostaglandin E2 (PGE-2) to instigate it, Paige et al. (2018) examined sex differences in the role of P2X4R and p38-MAPK, the same signaling mediators that are sex-differentiated in chronic neuropathic and inflammatory pain (see *What Do We Know about Sex Differences in Spinal Microglia and P2X4R Signaling?*). They found that when

TABLE 1

The role of P2X4R in various pain conditions

Pain Condition	Subjects Studied	Key Finding	Reference
Activity-induced muscle pain	Male and female C57BL/6 mice	Immunohistochemical staining shows P2X4R is upregulated in muscle macrophages, but the data were not split by sex. Intramuscular injection of the P2XR4 blocker 5-BDBD (5-(3-Bromophenyl)-1,3-dihydro-2H-benzofuro[3,2-e]-1,4-diazepin-2-one), or genetic knockdown of P2X4R with a macrophage-specific miRNA, prevented hyperalgesia in both sexes.	Oliveira-Fusaro et al., 2020
Peripheral diabetic neuropathy	Male Wistar rats	P2X4R protein is upregulated in the dorsal root ganglion, as detected by Western blot. Intrathecal or ganglionic injection of a P2X4R antagonist (PSB-15417), or of intrathecal P2X4R antisense oligodeoxynucleotide, reversed mechanical hypersensitivity.	Teixeira et al., 2019
Neuropathic pain (spared nerve injury and chronic constriction injury)	Male and female Sprague-Dawley rats	Male rats, but not females, with spared nerve injury had increased P2X4R protein expression in microglia of the dorsal spinal cord. P2X4R antagonist (TNP-ATP) attenuated allodynia in rats with chronic constriction injury but only in males and not females.	Mapplebeck et al., 2018
Inflammatory pain (intraplantar injection of complete Freund's adjuvant)	Male and female C57BL/6 mice	P2X4R expression was increased in sensory neurons of the dorsal root ganglion in male mice; females were not tested in this experiment. Nevertheless, P2X4R-deficient female mice did not exhibit hypersensitivity to CFA, unlike their wild-type counterparts.	Lalisse et al., 2018
Hyperalgesic priming (intraplantar injection of soluble interleukin-6 receptor followed by prostaglandin E2)	Male and female Swiss Webster mice	Intrathecal injection of TNP-ATP (P2X4R inhibitor) suppressed the initial response to soluble IL-6 receptor in both sexes. Intrathecal TNP-ATP blocked subsequent hyperalgesia to prostaglandin E2 only when given at the time of initial IL-6 receptor administration and only in males. TNP-ATP did not reverse established priming when given after prostaglandin E2 administration.	Paige et al., 2018
Partial sciatic nerve ligation	Male and female C57BL/6 mice	No sex difference was found in <i>P2rx4</i> transcript levels in purified peripheral afferent neurons or in isolated microglia after nerve injury.	Lopes et al., 2017
Herpetic pain	Female C57BL/6 mice	P2X4R mRNA level in the spinal dorsal horn and immunoreactivity in spinal microglia were increased. Selective P2X4R antagonist (NP-1815-PX) had an antiallodynic effect when administered intrathecally.	Matsumura et al., 2016
Spared nerve injury	Male and female CD1 mice	<i>P2rx4</i> mRNA level is upregulated in dorsal horn spinal cord of male mice but not in females.	Sorge et al., 2015
Cancer-induced bone pain	Female Sprague-Dawley rats	P2X4R mRNA and protein was upregulated in spinal microglia. Silencing receptor activity with intrathecal P2X4R siRNA attenuated mechanical hypersensitivity.	Jin et al., 2014
Inflammatory pain (intraplantar injection of complete Freund's adjuvant, carageenan, or formalin)	Male C57BL/6 mice	P2X4R-deficient mice show reduced hypersensitivity and impaired production of PGE2 after intraplantar complete Freund's adjuvant or carageenan. P2X4R knockout mice also exhibit modest reductions to acute formalin-induced pain but hypersensitivity at baseline on the hotplate assay.	Ulmann et al., 2010
Chronic fatigue syndrome comorbid with fibromyalgia syndrome	Male and female human subjects	After moderate exercise, chronic fatigue syndrome and chronic fatigue syndrome–fibromyalgia syndrome patients showed elevated level of P2RX4 in circulating leukocytes compared with control subjects. Baseline P2RX4 expression did not differ between patients and controls.	Light et al., 2009
Inflammatory pain (intraplantar injection of complete Freund's adjuvant or formalin or intraperitoneal injection of acetic acid) and neuropathic pain (L4 spinal nerve transection)	Male C57BL/6 mice	P2X4R knockout mice show reduced mechanical, but not cold, allodynia caused by nerve injury. P2X4R knockout also exhibit attenuated inflammatory pain. P2X4R knockout does not affect baseline mechanical or thermal measures, nor does it affect acute models of pain (visceral acetic acid or intraplantar formalin).	Tsuda et al., 2009
Inflammatory pain (formalin injection to dorsal paw)	Male Lewis rats	P2X4R + immunoreactivity was increased in the spinal cord dorsal horn 7 days after intraplantar formalin injection.	Guo et al., 2005
Spinal nerve injury	Male Wistar rats	P2X4R protein was selectively upregulated in spinal cord microglia. Knockdown of P2X4R with P2rx4 antisense oligodeoxynucleotide or P2X4R inhibition with TNP-ATP suppressed mechanical allodynia. Intraspinal injection to naïve rats of ATP-stimulated microglia produced allodynia.	Tsuda et al., 2003

P2X4R or p38-MAPK signaling was inhibited at the time of the initial injury via intrathecal TNP-ATP (2',3'-O-(2,4,6-Trinitrophenyl) adenosine 5'-triphosphate) or skepinone, respectively, the subsequent injury no longer resulted in a hyperalgesic response, indicating that the changes to nociceptive circuitry due to the initial injury were prevented. Similar to the nerve injury pain models, this intervention at P2X4R and p38-MAPK signaling suppressed the hyperalgesic responses only in males. Notably, it was critical that the intervention be administered at the time of the initial injury, as later administration at the time of the second injury did not prevent the hyperalgesia. A further point regarding the mechanism leading to persistent pain in this model of hyperalgesic priming is that the P2XR blocker, TNP-ATP, did suppress the development of the initial injury response in females and males, suggesting that P2X4R inhibition did have an effect. But, for females, this effect was only transient as they still remained "primed" for the second injury—i.e., TNP-ATP did not prevent the long-term consequences. As TNP-ATP is not totally specific to P2X4R (North and Jarvis, 2013), the prevention of the initial sensitization and/or of the priming may be mediated by other P2XRs. Thus, based on this work, P2X4R and p38-MAPK signaling is important in the establishment of hyperalgesic priming but only in males, whereas both of these signaling molecules are important for males and females for the expression of the initial injury.

To address the downstream step in the P2X4R–p38-MAPK pathway of pain within hyperalgesic priming, the same group of researchers asked whether BDNF signaling underlies the sex-dependent response in hyperalgesic priming (Moy et al., 2018). Again, using IL-6 as the inducer of priming and PGE-2 to unmask the priming, they found that inhibiting BDNF signaling using intrathecal TrkB-Fc, which sequesters BDNF, prevented priming in male mice only. When this experiment was done in rats, the sex difference disappeared, and females were also rescued from hyperalgesic priming. The mechanisms underlying this form of hyperalgesic priming then must be different between species, at least for females. Given that microglial BDNF is critical for male-specific neuropathic pain (Sorge et al., 2015), the Moy et al. (2018) study asked whether microglial BDNF was also required for the acute and priming phases of IL-6- and PGE-2-induced pain but found that removing BDNF from microglia had no observable effect on the pain phenotype. Together, these data further emphasize the need to study any purported pain mechanism across multiple pain conditions and species to determine its generalizability as well as unique response profile to different pathological states.

In a different pain-priming paradigm, after hindpaw incision during early life followed by a second incision in later life, animals show enhanced hyperalgesia and longer lasting pain (Walker et al., 2009). Moriarty et al. (2019) used this paradigm to study the sex-dependent involvement of microglia and p38-MAPK signaling in the establishment of incision-primed hyperalgesia in rats. They found that when microglia were inhibited at the time of neonatal incision injury via systemic minocycline treatment, or intrathecal p38-MAPK inhibition, using SB203580, incision in later life no longer caused enhanced hyperalgesia, but this was only in males and not in females. The authors further linked microglia to this sex-specific phenomenon by showing that after the second

incision in adulthood, gene expression of microglial markers increased in females to a greater extent than in males, who were rescued by early-life minocycline treatment. Together with the above examples, this suggests that even during early life, male rodents use microglia in priming injury to produce persistent and latent pain states. It will be interesting to determine whether P2X4R signaling is also involved in neonatal incision pain priming, as it is involved in adult primed states (Paige et al., 2018) and in chronic pain conditions (Inoue, 2019).

P2X4Rs could be involved in pain hypersensitivity independent of microglia via expression and activation of these receptors in other types of cells (Ulmann et al., 2010). Under such alternative (non-microglia-dependent) situations, P2X4R may then be involved in driving pain in females. For example, Lalisie et al. (2018) showed that a whole-body genetic knockout of P2X4R in a chronic inflammatory model, intraplantar complete Freund's adjuvant, prevented mechanical hypersensitivity in both males and females. The authors concluded that primary sensory neurons are the key source of P2X4R, which induces hypersensitivity in this inflammatory model. P2X4Rs have also been found to mediate activity-induced muscle pain via muscle macrophages; blocking P2X4R signaling in the muscle, or specifically in muscle macrophages, reduced hyperalgesia in both male and female mice (Oliveira-Fusaro et al., 2020). Thus, depending on the particular set of conditions and initiating stimuli, females also show pain hypersensitivity mediated by P2X4Rs.

Microglia may also be involved in females to drive pain under certain conditions. One report using a chronic constriction injury model of pain showed that microglial activation is correlated to hypersensitivity, with female mice showing longer maintenance of activated microglia and pain compared with males (Vacca et al., 2014). Additionally, in a spinal transection model of neuropathic pain, females were protected from pain after ablation of CX3C chemokine receptor 1 (CX3CR1)-positive cells, which include microglia and myeloid cells (Peng et al., 2016). Interestingly, during early pregnancy, female mice appear to rely on microglia to drive inflammatory and neuropathic pain, as minocycline was capable of reversing pain during this window (Rosen et al., 2017). In a separate study, mice with nerve injury-induced pain did not exhibit sex differences in gene expression in purified sensory neurons nor in microglia, including for *P2rx4* transcripts (Lopes et al., 2017). Together, these data suggest that sex differences in the involvement of P2X4R signaling, and possibly microglia more broadly in pain, are dependent on the type of pain condition being examined and the state of the animal itself. Although microglia and P2X4R signaling show a sex-dependent involvement in many chronic pain conditions, it remains essential to elucidate the details on under which conditions there are sex differences and what underlies the ability of female microglia to contribute to pain in one circumstance but not another.

Sex Differences in Microglia and P2X4R Signaling

As described above, spinal microglia and P2X4R signaling underlie multiple pain states and, in many cases, are sex-differentiated. Other than the spinal cord, central nervous system regions involved in pain-signal transmission also

exhibit basal sex differences in microglia, which may further underlie sex-specific responses after PNI.

Microglia have multiple functions in the healthy and diseased central nervous system (Kettenmann et al., 2011; Salter and Stevens, 2017). They can respond to the stimuli such as injury by producing proinflammatory cytokines, and they also can produce anti-inflammatory cytokines and show neuroprotective effects (Lucin and Wyss-Coray, 2009). However, sex differences in microglia state during development or after an injury remain unclear. During early development in mice and rats, male and female brain microglia differ in number (Mouton et al., 2002; Schwarz et al., 2012; Lenz et al., 2013) and in expression of cytokines and chemokines (Schwarz et al., 2012). The sex differences in these early-life stages might be due to the neonatal testosterone surge because there are limited sex differences in microglial density and morphology prior to this developmental period (Schwarz et al., 2012). Neonatal androgens are converted to estradiol by the local aromatase enzyme in the brain and result in brain masculinization and defeminization (McCarthy et al., 2009). Just after birth, during the neonatal testosterone surge (Clarkson and Herbison, 2016), microglia in males have higher chemokine levels of chemokine (C-C motif) ligand 4 and 20 compared with females, which may drive an increase in microglial cell number in males at early ages (Schwarz et al., 2012). Indeed, 4 days after birth, neonatal male mice have more microglia in the CA1 and CA3 of the hippocampus, the dentate gyrus, the parietal cortex, and the amygdala (Schwarz et al., 2012). Thus, the basal microglial reprogramming window may overlap with the critical hormonal period, specifically with the androgen surge. In the preoptic area of the rat brain, a region responsible for adult male sexual behavior, males exhibited increased secretion of a proinflammatory mediator (PGE-2) from microglia compared with females because of the higher level of estradiol in males during this critical period; this was necessary and sufficient for the masculinization of neuronal dendritic spine density and male copulatory behavior (Amateu and McCarthy, 2004; Lenz et al., 2013; Rahimian et al., 2019). Therefore, microglia that express PGE-2 and its receptor might be essential for studying developmental sex differences in the nervous system. Studies indicate the sex difference in microglia number exists after puberty; in adolescence and adulthood, female mice have higher number of total (Mouton et al., 2002) and activated (Schwarz et al., 2012) microglia than males.

With recent advancements in sequencing technology, our understanding of sex differences in microglia has been transformed. Multiple studies on microglia transcriptome (Guneykaya et al., 2018; Villa et al., 2018), translatoome (Kang et al., 2018), and proteome (Guneykaya et al., 2018) showed that microglia in the healthy adult brain is sexually differentiated. Two transcriptomic studies in mice found that male-derived microglia are more proinflammatory (Guneykaya et al., 2018; Villa et al., 2018) and that female-derived microglia exhibit a neuroprotective-like transcriptome (Villa et al., 2018). On the other hand, a different study found that female microglia express more genes involved in the inflammatory response (Thion et al., 2018). Although all studies showed that microglia are sexually differentiated, whether males or females differ in proinflammatory transcriptome signature is debated. Such contrasting findings highlight the sensitivity of microglia transcriptome to the technical isolation, developmental

stage, and the source of microglia. For reviews on this topic see Hanamsagar and Bilbo (2016), Kodama and Gan (2019), Villa et al. (2019).

Interestingly, Villa et al. (2018) showed that female microglia maintain a sex-specific transcriptome when implanted into the male brain, indicating the programmed long-term gene regulation in these cells irrespective of the circulating hormonal environment. Therefore, it is interesting to understand the mechanisms of gene regulation such as epigenetic modulation in the context of organizational sex differences in microglia. Despite the lack of consistency in the directionality of the sex differences, all studies to date comparing male and female microglia have unequivocally found sex differences in the transcriptomic signature, and these depend on age.

Environmental stimuli such as presence of LPS or antibiotic treatment further influence a sex-specific response in the microglial transcriptome. The study by Hanamsagar et al. (2017) showed that intraperitoneal injection of LPS to mice accelerates male microglial maturation. However, the same LPS challenge did not affect female-derived microglia (Hanamsagar et al., 2017). Other environmental stimuli, such as antibiotic treatment or the effect of the maternal microbiome on microglial maturation, have been studied. Thion et al. (2018) observed maternal microbiome depletion induced a sex-specific response. Male microglia were more vulnerable to the absence of a microbiome during early developmental stages, but female microglia were more sensitive in adulthood (Thion et al., 2018). Thus, environmental stimuli affect microglial function in a sex-specific way depending on the developmental stage.

The basal sex difference in purinergic receptor expression may determine the phenotypic state of microglia. Guneykaya et al. (2018) showed that in the healthy mouse brain, translation of purinergic receptors P2X4R, P2X7R, and P2Y12R had significantly higher levels in microglia in males compared with females. In a separate study, Crain et al. (2009) showed a sex difference in the expression of multiple purinergic receptors in freshly isolated microglia cells from mouse brain. Specifically, females express lower levels of *P2rx1*, *P2rx4*, *P2ry12*, and *P2ry13* in adulthood and higher levels of *P2rx3* (adulthood) and *P2rx5* and *P2ry4* (only at early ages, postnatal day 3) compared with males (Crain et al., 2009). Sex-dependent P2X4R expression might explain the sexually distinct pathways of P2X4R involvement in painful or disease conditions. For example, a study in ischemic stroke recovery showed that global knockout of P2X4R provided acute neuroprotection, i.e., reduced infarct volume, in both sexes compared with the wild-type mice (Verma et al., 2017). However, when the P2X4R knockout was specific to myeloid cells, only females showed acute neuroprotection after stroke. Observed sex differences were not related to the activational effects of estrogen, suggesting this sex difference is under the influence of the organizational effects of gonadal hormones or sex chromosomes (Verma et al., 2017).

Sex differences in brain microglia as they relate to pain modulation have been studied in the rat periaqueductal gray (PAG). Doyle et al. (2017) showed that naïve, female rat microglia have a more activated morphology compared with male despite no sex difference in the number of microglia in PAG. They also examined how these basal sex differences remodel after activation of TLR4, which is primarily expressed

in microglia. Interestingly, the sex difference in microglial activation at baseline potentiated after TLR4 activation was induced with LPS treatment. Doyle et al. (2017) also concluded that microglial TLR4 in the PAG contributes to the attenuated response to morphine observed in females, as blocking TLR4 in the PAG enhanced morphine efficacy in females but not in males. This underlying sex-differentiated neural response may explain why morphine is less effective in females compared with males (Cepeda and Carr, 2003).

Collectively, multiple studies show sex differences in microglia during healthy states and across different brain regions. These sex differences are sensitive to age, developmental stage, and environment. In a disease situation, there are sex differences in number, morphology, and gene expression of microglia remodels. To fully understand the role of microglia in disease conditions, such as chronic pain, it is crucial to know the basal state of the sex difference as well as the sex-specific responses to environmental challenges.

Summary

Microglia are critical for maintaining a healthy nervous system (Salter and Stevens, 2017), and when these cells malfunction because of disease or injury, they can cause pathological conditions such as chronic pain. As sex differences are normally present in microglia cell number, morphology, and gene expression, it stands to reason that sex differences are similarly present in microglia-dependent pathologies. Indeed, microglia are disease-mediating in models of chronic inflammatory and neuropathic pain but in a sex-specific manner. As reviewed here, data from studies link microglial dysfunction to chronic pain in only male rodents. To date, there has been a focus on the purinergic receptor P2X4R as the male-biased microglial mediator of chronic pain. Though there are a few examples of microglia possibly being involved in chronic pain in females, there is no report of microglial inhibition reducing pain in females but not in males. Thus, microglia-dependent chronic pain is male-biased, and females likely use alternative pathways, potentially T lymphocytes; currently, these female-biased pathways are understudied, and there is a great need to understand them further.

Additional noteworthy gaps remain in the area of sex differences in microglia-P2X4R signaling of chronic pain. What role do sex hormones play? What causes IRF5 to differentially bind to the *P2rx4* promoter between sexes after nerve injury? What makes P2X4R/microglia involved in female pain under certain conditions but not others? Do microglia in the female dorsal spinal cord expand after nerve injury because of CSF-1 as in males? And what role do expanded microglia in females play, if not pain? It is also of great interest to the broader pain field how, where, and under which circumstances P2X4Rs and microglia underlie pathological pain in humans, and whether these contributions are sexually differentiated.

We hope that by continuing the discussion of sex-specific mechanisms of chronic pain, researchers will recognize that it is vital to study and report on both sexes in preclinical work so that, ultimately, clinical therapies for chronic pain are designed for efficacy in every person.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Halievski, Ghazisaeidi, Salter.

References

- Amateau SK and McCarthy MM (2004) Induction of PGE2 by estradiol mediates developmental masculinization of sex behavior. *Nat Neurosci* **7**:643–650.
- Bernier LP, Ase AR, and Séguéla P (2018) P2X receptor channels in chronic pain pathways. *Br J Pharmacol* **175**:2219–2230.
- Burnstock G (2018) Purine and purinergic receptors. *Brain Neurosci Adv* **2**: 2398212818817494.
- Cepeda MS and Carr DB (2003) Women experience more pain and require more morphine than men to achieve a similar degree of analgesia. *Anesth Analg* **97**: 1464–1468.
- Clarkson J and Herbison AE (2016) Hypothalamic control of the male neonatal testosterone surge. *Philos Trans R Soc Lond B Biol Sci* **371**:20150115.
- Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, Gravel C, Salter MW, and De Koninck Y (2005) BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature* **438**:1017–1021.
- Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sîk A, De Koninck P, and De Koninck Y (2003) Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature* **424**:938–942.
- Crain JM, Nikodemova M, and Watters JJ (2009) Expression of P2 nucleotide receptors varies with age and sex in murine brain microglia. *J Neuroinflammation* **6**:24.
- Doyle HH, Eidsen LN, Sinkiewicz DM, and Murphy AZ (2017) Sex differences in microglia activity within the periaqueductal gray of the rat: a potential mechanism driving the dimorphic effects of morphine. *J Neurosci* **37**: 3202–3214.
- Ferrini F, Trang T, Mattioli TA, Laffray S, DelGuidice T, Lorenzo LE, Castonguay A, Doyon N, Zhang W, Godin AG, et al. (2013) Morphine hyperalgesia gated through microglia-mediated disruption of neuronal Cl⁻ homeostasis. *Nat Neurosci* **16**: 183–192.
- Fornasari D (2017) Pharmacotherapy for neuropathic pain: a review. *Pain Ther* **6** (Suppl 1):25–33.
- Guan Z, Kuhn JA, Wang X, Colquitt B, Solorzano C, Vaman S, Guan AK, Evans-Reinsch Z, Braz J, Devor M, et al. (2016) Injured sensory neuron-derived CSF1 induces microglial proliferation and DAP12-dependent pain. *Nat Neurosci* **19**: 94–101.
- Guneykaya D, Ivanov A, Hernandez DP, Haage V, Wojtas B, Meyer N, Maricos M, Jordan P, Buonfiglioli A, Gielniewski B, et al. (2018) Transcriptional and translational differences of microglia from male and female brains. *Cell Rep* **24**: 2773–2783.e6.
- Guo LH, Trautmann K, and Schluesener HJ (2005) Expression of P2X4 receptor by lesional activated microglia during formalin-induced inflammatory pain. *J Neuroimmunol* **163**:120–127.
- Hadley GR, Gayle JA, Ripoll J, Jones MR, Argoff CE, Kaye RJ, and Kaye AD (2016) Post-herpetic neuralgia: a review [published correction appears in *Curr Pain Headache Rep* (2016) **20**:28]. *Curr Pain Headache Rep* **20**:17.
- Hanamsagar R, Alter MD, Block CS, Sullivan H, Bolton JL, and Bilbo SD (2017) Generation of a microglial developmental index in mice and in humans reveals a sex difference in maturation and immune reactivity [published correction appears in *Glia* (2017) **66**:460]. *Glia* **65**:1504–1520.
- Hanamsagar R and Bilbo SD (2016) Sex differences in neurodevelopmental and neurodegenerative disorders: focus on microglial function and neuroinflammation during development. *J Steroid Biochem Mol Biol* **160**:127–133.
- Hildebrand ME, Xu J, Dedek A, Li Y, Sengar AS, Beggs S, Lombroso PJ, and Salter MW (2016) Potentiation of synaptic GluN2B NMDAR currents by fyn kinase is gated through BDNF-mediated disinhibition in spinal pain processing. *Cell Rep* **17**: 2753–2765.
- Inoue K (2018) A state-of-the-art perspective on microglial pain. *Open Biol* **8**.
- Inoue K (2019) Role of the P2X4 receptor in neuropathic pain. *Curr Opin Pharmacol* **47**:33–39.
- Jin XH, Wang LN, Zuo JL, Yang JP, and Liu SL (2014) P2X4 receptor in the dorsal horn partially contributes to brain-derived neurotrophic factor oversecretion and toll-like receptor-4 receptor activation associated with bone cancer pain. *J Neurosci Res* **92**:1690–1702.
- Kang SS, Ebbert MTW, Baker KE, Cook C, Wang X, Sens JP, Kocher JP, Petrucelli L, and Fryer JD (2018) Microglial translational profiling reveals a convergent APOE pathway from aging, amyloid, and tau. *J Exp Med* **215**: 2235–2245.
- Kettenmann H, Hanisch UK, Noda M, and Verkhratsky A (2011) Physiology of microglia. *Physiol Rev* **91**:461–553.
- Kodama L and Gan L (2019) Do microglial sex differences contribute to sex differences in neurodegenerative diseases? *Trends Mol Med* **25**:741–749.
- Lalisse S, Hua J, Lenoir M, Linck N, Rassenclren F, and Ulmann L (2018) Sensory neuronal P2RX4 receptors controls BDNF signaling in inflammatory pain. *Sci Rep* **8**:964.
- Lenz KM, Nugent BM, Haliyur R, and McCarthy MM (2013) Microglia are essential to masculinization of brain and behavior. *J Neurosci* **33**:2761–2772.
- Light AR, White AT, Huguen RW, and Light KC (2009) Moderate exercise increases expression for sensory, adrenergic, and immune genes in chronic fatigue syndrome patients but not in normal subjects. *J Pain* **10**:1099–1112.
- Lopes DM, Malek N, Edey M, Jager SB, McMurray S, McMahan SB, and Denk F (2017) Sex differences in peripheral not central immune responses to pain-inducing injury. *Ki Rep* **7**:16460.
- Lucin KM and Wyss-Coray T (2009) Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron* **64**:110–122.

- Luo X, Fitzsimmons B, Mohan A, Zhang L, Terrando N, Kordasiewicz H, and Ji RR (2018) Intrathecal administration of antisense oligonucleotide against p38 α but not p38 β MAP kinase isoform reduces neuropathic and postoperative pain and TLR4-induced pain in male mice. *Brain Behav Immun* **72**:34–44.
- Lynch ME (2011) The need for a Canadian pain strategy. *Pain Res Manag* **16**:77–80.
- Mapplebeck JC, Beggs S, and Salter MW (2016) Sex differences in pain: a tale of two immune cells. *Pain* **157** (Suppl 1):S2–S6.
- Mapplebeck JC, Beggs S, and Salter MW (2017) Molecules in pain and sex: a developing story. *Mol Brain* **10**:9.
- Mapplebeck JCS, Dalgarno R, Tu Y, Moriarty O, Beggs S, Kwok CHT, Halievski K, Assi S, Mogil JS, Trang T, et al. (2018) Microglial P2X4R-evoked pain hypersensitivity is sexually dimorphic in rats. *Pain* **159**:1752–1763.
- Mapplebeck JCS, Lorenzo LE, Lee KY, Gauthier C, Muley MM, De Koninck Y, Prescott SA, and Salter MW (2019) Chloride dysregulation through down-regulation of KCC2 mediates neuropathic pain in both sexes. *Cell Rep* **28**: 590–596.e4.
- Marchand F, Perretti M, and McMahon SB (2005) Role of the immune system in chronic pain. *Nat Rev Neurosci* **6**:521–532.
- Masuda T, Iwamoto S, Yoshinaga R, Tozaki-Saitoh H, Nishiyama A, Mak TW, Tamura T, Tsuda M, and Inoue K (2014) Transcription factor IRF5 drives P2X4R+ reactive microglia gating neuropathic pain. *Nat Commun* **5**:3771.
- Masuda T, Ozono Y, Mikuriya S, Kohro Y, Tozaki-Saitoh H, Iwatsuki K, Uneyama H, Ichikawa R, Salter MW, Tsuda M, et al. (2016) Dorsal horn neurons release extracellular ATP in a VNUT-dependent manner that underlies neuropathic pain. *Nat Commun* **7**:12529.
- Masuda T, Tsuda M, Yoshinaga R, Tozaki-Saitoh H, Ozato K, Tamura T, and Inoue K (2012) IRF8 is a critical transcription factor for transforming microglia into a reactive phenotype. *Cell Rep* **1**:334–340.
- Matsumura Y, Yamashita T, Sasaki A, Nakata E, Kohno K, Masuda T, Tozaki-Saitoh H, Imai T, Kuraishi Y, Tsuda M, et al. (2016) A novel P2X4 receptor-selective antagonist produces anti-allodynic effect in a mouse model of herpetic pain. *Sci Rep* **6**:32461.
- McCarthy MM, Wright CL, and Schwarz JM (2009) New tricks by an old dogma: mechanisms of the organizational/activational hypothesis of steroid-mediated sexual differentiation of brain and behavior. *Horm Behav* **55**:655–665.
- Moriarty O, Tu Y, Sengar AS, Salter MW, Beggs S, and Walker SM (2019) Priming of adult incision response by early-life injury: neonatal microglial inhibition has persistent but sexually dimorphic effects in adult rats. *J Neurosci* **39**:3081–3093.
- Mouton PR, Long JM, Lei DL, Howard V, Jucker M, Calhoun ME, and Ingram DK (2002) Age and gender effects on microglia and astrocyte numbers in brains of mice. *Brain Res* **956**:30–35.
- Moy JK, Szabo-Pardi T, Tillu DV, Megat S, Pradhan G, Kume M, Asiedu MN, Burton MD, Dussor G, and Price TJ (2018) Temporal and sex differences in the role of BDNF/TrkB signaling in hyperalgesic priming in mice and rats. *Neurobiol Pain* **5**: 100024.
- North RA and Jarvis MF (2013) P2X receptors as drug targets. *Mol Pharmacol* **83**: 759–769.
- Oliveira-Fusaro MC, Gregory NS, Kolker SJ, Rasmussen L, Allen LH, and Sluka KA (2020) P2X4 receptors on muscle macrophages are required for development of hyperalgesia in an animal model of activity-induced muscle pain. *Mol Neurobiol* DOI: 10.1007/s12035-019-01852-x [published ahead of print].
- Paalme V, Rump A, Mädo K, Teras M, Truumees B, Aitai H, Ratas K, Bourge M, Chiang CS, Ghalali A, et al. (2019) Human peripheral blood eosinophils express high levels of the purinergic receptor P2X4. *Front Immunol* **10**:2074.
- Paige C, Maruthy GB, Mejia G, Dussor G, and Price T (2018) Spinal inhibition of P2XR or p38 signaling disrupts hyperalgesic priming in male, but not female, mice. *Neuroscience* **385**:133–142.
- Peng J, Gu N, Zhou L, B Eyo U, Murugan M, Gan WB, and Wu LJ (2016) Microglia and monocytes synergistically promote the transition from acute to chronic pain after nerve injury. *Nat Commun* **7**:12029.
- Rahimian R, Cordeau P Jr., and Kriz J (2019) Brain response to injuries: when microglia go sexist. *Neuroscience* **405**:14–23.
- Rosen SF, Ham B, Drouin S, Boachie N, Chabot-Dore AJ, Austin JS, Diatchenko L, and Mogil JS (2017) T-cell mediation of pregnancy analgesia affecting chronic pain in mice. *J Neurosci* **37**:9819–9827.
- Salter MW and Beggs S (2014) Sublime microglia: expanding roles for the guardians of the CNS. *Cell* **158**:15–24.
- Salter MW and Stevens B (2017) Microglia emerge as central players in brain disease. *Nat Med* **23**:1018–1027.
- Schopflocher D, Taenzer P, and Jovey R (2011) The prevalence of chronic pain in Canada. *Pain Res Manag* **16**:445–450.
- Schwarz JM, Sholar PW, and Bilbo SD (2012) Sex differences in microglial colonization of the developing rat brain. *J Neurochem* **120**:948–963.
- Sorge RE, LaCroix-Fralish ML, Tuttle AH, Sotocinal SG, Austin JS, Ritchie J, Chanda ML, Graham AC, Topham L, Beggs S, et al. (2011) Spinal cord Toll-like receptor 4 mediates inflammatory and neuropathic hypersensitivity in male but not female mice. *J Neurosci* **31**:15450–15454.
- Sorge RE, Mapplebeck JC, Rosen S, Beggs S, Taves S, Alexander JK, Martin LJ, Austin JS, Sotocinal SG, Chen D, et al. (2015) Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *Nat Neurosci* **18**: 1081–1083.
- Taves S, Berta T, Liu DL, Gan S, Chen G, Kim YH, Van de Ven T, Laufer S, and Ji RR (2016) Spinal inhibition of p38 MAP kinase reduces inflammatory and neuropathic pain in male but not female mice: sex-dependent microglial signaling in the spinal cord. *Brain Behav Immun* **55**:70–81.
- Teixeira JM, Dos Santos GG, Neves AF, Athie MCP, Bonet IJM, Nishijima CM, Farias FH, Figueiredo JG, Hernandez-Olmos V, Alshaibani S, et al. (2019) Diabetes-induced neuropathic mechanical hyperalgesia depends on P2X4 receptor activation in dorsal root ganglia. *Neuroscience* **398**:158–170.
- Thion MS, Low D, Silvin A, Chen J, Grisel P, Schulte-Schrepping J, Blecher R, Ulas T, Squarzon P, Hoeffel G, et al. (2018) Microbiome influences prenatal and adult microglia in a sex-specific manner. *Cell* **172**:500–516.e16.
- Trang T, Beggs S, Wan X, and Salter MW (2009) P2X4-receptor-mediated synthesis and release of brain-derived neurotrophic factor in microglia is dependent on calcium and p38-mitogen-activated protein kinase activation. *J Neurosci* **29**: 3518–3528.
- Tsuda M, Inoue K, and Salter MW (2005) Neuropathic pain and spinal microglia: a big problem from molecules in “small” glia. *Trends Neurosci* **28**:101–107.
- Tsuda M, Kuboyama K, Inoue T, Nagata K, Tozaki-Saitoh H, and Inoue K (2009) Behavioral phenotypes of mice lacking purinergic P2X4 receptors in acute and chronic pain assays. *Mol Pain* **5**:28.
- Tsuda M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, and Inoue K (2003) P2X4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* **424**:778–783.
- Ulmann L, Hirbec H, and Rassendren F (2010) P2X4 receptors mediate PGE2 release by tissue-resident macrophages and initiate inflammatory pain. *EMBO J* **29**: 2290–2300.
- Vacca V, Marinelli S, Pieroni L, Urbani A, Luvisetto S, and Pavone F (2014) Higher pain perception and lack of recovery from neuropathic pain in females: a behavioural, immunohistochemical, and proteomic investigation on sex-related differences in mice. *Pain* **155**:388–402.
- Verma R, Cronin CG, Hudobenko J, Venna VR, McCullough LD, and Liang BT (2017) Deletion of the P2X4 receptor is neuroprotective acutely, but induces a depressive phenotype during recovery from ischemic stroke. *Brain Behav Immun* **66**:302–312.
- Villa A, Della Torre S, and Maggi A (2019) Sexual differentiation of microglia. *Front Neuroendocrinol* **52**:156–164.
- Villa A, Gelosa P, Castiglioni L, Cimino M, Rizzi N, Pepe G, Lolli F, Marcello E, Sironi L, Vegeto E, et al. (2018) Sex-specific features of microglia from adult mice. *Cell Rep* **23**:3501–3511.
- Walker SM, Tochiki KK, and Fitzgerald M (2009) Hindpaw incision in early life increases the hyperalgesic response to repeat surgical injury: critical period and dependence on initial afferent activity. *Pain* **147**:99–106.
- Yang Y, Li H, Li TT, Luo H, Gu XY, Lü N, Ji RR, and Zhang YQ (2015) Delayed activation of spinal microglia contributes to the maintenance of bone cancer pain in female Wistar rats via P2X7 receptor and IL-18. *J Neurosci* **35**:7950–7963.
- Yu X, Liu H, Hamel KA, Morvan MG, Yu S, Leff J, Guan Z, Braz JM, and Basbaum AI (2020) Dorsal root ganglion macrophages contribute to both the initiation and persistence of neuropathic pain. *Nat Commun* **11**:264.

Address correspondence to: Michael W. Salter, The Hospital for Sick Children, 686 Bay St., Toronto, ON M5G 0A4, Canada. E-mail: michael.salter@sickkids.ca