

Glutamate Buffering Capacity and Blood-Brain Barrier Protection of Opioid Receptor Agonists Biphalin and Nociceptin

Saeideh Nozohouri,¹ Yong Zhang,¹ Thamer H. Albekairi, Bhuvaneshwar Vaidya, and Thomas J. Abbruscato

Department of Pharmaceutical Sciences, School of Pharmacy, Texas Tech University Health Sciences Center, Amarillo, Texas (S.N., Y.Z., T.H.A., B.V., T.J.A.); and Department of Pharmacology and Toxicology, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia (T.H.A.)

Received July 12, 2021; accepted September 29, 2021

ABSTRACT

Opioids play crucial roles in the regulation of many important brain functions including pain, memory, and neurogenesis. Activation of opioid receptors is reported to have neuroprotective effects after ischemic reperfusion injury. The objective of this study was to understand the role of biphalin and nociceptin, opioid receptor agonists, on blood-brain barrier (BBB) integrity during ischemic stroke. In this study, we aimed to measure the effect of biphalin and nociceptin on astrocytic glutamate uptake and on expression of excitatory amino acid transporter to study the indirect role of astrocytes on opioid receptor-mediated BBB protection during in vitro stroke conditions. We used mouse brain endothelial cells (bEnd.3) and primary astrocytes as an in vitro BBB model. Restrictive BBB properties were evaluated by measuring [¹⁴C] sucrose paracellular permeability and the redistribution of the tight junction proteins. The protective effect of biphalin and nociceptin on BBB integrity was assessed after exposing cells to oxygen glucose deprivation (OGD) and glutamate. It was observed that combined stress (2 mM glutamate and 2 hours of OGD) significantly reduced glutamate uptake by astrocytes; however, biphalin and nociceptin treatment

increased glutamate uptake in primary astrocytes. This suggests a role of increased astrocytic buffering capacity in opioid-mediated protection of the BBB during ischemic stroke. It was also found that the combined stress significantly increased [¹⁴C] sucrose paracellular permeability in an in vitro BBB model. Biphalin and nociceptin treatment attenuated the effect of the combined stress, which was reversed by the opioid receptor antagonists, suggesting the role of opioid receptors in biphalin and nociceptin's BBB modulatory activity.

SIGNIFICANT STATEMENT

There is an unmet need for discovering new efficacious therapeutic agents to offset the deleterious effects of ischemic stroke. Given the confirmed roles of opioid receptors in the regulation of central nervous system functions, opioid receptor agonists have been studied as potential neuroprotective options in ischemic conditions. This study adds to the knowledge about the cerebrovascular protective effects of opioid receptor agonists and provides insight about the mechanism of action of these agents.

Introduction

Ischemic stroke, one of the major causes of disability and death worldwide, comprises a series of complex pathogenic processes such as reactive oxygen species production, excitotoxicity, inflammation, blood-brain barrier (BBB) disruption, and eventually neuron death. Amelioration of these deleterious processes will improve postischemic outcomes in patients and ultimately neuronal recovery. However, only one Food and Drug Administration-approved therapeutic drug exists for ischemic stroke, tissue plasminogen activator (t-PA), which has limitations regarding side effects and patient eligibility. Therefore, there is an unmet need for discovering new

efficacious therapeutic agents to offset the deleterious effects of ischemic stroke (Nozohouri et al., 2020).

After ischemic injury, the major excitatory neurotransmitter, glutamate, is increased through excess release from activated presynaptic neurons and malfunctioned clearance by the astrocytic glutamate transporters. Excess extracellular glutamate is excitotoxic to neurons. It has also been reported to increase BBB permeability in cultured brain endothelial cells through activation of *N*-methyl-D-aspartate (NMDA) receptors (Xhima et al., 2016). The BBB functions as a highly specialized interface, which comprises cellular and vascular components. The selective nature of the BBB allows the maintenance of a unique extracellular milieu within the brain that is essential for normal central nervous system (CNS) function. Astrocytes play an important role in BBB integrity through regulating cerebral endothelial cell function, buffering extracellular neurotransmitter concentrations, and modulating CNS ionic balances and fluid accumulation. Due to its toxicity at higher

This work was funded from resources provided by the Department of Pharmaceutical Sciences of the Jerry H. Hodge School of Pharmacy.

¹S.N. and Y.Z. contributed equally to the study.

dx.doi.org/10.1124/jpet.121.000831.

ABBREVIATIONS: BBB, blood-brain barrier; BMEC, brain microvascular endothelial cell; CNS, central nervous system; EAAT, excitatory amino acid transporter; GLAST, glutamate-aspartate transporter; mAb, monoclonal antibody; NMDA, *N*-methyl-D-aspartate; NOP, nociceptin opioid receptor; OGD, oxygen glucose deprivation; OR, opioid receptor; t-PA, tissue plasminogen activator.

concentrations, extracellular glutamate is required to be rapidly removed by a set of excitatory amino acid transporters (EAATs) primarily expressing in astrocytes including glutamate-aspartate transporter (GLAST, also known as EAAT1) and glutamate transporter-1 (GLT-1, also known as EAAT2). Importantly, increased glial EAAT1 (but not EAAT2) has been observed in patients with traumatic brain injury, which reflects a neuroprotective potential of EAAT1 (Choi et al., 2015). The neuroprotective role of EAAT1 has been also reported in postischemic brain damage in EAAT1-deficient animals and in patients with stroke (Yamashita et al., 2006; Chen et al., 2015). These studies suggest a superior role of astrocytic EAAT1 over EAAT2 in maintaining the excitatory neurotransmitters hemostasis after brain injuries. Therefore, developing novel agents to regulate astrocytic EAAT1 and glutamate buffering capacity could be a promising neurovascular protective strategy in the management of ischemic stroke. Targeting glutamate clearance may provide an advantage over glutamate receptor antagonists, which have been found to decrease neurodegeneration in some nondamaged areas, which produced failed results in clinical trials (Ikonomidou and Turski, 2002).

Opioids play crucial roles in the regulation of CNS functions and hemodynamics through activation of the opioid receptors (ORs) distributing widely throughout the brain. ORs comprise four subtypes, i.e., the classic mu-OR, delta-OR, kappa-OR, and the newer nociceptin opioid peptide receptor (NOP). The classic ORs have gained extensive interest of researchers for their role as neuroprotective targets and have been reported to demonstrate potential neuroprotective activities in ischemic stroke by several overlapping mechanisms (Vaidya et al., 2018). There are numerous studies regarding neuroprotective effects of both opioid agonists and antagonists. Liao et al. have shown that naloxone decreases brain infarction, chemokine expression and neutrophil accumulation through blocking mu opioid receptors after ischemic injury (Liao et al., 2003). Although earlier preclinical reports have shown neuroprotective activity of OR antagonists, clinical trials failed to verify those findings due to the adverse effects of high doses of antagonists on neuronal functions (Vaidya et al., 2018). Our laboratory has published several studies reporting the neuroprotective effects of a nonselective OR agonist buprenorphine, which is associated with reduced reactive oxygen species production, brain edema, infarct size, and neuron death, tested in ischemic stroke models (Cowell and Lee, 2016; Yang et al., 2015; Islam et al., 2016). The structure of buprenorphine is a combination of two tetrapeptides derived from enkephalin pharmacophore, which have high affinity to delta-OR and mu-OR and low affinity to kappa-OR but no affinity to NOP (Yang et al., 2015). Nociceptin is a natural ligand of NOP with no affinity to the classic ORs. Furthermore, a recent study reports that activation of NOP serves as a regulator of EAAT1 and glutamate uptake in developing astrocytes (Meyer et al., 2017). This study raises the questions 1) if a potent OR activator, such as buprenorphine, has similar regulating capacity of astrocytic EAAT1 and glutamate uptake and 2) if the glutamate buffering activity of OR agonists ameliorate ischemia/glutamate-induced BBB damage.

In the present study, we examine glutamate buffering activity of buprenorphine and nociceptin under ischemic conditions with primary astrocytes and their protective potential against

ischemia/glutamate-induced paracellular permeability with an endothelial cells/astrocytes coculture BBB model.

Materials and Methods

Cell Culture and bEnd3/Astrocyte Coculture. bEnd3 cells (passages 27–29) were cultured in Dulbecco's modified Eagle's medium (Sigma, St. Louis, MO) supplemented by FBS (Atlanta Biologicals, Minneapolis, MN) and nonessential amino acid and penicillin/streptomycin (Sigma, St. Louis, MO). Cells were maintained in a humidified incubator at 37°C and with 5% CO₂. Mouse primary astrocytes were obtained from the cerebral cortices of one day old CD-1 mouse pups (Charles Rivers Laboratory) according to the method explained previously by Du et al. (2010). After isolating the brain, cortices were removed and placed in Hanks' balanced salt solution supplemented with gentamycin (10 µg/ml). Then cortices were digested with 0.25% trypsin for 10–15 minutes at 37°C followed by neutralizing with FBS containing Dulbecco's modified Eagle's medium. The cells were then seeded into a cell culture flask, and the medium was refreshed every 3 days until reaching confluency.

For bEnd3 and astrocyte co-culture, the Transwell inserts (0.4-µm pore size, 12-well; Corning, Lowell, MA) were inverted, and astrocytes with a density of 150,000 cells per filter were seeded onto the basolateral side of the filter membrane and allowed to adhere. After 4 hours, the Transwells were inverted back, and astrocytes were allowed to grow for 2 more days in astrocyte medium. Then, bEnd3 cells with a density of 50,000 cells per filter were added onto the apical side of the Transwell filter. The co-culture of primary astrocytes and bEnd3 cells was grown for 8 days. Media were changed every other day (Li et al., 2010). Barrier integrity was assessed by measuring transendothelial electrical resistance by EVOM resistance meter (World Precision Instruments, Sarasota, FL) using the STX-2 electrodes.

Oxygen Glucose Deprivation. To simulate the stroke condition in vitro, cells were exposed to an oxygen glucose deprivation (OGD) condition following an established protocol (Yang et al., 2015). For this purpose, cell culture medium was removed, and cells were rinsed with Dulbecco's phosphate buffered saline solution and glucose-free Earle's balanced salt solution (in mM 140 NaCl, 0.83 MgSO₄, 5.36 KCl, 1.02 NaH₂PO₄, 1.18 CaCl₂, 26.19 NaHCO₃) bubbled with 95% N₂/5% CO₂ was added to the cells to create an aglycemic condition. Then cells were transferred to a custom-made hypoxic chamber (Coy Laboratories, Grasslake, MI) with 95% N₂ and 5% CO₂ at 37°C to induce hypoxia (1% oxygen). Cells were exposed to the OGD condition for 2 hours.

Glutamate ± OGD ± Buprenorphine and Nociceptin Treatment on the In Vitro BBB Models. The permeability experiments were performed on days 8–10 after mono- or co-culture establishment. On the day of experiment, the in vitro BBB model was incubated with 1 µM naltrexone in HEPES buffer (in mM 120 NaCl, 1 CaCl₂, 25 HEPES, 1 KH₂PO₄, 2 KCl, 1 MgSO₄, 10 D-glucose) in both compartments for 30 minutes; then the medium was changed to 1 µM naltrexone and 0.1 µM buprenorphine in HEPES. After 30 minutes, naltrexone, buprenorphine, and 2 mM glutamate were added to the cells in Earle's balanced salt solution, which simulates the glutamate being released from neurons. Then the cells were transferred to the OGD chamber. The same treatment strategy was used to evaluate the effect of 1 µM nociceptin and its inhibitor. For the co-culture of bEnd3 cells and astrocytes, the treatment was only added to the basolateral side of the Transwell insert to study the role of astrocytes in the BBB protection.

Permeability Measurement. The paracellular permeability was assessed by measuring the diffusion profile of [¹⁴C] sucrose (Nozohouri et al., 2021). Briefly, after 2 hours of OGD, the medium was removed from both compartments; 1.5 ml HEPES was added to the basolateral compartment, and 0.5 ml HEPES containing 0.1 µCi/ml of [¹⁴C] sucrose was added to apical compartment for evaluation of BBB permeability. Samples (100 µl) were removed from the basolateral chamber at (5, 15, 30, 60, and 120 minutes) and replaced with fresh HEPES. The radioactivity of collected samples were evaluated using a




Fig. 1. Concentration- and time-dependent effect of biphalin and nociceptin in glutamate uptake and EAAT1 expression in primary cortical astrocytes. Increasing concentrations of biphalin (A) and nociceptin (B) increase [³H] glutamate uptake in primary cortical astrocytes in normoxic condition. (C) This is due to the increased expression of glutamate transporter, EAAT1, in these cells. (D) Biphalin and nociceptin increase the expression of EAAT1 in a time-dependent manner, and different exposure time results in overexpression of EAAT1 accordingly. (E) The overexpression of EAAT1 is blocked by the antagonists of NOP and classic opioid receptors, which suggests that the role of these compounds in increasing the expression of glutamate transporter is mediated through opioid receptors. Data represented as means ± S.D. of 3 to 4 independent replicates. ***P* < 0.01; *****P* < 0.0001; ns, not significant; Bi, biphalin; NOC, nociceptin; NOC-I, nociceptin inhibitor; NTX, naltrexone.

liquid scintillation counter (Beckman Coulter, Brea, CA). Then the permeability coefficient was calculated using the following formula:

$$PC = \frac{dQ}{dt} \times \frac{1}{C_0 \times A}$$

where *dQ/dt* is the diffusion rate of [¹⁴C] sucrose across the membrane, *A* is the area of Transwell insert, and *C*₀ is the initial concentration of the paracellular marker added in the donor compartment.

Western Blot Analysis. Astrocytes and bEnd3 cultures were lysed using radioimmunoprecipitation assay buffer (RIPA) buffer supplemented with proteinase and phosphatase inhibitor cocktails and then collected. Protein concentration in each sample was measured using BCA protein assay. After solubilizing samples using Laemmli buffer, equal quantities of proteins (20 μg per lane) were loaded and separated by SDS-PAGE in 10% acrylamide and then transferred to a polyvinylidene fluoride (PVDF) membrane for 2 hours at 100 V. Then the membrane was incubated in blocking solution containing 5% non-fat dry milk to block nonspecific binding. Then the membranes were

immunoblotted with the appropriate primary antibodies overnight at 4°C with the following dilutions: anti-EAAT1/GLAST (1:1000) rabbit monoclonal antibody (mAb) (Cell Signaling Technology, Danvers, MA), anti-claudin-5 (1:500) mouse mAb (Invitrogen, Waltham, MA), and anti-actin (1:10000) mouse mAb (Sigma, St. Louis, MO). After rinsing, the membrane was incubated with the appropriate anti-rabbit or anti-mouse secondary antibody (Cell Signaling Technology) for 2 hours, and the bands were visualized with enhanced chemiluminescence. Beta-actin immunoreactivity was used as the loading control in each sample and was used to normalize the corresponding data. Data were analyzed using NIH ImageJ software and are reported as ratios of the control group.

[³H] Glutamate Uptake. For this sets of experiments, nociceptin (1 μM), biphalin, and selective opioid receptor agonists (0.1 μM) with or without the nonselective opioid receptor antagonist naltrexone (1 μM) and nociceptin receptor antagonist BAN ORL 24 (0.1 μM) were added to the primary astrocyte culture grown on six-well plates. After overnight incubation, the selected groups were transferred to the OGD chamber following the above-mentioned method, and after 2 hours of OGD, [³H] glutamate uptake assay was performed on all




Fig. 2. Effect of biphalin and nociceptin pretreatment in [^3H] glutamate uptake and EAAT1 expression in primary cortical astrocytes after exposure to 2 hours of OGD. Two hours of OGD increase the capacity of astrocytes in glutamate uptake, and pretreatment with biphalin (A) or nociceptin (C) further increase this capacity, which is inhibited by the opioid receptor antagonists. This effect is mediated by significant upregulation of EAAT1 in primary cortical astrocytes [(B), (D)]. Data represented as means \pm S.D. of 3 to 4 independent replicates. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. Bi, biphalin; NOC, nociceptin; NOC-I, nociceptin inhibitor; NTX, naltrexone.

groups. In one set of experiments, the astrocytes were incubated with 10 μM of EAAT1 inhibitor, UCPH 101 (Abcam, Cambridge, UK), for 15 minutes after the OGD and before performing the glutamate uptake assay. First, a 20 mM stock solution of UCPH 101 was prepared in dimethyl sulfoxide, and the final concentration of 10 μM was prepared in HEPES and then was added to cells (Liang et al., 2014). The uptake was measured by incubating the cells with [^3H] glutamate (0.1 $\mu\text{Ci/ml}$) for 7 minutes. An incubation time of 7 minutes was selected based on the results of the time-dependent uptake study performed in our laboratory (data not shown). The uptake study was conducted by incubating the cells grown on six-well plates with HEPES containing 25 μM glutamate, which mimics the physiologic concentration of glutamate. The uptake was terminated by removing radioactive HEPES buffer and rinsing the cells with ice-cold HEPES. Cells were then lysed using 1% Triton X-100 (ARCOS). Radioactivity was then measured using a liquid scintillation machine (Beckman Coulter). Protein estimation was performed on lysates using a BCA protein estimation kit (Pierce BCA Protein Assay Kit) following the manufacturer's protocol. The data were normalized by protein estimation and presented as the percentage change compared with the control group.

Immunocytochemistry. The bEnd3 cells were seeded into four-well chamber slides (Laboratory-Tek II Chamber slides). After reaching confluency, the cells were exposed to OGD and glutamate with or without treatment with biphalin and nociceptin. Afterward, the cells were fixed with 4% paraformaldehyde, washed, and permeabilized with 0.1% Triton X-100 for 5 minutes. Cells were incubated with blocking solution containing 1% bovine serum albumin and 2% goat serum for 1 hour at room temperature followed by incubation with primary antibody (1:100) in the same blocking solution overnight at 4°C. Next day, the cells were washed with 0.1% bovine serum albumin solution in PBS and were stained with fluorescence-tagged corresponding secondary antibody at room temperature for 2 hours followed by washing three times with PBS. Slides were mounted with 4',6-diamidino-2-phenylindole (DAPI) containing Prolong Gold antifade mounting medium (Invitrogen). Fluorescent Images were obtained using a multiphoton microscope (A1R; Nikon, Melville, NY) in the confocal mode, using a 60 \times objective.

Statistical Analysis. All the data are presented as means \pm S.D. of at least three independent replicates. Unpaired Student's t test was used to compare two groups, whereas one-way ANOVA




Fig. 3. Effect of biphalin and nociceptin on glutamate uptake in astrocytes exposed to OGD and glutamate combined stress condition. (A) Exposure of astrocytes to 2 mM of glutamate along with 2 hours of OGD significantly decreased the capacity of astrocytes in uptake of glutamate. (B) Pretreatment of cortical astrocytes with 0.1 μM of biphalin or 1 μM of nociceptin significantly increased the uptake of [³H] glutamate in damaged astrocytes exposed to 2 hours of OGD and 2 mM glutamate compared with untreated groups. (C) This effect is mediated through upregulation of EAAT1 in primary cortical astrocytes. (D) The effect of EAAT1 in the transport of extracellular glutamate in biphalin- and nociceptin-treated astrocytes was significantly blocked by the selective inhibitor of EAAT1, UCPH 101 (10 μM), reflecting the major contribution of this transporter in the clearance of glutamate. Data represented as means ± S.D. of 3 to 4 independent replicates. ***P* < 0.01; *****P* < 0.0001. Bi, biphalin; Glut, glutamate; NOC, nociceptin.

followed by Tukey’s post hoc multiple comparison test was used to compare more than two groups (Prism, version 8.0; GraphPad Software Inc., San Diego, CA). *P* values <0.05 were considered statistically significant.

Results

Biphalin and Nociceptin Increase Glutamate Uptake and Upregulate Glutamate Transporter EAAT1 Expression in Primary Astrocytes in a Concentration- and Time-Dependent Manner. To investigate the effect of biphalin and nociceptin on the glutamate buffering capacity of astrocytes, increasing concentrations of biphalin and nociceptin were added to mouse primary cortical astrocytes, and [³H] glutamate uptake was measured in these cells. As shown in Fig. 1, A and B, biphalin and nociceptin increase the uptake of glutamate by astrocyte in a concentration-dependent manner in

normoxic condition. We suggest that biphalin and nociceptin mediate glutamate uptake through upregulation of glutamate transporter, EAAT1/GLAST, in astrocytes. As depicted in Fig. 1C, increasing concentrations of biphalin and nociceptin upregulate the expression of EAAT1 in primary astrocytes and hence increase the glutamate buffering capacity of these cells. With these experiments, one concentration of biphalin (0.1 μM) and nociceptin (1μM) was selected, and the cells were incubated with that concentration for various time points. It was observed that biphalin and nociceptin increase the expression of EAAT1 in a time-dependent manner as well (Fig. 1D). To further evaluate the mechanism involved in this upregulation, we preincubated the cells with antagonists of opioid receptors, naltrexone for biphalin and BAN ORL 24 (a nociceptin inhibitor) for nociceptin. The results indicated that the overexpression of EAAT1 is inhibited by the antagonists of NOP and classic opioid receptors, suggesting that the role of nociceptin and biphalin in




Fig. 4. Comparison between the effect of biphalin and selective opioid receptor agonists in inducing the glutamate uptake by primary astrocytes. (A) With the same concentration (0.1 μ M), biphalin increases the uptake of [³H] glutamate in astrocytes more than selective kappa (U-50488), delta (DPDPE), and mu (DAMGO) opioid receptor agonists confirming better protective effect of biphalin compared with selective agonists. (B) Also, biphalin increases the expression of EAAT1 more than selective opioid receptor agonists. Data represented as means \pm S.D. of 3 to 4 independent replicates. ** P < 0.01; *** P < 0.001. Bi, biphalin.

increasing the expression of glutamate transporter is mediated through classic and nociceptin opioid receptors (Fig. 1E). Based on the above-mentioned findings, we selected 12 hours of treatment with 0.1 μ M biphalin and 1 μ M nociceptin for future experiments of this study.

Biphalin and Nociceptin Pretreatment Increases the [³H] Glutamate Uptake in Primary Cortical Astrocytes Challenged with 2 Hours of OGD by Increasing the Expression of EAAT1. After confirming the effect of biphalin and nociceptin in increasing the expression of EAAT1 and glutamate uptake by astrocytes in normoxic condition, we studied their effect on OGD-exposed cells as well to mimic ischemic stroke condition. As depicted in Fig. 2, A and C, 2 hours of OGD increases the capacity of astrocytes in clearing the excess extracellular glutamate by upregulating the level of EAAT1 in these cells; however, pretreatment with 0.1 μ M biphalin or 1 μ M nociceptin further increases glutamate uptake by primary astrocytes, which is mediated by significant upregulation of glutamate transporter, EAAT1 (Fig. 2, B and D). Also, like normoxic condition, this protective effect is reversed by corresponding antagonists of opioid or NOP receptors during the OGD condition.

Biphalin and Nociceptin Improve Glutamate Uptake in Astrocytes Exposed to OGD and Glutamate Combined Stress Condition. Two hours of OGD alone did not compromise the capacity of astrocytes in glutamate uptake

and in fact increased the uptake; however, exposure of astrocytes to 2 mM of glutamate along with 2 hours of OGD significantly decreased the capacity of astrocytes in uptake of glutamate by 45% compared with the control group (P < 0.0001) (Fig. 3A). Interestingly, pretreatment of cortical astrocytes with 0.1 μ M of biphalin or 1 μ M of nociceptin significantly increased the uptake of [³H] glutamate in damaged astrocytes exposed to 2 hours of OGD and 2 mM glutamate compared with untreated groups (Fig. 3B). Moreover, compatible with the uptake study, in these groups of astrocytes the expression of EAAT1 was improved significantly by biphalin and nociceptin pretreatment compared with the untreated group (3C). In another set of experiments, primary astrocytes were preincubated with 10 μ M EAAT1 inhibitor, UCPH 101, prior to glutamate uptake to evaluate the role of EAAT1 in the clearance of extracellular glutamate. As depicted in Fig. 3D, 15-minute exposure of astrocytes to UCPH 101 significantly decreased [³H] glutamate uptake in the biphalin- and nociceptin-treated cells by blocking the transport of glutamate via EAAT1.

Nonselective OR Agonist, Biphalin, Shows Superior Glutamate Buffering Activity Compared with Subtype-Selective OR Agonists. To further investigate the potency of biphalin, we compared the effect of it with different selective opioid receptor agonists. As shown in Fig. 4, with the same concentration (0.1 μ M), biphalin increases the uptake of [³H] glutamate in astrocytes 44%, 52%, and 45% more than selective kappa (U-50488), delta ([D-Pen2,5]Enkephalin, [D-Pen2,D-Pen5]Enkephalin, DPDPE), and mu ([D-Ala2, N-MePhe4, Gly-ol]-enkephalin, DAMGO) opioid receptor agonists, respectively, confirming a better excitatory amino acid protective effect of biphalin compared with selective agonists. Similarly, biphalin increases the expression of EAAT1 more than selective opioid receptor agonists.

Biphalin and Nociceptin Stabilize BBB Integrity in an In Vitro Co-culture Model Exposed to OGD and Glutamate Combined Stress Condition. To investigate the possible role of astrocytes in increasing the barrier properties of the BBB and buffering the ions and transmitters, we established and used in vitro coculture model of primary astrocytes and bEnd3 cells. We exposed this model to the combined stressor of 2 mM glutamate and 2 hours of OGD, which was intended to compromise the BBB and increase the paracellular permeability. We found that the percentage increase in the paracellular permeability in the monoculture of bEnd3 cells was 54% (0.00059 cm/min to 0.00091 cm/min), whereas in the coculture model this change was 32% (0.0005 cm/min to 0.00066 cm/min) (Fig. 5A). These data suggest that astrocytes have an important role in influencing the BBB integrity during in vitro stroke conditions and with elevated levels of glutamate. To potentiate the protective role of astrocytes in maintaining the BBB integrity, the co-culture of bEnd3 cells and astrocytes was pretreated with 0.1 μ M biphalin or 1 μ M nociceptin in the basolateral side of the model, where astrocytes reside. It was found that biphalin pretreatment significantly decreased the paracellular permeability compared with the group exposed to 2 hours of OGD and 2 mM glutamate (0.00058 cm/min versus 0.00041 cm/min; P < 0.0001). To confirm that this effect is mediated through opioid receptors we further pretreated the coculture with 1 μ M naltrexone prior to the experiment. Interestingly, we found that naltrexone significantly negates the effect of biphalin, suggesting the role of




Fig. 5. (A) The percentage increase in the paracellular permeability in response to glutamate and OGD. It was found that the monoculture of bEnd3 cells was almost 20% more permeable than the coculture model, which is much resistant to combined stress condition indicating the role of astrocytes in BBB integrity and protection. (B) Role of opioid receptors in the leakiness of the BBB in coculture model. Biphalin significantly stabilized the BBB and attenuated the increase in paracellular permeability due to the combined stress condition. Naltrexone pretreatment negated this effect indicating that the action is mediated through opioid receptors. Nociceptin treatment also showed similar effect on the BBB permeability confirming the involvement of opioid receptors in the integrity of BBB. Data represented as means \pm S.D. of 3 to 4 independent replicates. $**P < 0.01$; $***P < 0.001$; ns, not significant; NOC, nociceptin; PC, permeability coefficient.

opioid receptors in the mechanism of biphalin protection (0.00041 cm/min versus 0.00053 cm/min; $P < 0.01$) (Fig. 5B). Similar to biphalin, nociceptin treatment decreased the permeability of paracellular marker and improved the BBB integrity compared with the combined stress-exposed group (0.00058 cm/min versus 0.00036 cm/min; $P < 0.0001$), and this effect was partially blocked by the nociceptin receptor’s inhibitor, BAN ORL 24.

Biphalin and Nociceptin Attenuate the Increased BBB Permeability and Improve the Expression of Tight Junction Protein in bEnd3 Monolayers After Combined Stress Condition (2 Hours of OGD + 2 mM Glutamate). To evaluate the protective effect of biphalin and nociceptin directly on bEnd3 cells, monolayers of bEnd3 cells were exposed to either 2 mM glutamate at normoxic condition or 2 mM of glutamate and 2 hours of OGD. As depicted in Fig. 6A, 2 hours of OGD alone did not significantly increase the paracellular permeability of [¹⁴C] sucrose compared with the normoxic group; however, addition of 2 mM glutamate increased the BBB permeability significantly (0.00074 cm/min versus 0.00051 cm/min; $P < 0.01$). The most significant level

compared with normoxia was obtained when both 2 hours of OGD and 2 mM glutamate were combined (0.00091 cm/min versus 0.00051 cm/min; $P < 0.001$). Pretreatment of bEnd3 monolayers with 0.1 μ M biphalin or 1 μ M nociceptin significantly attenuated glutamate-mediated increase in BBB permeability in the OGD condition (0.00077 cm/min versus 0.00052 cm/min; $P < 0.05$) (Fig. 6B).

The most prominent structural and molecular components of the BBB are tight junctional proteins between the brain microvascular endothelial cells (BMECs). To provide further mechanistic insight on the protective effect of biphalin and nociceptin on barrier function, in another set of experiments we evaluated the expression of tight junction protein claudin-5 in bEnd3 cells after exposing the cells to 2 hours of OGD and 2 mM of glutamate using Western blotting and immunofluorescence staining. It was observed that pretreatment of bEnd3 cells with biphalin or nociceptin significantly improved the expression of claudin-5 compared with the untreated group after exposure to the combined stress condition. Representative images of claudin-5 appear in Fig. 6C. Significant disruptions were seen in the distribution and expression of claudin-5




Fig. 6. Effect of biphalin and nociceptin on the BBB permeability and the expression of tight junction proteins in bEnd3 monolayers after combined stress condition (2 hours of OGD + 2 mM glutamate). (A) Two hours of OGD did not cause significant alteration in [¹⁴C] sucrose paracellular permeability, but the addition of 2 mM glutamate increased the permeability of paracellular marker in normoxic condition. Combination of 2 mM glutamate and 2 hours of OGD significantly increased the permeability of radiolabeled marker. (B) Biphalin or nociceptin pretreatment significantly attenuated the effect of 2 mM glutamate and 2 hours of OGD by decreasing the paracellular permeability of [¹⁴C] sucrose compared with the untreated group, suggesting the involvement of glutamate in the protective mechanism of action of these opioid agonists. (C) Representative immunofluorescence images showing protein expression and localization of claudin-5 in bEnd3 cells after exposure to combined stress condition and treatment with biphalin or nociceptin. (D) Western blot studies also confirmed that biphalin and nociceptin treatment increase the expression of claudin-5 in cells exposed to combined stress conditions. Data represented as means ± S.D. of 3 to 4 independent replicates. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; ns, not significant; PC, permeability coefficient.

after combined stress, which is attenuated with biphalin or nociceptin pretreatment since they resulted in more intense staining of claudin-5 at cell-to-cell contact points.

Discussion

The main challenge in developing new therapeutics for the management of ischemic stroke is the complex pathophysiology of this disorder along with the incomplete understanding of contributing cellular and molecular mechanisms (Nozohouri et al., 2020). It is well known that paracellular permeability of compounds across BBB increases significantly after ischemic injury, which is due to the opening of the tight junction proteins. Endothelial cells of the cerebral microvasculature function to protect neurons and glial cells from harmful insults. Among tight junction proteins, claudins, a family of transmembrane proteins, play a crucial role in sealing the gaps between cells and limiting paracellular passage of ions. Among various

forms of claudin, claudin-5 is mainly expressed in cerebral microvascular endothelial cells.

Since ischemic stroke initiates multiple pathways resulting in neuronal cell death, successful neuroprotection needs stimulation of several pathways. Therefore, activation of opioid receptor subtypes to simultaneously target various neuroprotective pathways is a rational therapeutic strategy. The neuroprotective role of opioid receptor agonists in ischemic stroke has been widely reported by different research groups. Opioid receptor activation attenuates several pathophysiological factors such as excitotoxicity, inflammation, and anoxic depolarization. Moreover, these receptors prevent BBB disruption, regulate astrocytic alterations, and offset potential adverse effects of t-PA. It has also been reported that nonselective opioid receptor agonists improve stroke outcomes better than selective agonists (Popielek-Barczyk et al., 2017; Yang et al., 2015).

However, opioid agonists have also contributed to ischemic stroke pathogenesis. Along with common side effects of abuse

and addiction related to opioids, long-term pain management using prescribed opioids presents neurovascular complications that contribute to ischemic stroke (Voon et al., 2018; Fallon et al., 2018). This class of drugs, such as morphine, acts primarily by activating mu-ORs and result in mitochondrial dysfunction, oxidative stress, and thus BBB impairment (Feng et al., 2012; Woller et al., 2012; Hamzei Moqaddam et al., 2009). Therefore, opioid antagonists could also provide an appealing therapeutic strategy for the management of ischemic stroke (Anttila et al., 2018; Chen et al., 2001; Liao et al., 2003).

Biphalin is an example of a nonselective, potent opioid receptor agonist that has high affinity to mu and delta opioid receptors. Nociceptin, on the other hand, is an endogenous peptide that binds to the G-protein-coupled receptor, opioid receptor like-1, which is another member of the opioid receptor family. Despite the structural similarity of nociceptin to dynorphin A, it does not have binding affinity to the classic opioid receptors. In this study, we evaluated the cerebrovascular protective properties of biphalin and nociceptin.

This study focused on understanding how BMECs respond to *in vitro* ischemic conditions after treatment with opioid receptor agonists. To accomplish this goal, we used *in vitro* mono- and co-culture models, which are composed of bEnd3 cells cultured with astrocytes positioned opposed on a polyester Transwell inserts. To simulate ischemia *in vitro*, we optimized the timing of BBB disruption, which was exposure of cells to 2 hours of hypoxic condition with the addition of 2 mM glutamate, as glutamate level was reported to increase up to 10 mM during ischemic injury (András et al., 2007). BMEC monolayers and co-culture of bEnd3 cells and astrocytes exposed to OGD and glutamate showed a significant increase in [¹⁴C] sucrose permeability compared with control groups. These data are consistent with other published findings claiming increased permeability in BMECs treated with hypoxia for various durations (Mark and Davis, 2002).

One important finding was that biphalin and nociceptin pretreatment prevented the increase in BBB opening through opioid receptors since the protective effect was negated by their corresponding inhibitors. The protective effect of biphalin and nociceptin on the BBB integrity is proposed to be mediated through glutamate. Therefore, it is crucial to understand how the increased level of glutamate during ischemia-reoxygenation disrupts the BBB mechanistically. The effect of glutamate-induced alterations of the tight junctional proteins in cultured brain microvascular endothelial cells occurred after exposure to glutamate and resulted in cellular redistribution of these proteins followed by a decrease in their expression level. Interestingly, biphalin and nociceptin treatments were effective in improving the expression of claudin-5 hence attenuating the paracellular permeability of [¹⁴C] sucrose. However, pretreatment with biphalin and nociceptin did not improve the expression level and distribution of zonula occludens-1 (ZO-1) and occludin in damaged cells (data not shown).

Since stroke is both a vascular and neuronal disorder, it is advantageous to activate opioid receptors in different sites of neurovascular unit, mainly neurons and astrocytes, to exhibit the maximum beneficial effects. Astrocytes have been reported to express opioid receptors, and the expression level of these receptors can be modulated by the exposure to different stimuli such as glutamate (Thorlin et al., 1997). Astrocytes provide various housekeeping functions including formation of the BBB, modulation of cellular communications, and protection

against oxidative stress. Moreover, astrocytes play crucial roles in the removal of extracellular glutamate and potassium during synaptic plasticity (Yao et al., 2014). It is well known that in the ischemic brain, the most common neurotoxic transmitter is glutamate and that an excess amount of it results in excitotoxicity and neuronal damage. Various approaches such as NMDA receptor antagonists have been studied in clinical settings for the prevention of glutamate-induced excitotoxicity. However, application of NMDA antagonists for cerebral ischemia has failed in these trials since these antagonists hinder neuronal survival by blocking the synaptic transmission (Ikonomidou and Turski, 2002; Wang and Shuaib, 2005). Another therapeutic approach to inhibit glutamate-mediated excitotoxicity is to decrease extracellular glutamate concentration through increasing the uptake of glutamate by EAATs (Simantov et al., 1999; Williams et al., 2005). The glutamate-aspartate transporter, named EAAT1 in humans, is one of the main transporters during brain development and plays an important role in protecting brain against glutamate-induced cytotoxicity. These transporters are predominantly expressed on astrocytes, and their expression can be regulated pharmacologically. The upregulation of EAAT on astrocytes has been shown to improve neurologic score and decrease infarct volume in experimental stroke animal models (Verma et al., 2010; Beschorner et al., 2007). In our study we also focused on evaluating the protective effect of biphalin and nociceptin on the glutamate clearance after glutamate-induced injury in primary cortical astrocytes. To perform these experiments, we measured the uptake of [³H] glutamate in primary cortical astrocytes. First, we validated the published method reported in primary rat cortical astrocytes in mice cortices using nociceptin treatment as a positive control since nociceptin regulates the levels of the glutamate/aspartate transporters GLAST/EAAT1 in both human and rodent brain astrocytes (Meyer et al., 2017). GLAST/EAAT1 upregulation by nociceptin is mediated by nociceptin opioid receptors and the downstream participation of a complex signaling cascade that involves the interaction of several kinase systems.

Interestingly, biphalin acted similarly as nociceptin and increased the glutamate uptake in primary cortical astrocytes. In addition, biphalin effect on glutamate uptake was observed at 0.1 μM concentration, which is the same effective concentration that stabilized the BBB permeability compromised by OGD and glutamate. Exposure of primary cortical astrocytes to 2 hours of OGD increased the glutamate uptake; however, combined stress decreased the capacity of astrocytes in [³H] glutamate uptake significantly. Pretreatment with 0.1 μM biphalin and 1 μM nociceptin increased the protective effect of astrocytes, and their effect was negated by opioid receptor and nociceptin receptor antagonists. Interestingly, biphalin was more effective than nociceptin and selective kappa (U-50488), delta (DPDPE), and mu (DAMGO) opioid receptor agonists in increasing the capacity of astrocytes for glutamate uptake and nociceptin showed similar effect to the selective OR agonists. The mechanism behind this effect was evaluated by Western blotting to investigate the expression level of glutamate transporters.

As expected, pretreatment with biphalin and nociceptin increased the expression of astrocytic EAAT1. Additionally, the effect of upregulated EAAT1 was significantly blocked by the selective inhibitor of this transporter, which further confirms that the effects of biphalin and nociceptin in glutamate uptake

is mediated through EAAT1 overexpression and that EAAT1 is the major contributor to glutamate clearance in astrocytes. Overall, the BBB can be considered as one of the main targets of biphalin and nociceptin when treating ischemic stroke. These opioid receptor agonists have protective effects on both the function and structure of the BBB under stress conditions. Biphalin and nociceptin maintain the integrity of tight junctions by improving the expression of claudin-5. Moreover, they enhance the protective function of astrocytes in clearing the extracellular glutamate by increasing the expression of glutamate transporter, EAAT1. These results have important clinical implications for possible prevention of vasogenic edema, which can be a life-threatening result of ischemic stroke.

Future studies are warranted to investigate the effect of nociceptin and biphalin on neurons in ischemic conditions. It is also crucial to consider the signaling mechanisms behind the increased expression of EAAT1 during stress condition. Since BBB comprises various cell types, studies using a triple culture model of the neurovascular unit cells would be remarkable to evaluate the impact of other cells on the expression of EAAT1 and the role of opioid receptors. The results of the current study provide an opportunity to design a preclinical study to investigate the BBB protection potential of opioid receptor agonists, as putative stroke treatments alone or in combination with thrombectomy or t-PA administration.

Authorship Contributions

Participated in research design: Nozohouri, Zhang, Albekairi, Vaidya, Abbruscato.

Conducted experiments: Nozohouri, Zhang, Albekairi.

Contributed new reagents or analytical tools: Vaidya, Abbruscato.

Performed data analysis: Nozohouri, Zhang, Albekairi.

Wrote or contributed to the writing of the manuscript: Nozohouri, Zhang, Albekairi, Vaidya, Abbruscato.

References

- András IE, Deli MA, Veszelka S, Hayashi K, Hennig B, and Toborek M (2007) The NMDA and AMPA/KAR receptors are involved in glutamate-induced alterations of occludin expression and phosphorylation in brain endothelial cells. *J Cereb Blood Flow Metab* **27**:1431–1443.
- Anttila JE, Albert K, Wires ES, Mätlik K, Loram LC, Watkins LR, Rice KC, Wang Y, Harvey BK, and Airavaara M (2018) Post-stroke intranasal (+)-naloxone delivery reduces microglial activation and improves behavioral recovery from ischemic injury. *eNeuro* **5**:ENEURO.0395-17.2018.
- Beschorner R, Simon P, Schauer N, Mittelbronn M, Schluesener HJ, Trautmann K, Dietz K, and Meyermann R (2007) Reactive astrocytes and activated microglial cells express EAAT1, but not EAAT2, reflecting a neuroprotective potential following ischaemia. *Histopathology* **50**:897–910.
- Chen C-J, Liao S-L, Chen W-Y, Hong J-S, and Kuo J-S (2001) Cerebral ischemia/reperfusion injury in rat brain: effects of naloxone. *Neuroreport* **12**:1245–1249.
- Chen H-J, Shen Y-C, Shiao Y-J, Liou K-T, Hsu W-H, Hsieh P-H, Lee C-Y, Chen Y-R, and Lin Y-L (2015) Multiplex brain proteomic analysis revealed the molecular therapeutic effects of Buyang Huanwu decoction on cerebral ischemic stroke mice. *PLoS One* **10**:e0140823.
- Choi J, Stradmann-Bellinghausen B, Yakubov E, Savaskan NE, and Régnier-Vigouroux A (2015) Glioblastoma cells induce differential glutamatergic gene expressions in human tumor-associated microglia/macrophages and monocyte-derived macrophages. *Cancer Biol Ther* **16**:1205–1213.
- Du F, Qian ZM, Zhu L, Wu XM, Qian C, Chan R, and Ke Y (2010) Purity, cell viability, expression of GFAP and bystin in astrocytes cultured by different procedures. *J Cell Biochem* **109**:30–37.
- Fallon M, Giusti R, Aielli F, Hoskin P, Rolke R, Sharma M, and Ripamonti C (2018) Management of cancer pain in adult patients: ESMO Clinical Practice Guidelines. *Ann Oncol* **29**:iv166–iv191.
- Feng Y, He X, Yang Y, Chao D, Lazarus LH, and Xia Y (2012) Current research on opioid receptor function. *Curr Drug Targets* **13**:230–246.
- Hamzei Moqaddam A, Ahmadi Musavi SM, and Khademizadeh K (2009) Relationship of opium dependency and stroke. *Addict Health* **1**:6–10.
- Ikonomidou C and Turski L (2002) Why did NMDA receptor antagonists fail clinical trials for stroke and traumatic brain injury? *Lancet Neurol* **1**:383–386.
- Li G, Simon MJ, Cancel LM, Shi Z-D, Ji X, Tarbell JM, Morrison 3rd B, and Fu BM (2010) Permeability of endothelial and astrocyte cocultures: in vitro blood-brain barrier models for drug delivery studies. *Ann Biomed Eng* **38**:2499–2511.
- Liang J, Chao D, Sandhu HK, Yu Y, Zhang L, Balboni G, Kim DH, and Xia Y (2014) δ -Opioid receptors up-regulate excitatory amino acid transporters in mouse astrocytes. *Br J Pharmacol* **171**:5417–5430.
- Liao S-L, Chen W-Y, Raung S-L, and Chen C-J (2003) Neuroprotection of naloxone against ischemic injury in rats: role of mu receptor antagonism. *Neurosci Lett* **345**(3):169–172.
- Cowell SM and Lee YS (2016) Biphalin: the foundation of bivalent ligands. *Curr Med Chem* **23**:3267–3284.
- Mark KS and Davis TP (2002) Cerebral microvascular changes in permeability and tight junctions induced by hypoxia-reoxygenation. *Am J Physiol Heart Circ Physiol* **282**:H1485–H1494.
- Meyer LC, Paisley CE, Mohamed E, Bigbee JW, Kordula T, Richard H, Lutfy K, and Sato-Bigbee C (2017) Novel role of the nociceptin system as a regulator of glutamate transporter expression in developing astrocytes. *Glia* **65**:2003–2023.
- Nozohouri S, Noorani B, Al-Ahmad A, and Abbruscato TJ (2021) Estimating brain permeability using in vitro blood-brain barrier models. *Methods Mol Biol* **2367**:47–72.
- Nozohouri S, Sifat AE, Vaidya B, and Abbruscato TJ (2020) Novel approaches for the delivery of therapeutics in ischemic stroke. *Drug Discov Today* **25**:535–551.
- Popiolek-Barczyk K, Piotrowska A, Makuch W, and Mika J (2017) Biphalin, a dimeric enkephalin, alleviates LPS-induced activation in rat primary microglial cultures in opioid receptor-dependent and receptor-independent manners. *Neural Plast* **2017**:3829472.
- Islam MR, Yang L, Lee YS, Hruba VJ, Karamyan VT, and Abbruscato TJ (2016) Enkephalin-fentanyl multifunctional opioids as potential neuroprotectants for ischemic stroke treatment. *Curr Pharm Des* **22**:6459–6468.
- Simantov R, Crispino M, Hoe W, Broutman G, Tocco G, Rothstein JD, and Baudry M (1999) Changes in expression of neuronal and glial glutamate transporters in rat hippocampus following kainate-induced seizure activity. *Brain Res Mol Brain Res* **65**:112–123.
- Thorlin T, Eriksson PS, Hansson E, and Rönnbäck L (1997) [D-Pen^{2,5}]enkephalin and glutamate regulate the expression of δ -opioid receptors in rat cortical astrocytes. *Neurosci Lett* **232**:67–70.
- Vaidya B, Sifat AE, Karamyan VT, and Abbruscato TJ (2018) The neuroprotective role of the brain opioid system in stroke injury. *Drug Discov Today* **23**:1385–1395.
- Verma R, Mishra V, Sasmal D, and Raghurib R (2010) Pharmacological evaluation of glutamate transporter 1 (GLT-1) mediated neuroprotection following cerebral ischemia/reperfusion injury. *Eur J Pharmacol* **638**:65–71.
- Voon P, Greer AM, Amlani A, Newman C, Burmeister C, and Buxton JA (2018) Pain as a risk factor for substance use: a qualitative study of people who use drugs in British Columbia, Canada. *Harm Reduct J* **15**:35.
- Wang CX and Shuaib A (2005) NMDA/NR2B selective antagonists in the treatment of ischemic brain injury. *Curr Drug Targets CNS Neurol Disord* **4**:143–151.
- Williams SM, Sullivan RK, Scott HL, Finkelstein DI, Colditz PB, Lingwood BE, Dodd PR, and Pow DV (2005) Glial glutamate transporter expression patterns in brains from multiple mammalian species. *Glia* **49**:520–541.
- Woller SA, Moreno GL, Hart N, Wellman PJ, Grau JW, and Hook MA (2012) Analgesia or addiction?: implications for morphine use after spinal cord injury. *J Neurotrauma* **29**:1650–1662.
- Xhima K, Weber-Adrian D, and Silburt J (2016) Glutamate induces blood-brain barrier permeability through activation of N-methyl-D-aspartate receptors. *J Neurosci* **36**:12296–12298.
- Yamashita A, Makita K, Kuroiwa T, and Tanaka K (2006) Glutamate transporters GLAST and EAAT4 regulate posts ischemic Purkinje cell death: an in vivo study using a cardiac arrest model in mice lacking GLAST or EAAT4. *Neurosci Res* **55**:264–270.
- Yang L, Islam MR, Karamyan VT, and Abbruscato TJ (2015) In vitro and in vivo efficacy of a potent opioid receptor agonist, biphalin, compared to subtype-selective opioid receptor agonists for stroke treatment. *Brain Res* **1609**:1–11.
- Yao Y, Chen Z-L, Norris EH, and Strickland S (2014) Astrocytic laminin regulates pericyte differentiation and maintains blood brain barrier integrity. *Nat Commun* **5**:3413.

Address correspondence to: Thomas J. Abbruscato, Department of Pharmaceutical Sciences, Texas Tech University Health Sciences Center, 1300 S Coulter St, Amarillo, TX 79106. E-mail: Thomas.abbruscato@ttuhsc.edu