

JPET #53496

METABOLISM OF FLAVONOIDS VIA ENTERIC RECYCLING: MECHANISTIC
STUDIES OF DISPOSITION OF APIGENIN IN CACO-2 CELL CULTURE MODEL

Ming Hu, June Chen, Huimin Lin

Department of Pharmaceutical Sciences, College of Pharmacy, Washington State

University, Pullman, WA 99164

JPET #53496

Running title: Intestinal Disposition Mechanisms of Apigenin

Address correspondence to:

Dr. Ming Hu

College of Pharmacy

Washington State University

Pullman, WA 99164-6510

Phone: (509)-335-4184

Fax: (509)-335-0162

E-mail: minghu@wsu.edu

Number of Text Page:	28
Number of Tables:	0
Number of Figures:	6
Number of References:	28
Number of Words in Abstract:	237
Number of Words in Introduction:	534
Number of Words in Discussion:	1011

List of Abbreviations:

HBSS: Hank's balanced salt solution; **MRP:** Multidrug resistance related protein; **OAT:** organic anion transporter; **UGT:** Uridine diphosphate-5'-glucuronosyltransferase or UDP-glucuronosyltransferase;

JPET #53496

ABSTRACT

The purpose of this study was to determine the mechanisms responsible for intestinal disposition of apigenin in the human Caco-2 cell culture model. The results indicated that most of the absorbed apigenin (10 μM) were conjugated and only a small fraction was transported intact. The amounts of conjugates excreted, especially that of the sulfate, were dependent on days-post-seeding. Apical efflux of apigenin sulfate did not change with concentration of apigenin (4 to 40 μM), whereas its basolateral efflux increased ($p < 0.01$) with concentration and plateaued at about 25 μM . In contrast, sulfate formation rates in cell lysate increased with concentration and plateaued at 25 μM and were 4-6 times faster than the corresponding excretion rates. Formation and polarized excretion rates of glucuronidated apigenin increased with apigenin concentration but formation rates were usually 2.5-6 times faster than the corresponding excretion rates. Inhibitors of multidrug resistance related proteins (MRPs) such as leukotriene C₄ and MK-571, which inhibited glucuronidation of apigenin at high concentration ($\geq 25 \mu\text{M}$), significantly decreased excretion of both apigenin conjugates, and higher concentrations of MK-571 increased the extent of inhibition. In contrast, an organic anion transporter (OAT) inhibitor oestrone sulfate only inhibited excretion of apigenin sulfate. In conclusion, we have showed for the first time that intestinal efflux is the rate-limiting step in the intestinal excretion of phase II conjugates of flavones. Furthermore, MRP and OAT are involved in the intestinal efflux of these hydrophilic phase II conjugates.

JPET #53496

Apigenin is an important and bioactive common flavonoid that is present in a variety of plants, vegetables, fruits, and herbs as well as their extracts, which are commonly marketed as diet and herbal supplements. Consumption of a diet rich in flavonoids is associated with a variety of health benefits (see reviews by Birt et al, 2001; Yang et al, 2001; and Harborne and Williams, 2000). Apigenin has demonstrated anticancer activities (e.g., Gupta et al, 2001; Wang et al, 2000; Ross and Kasum, 2002). Recently, apigenin has been shown to have the potential to prevent prostate cancer via cell cycle regulation, apoptosis, inhibition of androgen synthesis, or inhibition of oncogenic protein kinases (Gupta et al, 2001; 2002; Knowles et al, 2000; Kobayashi et al, 2002; Lee et al, 1998; Makela et al, 1998; Wang et al, 2000). However, apigenin, like many other flavonoids that are part of popular herbal supplements such as ginkgo biloba, is poorly bioavailable (Birt et al, 2001, Ross et al, 2002), and little is known about its absorption and metabolism. Like many other flavonoids, the in vitro anticancer effects of apigenin are represented by IC_{50} or EC_{50} values in the μM range, significantly higher than its in vivo concentration achievable after diet. Studies in rats and Caco-2 cells suggest that it is rapidly absorbed but more extensively metabolized than genistein, an isoflavone analog of apigenin (Liu and Hu, 2002, Chen et al, 2003). In HepG2 cells, apigenin was also metabolized rapidly via UGT1A1 (Galigatovic et al, 1999; Walle and Walle, 2002). Transdermal delivery of apigenin to prevent skin cancer has been used to avoid extensive metabolism and the results indicated that it was not metabolized significantly in skin (Li et al, 1996).

JPET #53496

Previously, we have proposed that repeated shuffling of flavonoids through a duo recycling scheme involving both enteric and enterohepatic recycling is the reason for extensive metabolism of apigenin (Liu and Hu, 2002). This metabolic scheme is supported by studies from independent groups of investigators, who also showed that intestine conjugated flavonoids and excreted their conjugates (Crespy et al, 1999; Andlauer et al, 2000). More recently, we have shown that intestinal disposition of apigenin is more important than hepatic disposition of apigenin, in that more apigenin conjugates were excreted into the rat lumen directly than those excreted via the bile (Chen et al, 2003). However, intestinal metabolism of flavonoids is poorly understood, and mechanistic studies that systemically study the intestinal metabolism of a flavone and enteric excretion of its phase II conjugates are lacking. For example, mechanisms responsible for efflux of phase II conjugates of flavonoids from the enterocytes have not been determined. Studies of effects of cellular differentiation and of changes in apigenin concentration on intestinal phase II conjugates of flavones are also lacking.

In this report, we systemically determined the pathways and kinetics of apigenin disposition and mechanisms responsible for metabolism of apigenin and excretion of apigenin conjugates in the Caco-2 cell monolayers. We have chosen Caco-2 cell model system because it is an established model of human intestinal epithelium. It was also chosen because apigenin is rapidly metabolized in this model, which allowed us to accurately measure amounts of apigenin metabolites excreted as a function of time.

JPET #53496

MATERIALS AND METHODS

Materials. Cloned Caco-2 cells, TC7, were a kind gift from Dr. Moniqué Rousset of INSERM U178 (Villejuif, France). MK-571 and leukotriene C₄ were purchased from BIOMOL (Plymouth Meeting, PA). Apigenin was purchased from Indofine Chemicals (Somerville, NJ). β -Glucuronidase with (Catalog#G1512) or without sulfatase (Catalog#G7396), sulfatase without glucuronidase (Catalog#S1629), uridine diphosphoglucuronic acid, alamethicin, D-saccharic-1,4-lactone monohydrate, magnesium chloride, oestrone sulfate, and Hank's balanced salt solution (HBSS, powder form) were purchased from Sigma Chem. Co. (St Louis, MO). All other materials (typically analytical grade or better) were used as received.

Cell Culture. The culture conditions for growing Caco-2 cells have been described previously (Hu et al, 1994a,b; Liu and Hu, 2002). The seeding density (100,000 cells/cm²), growth media (DMEM supplemented with 10% fetal bovine serum), and quality control criteria were all implemented in the present study as they were described previously (Hu et al, 1994a, b, Liu and Hu, 2002). Caco-2 TC7 cells fed every other day, and the monolayers were ready for experiments from 19-22 days post seeding except in days-post-seeding studies.

Transport Experiments in the Caco-2 Cell Culture Model. Experiments in triplicate were performed in pH 7.4 Hank's balanced salt solution (or HBSS) (Hu et al, 1994a, b). The protocol for performing cell culture experiments was from those described

JPET #53496

previously (Liu and Hu, 2002). Briefly, the cell monolayers were washed three times with 37°C pH 7.4 HBSS. The transepithelial electrical resistance values were measured, and those with transepithelial electrical resistance values less than 500 ohms \times cm² were discarded. The monolayers were incubated with the buffer for 1 hour and the incubation medium was then aspirated. Afterwards, the solution containing the compound of interest was loaded on to the APICAL side and amounts of transepithelial transport were followed as a function of time. Four donor samples (650 μ l) and four receiver samples (650 μ l) were taken every 30 min, followed by the addition of 650 μ l of donor solution to the donor side or 650 μ l of fresh buffer at the receiver side. A 200 μ l of methanol/acetonitrile containing 50 μ M of testosterone was added to each sample as internal standard and preservative. Afterwards, the mixture centrifuged at 13,000 rpm for 15 minutes, and the supernatant was analyzed by HPLC (see below).

Caco-2 Cell Lysate. Caco-2 cell monolayers (6) were put in 6 ml potassium phosphate (pH7.4 buffer), and sonicated in Aquasonic 150D sonicator (VWR Scientific, Bristol, CT) for 20 min at the maximum power (135 average watts) in an ice-cold water bath. The resulting cell lysate is then harvest and used fresh in the experiments.

Lysate Protein Concentration. Protein concentration of microsomal protein was determined using the Bio-Rad protein assay, using bovine serum albumin as standards.

Measurement of UGT Activities Using Cell Lysate. The incubation procedures for measuring UDP-glucuronosyltransferase (UGT) activities using cell lysate were as

JPET #53496

follows: (1) mix 135 μ l of cell lysate (final concentration \approx 0.7 mg protein/ml), magnesium chloride (0.88 mM), saccharolactone (4.4 mM), alamethicin (0.022 mg/ml); different concentrations of substrates in a 50 mM potassium phosphate buffer (pH 7.4); and uridine diphosphoglucuronic acid (3.5 mM, add last) to a final volume of 200 μ l; (2) incubate the mixture at 37°C for 10 or 30 min; and (3) stop the reaction by the addition of 50 μ l 94% acetonitrile/6% glacial acetic acid containing 100 μ M testosterone as an internal standard.

Measurement of Sulfotransferase Activities Using Cell Lysate. The incubation conditions for measuring sulfotransferases activities were as follows: (1) mix 90 μ l cell lysate (final concentration: 0.9 mg protein/ml), 3'-phosphoadenylyl sulfate (100 μ M), and substrate of an appropriate concentration (total volume 100 μ l) for 30 min; (2) stop the reaction by the addition of 25 μ l 94% acetonitrile/6% glacial acetic acid containing 100 μ M testosterone as an internal standard; (3) centrifuge the sample at 13,000 rpm for 15 min; (4) assay the sample via HPLC as described below.

HPLC Analysis of Apigenin and Its Conjugates. The conditions for HPLC analysis of isoflavones were modified based on a previously published method (Chen et al, 2003). The conditions for analyzing apigenin and its conjugates were: system, HP 1090 with diarray detector and HP Chemstation; column, Aqua (Phenomenex, Guilory, CA), 5 μ m, 150x0.45 cm; mobile phase A, 0.04% (w/v) phosphoric acid plus 0.06% (v/v) triethylamine (pH2.8); mobile phase B, 100% acetonitrile; gradient, 0-3 min, 20%B, 2-22, 20%-49%B, 22-23, 49%B; wavelength 340 nm; injection volume, 200 μ l. There was a five-minute interval between the end of the run and the next injection to allow the

JPET #53496

column to be re-equilibrated with 20% mobile phase B. The retention time of apigenin was 14.3 min, apigenin sulfate 11.0 min, and apigenin glucuronic acid 4.7 min.

Data Analysis.

Rate of apigenin transport or rate of apigenin conjugate excretion was determined using linear regression analysis of amounts transported or excreted versus time plots.

One-way ANOVA or an unpaired Student's t-test (Microsoft Excel) was used to analyze the data. The prior level of significance was set at 5%, or $p < 0.05$.

RESULTS

Apical to Basolateral Transport and Metabolite Excretion as a Function of Days-Post-Seeding in Intact Caco-2 Cell Monolayers.

We followed the transport and metabolism of apigenin (10 μ M) as a function of days-post-seeding (Fig.1). The results indicated that amounts of intact apigenin transported from the apical to basolateral side, rate of excretion of conjugates from both the apical and basolateral sides increased approximately linearly with time (Fig.1). The total amounts of sulfate and glucuronidated metabolites excreted were significantly ($p < 0.05$) dependent on days-post-seeding (Fig.2A). Maximal amounts of metabolites were excreted at 15 days-post-seeding, and the maximal amounts of intact apigenin were transported at 9 days-post-seeding (Fig.2A). The rates of metabolite excretion were also significantly dependent on days-post-seeding and the maximal excretion rate or maximal amounts excreted occurred at 15 days-post-seeding (Fig.2B). In contrast to the total rate, only basolateral excretion rates reached maximum at 15 days-post-seeding whereas apical excretion rates appear to have two maximums at 15 days-post-seeding and 24 days-post-seeding (Fig.2C). In addition to the dependence on days-post-seeding, more glucuronidated metabolites were excreted into the basolateral media, whereas more sulfated metabolites were excreted into the apical media (Fig.1). Lastly, higher total amounts of sulfated metabolites were excreted than glucuronidated metabolites at an apigenin concentration of 10 μ M (Fig.2B), which was primarily due to much faster apical excretion of sulfates (Fig.2C).

JPET #53496

Effects of Concentrations on the Disposition of Apigenin in Intact Cell Monolayers.

An increase in apical concentrations of apigenin from 5 μM to 40 μM significantly increase the apical to basolateral transport rates of apigenin, basolateral efflux rates of glucuronidated and sulfated conjugates, and apical efflux rates of glucuronidated conjugates, but not apical efflux rates of sulfated conjugates. At concentrations of 5 μM or less, the rates of apigenin transport (in intact form) were quite low. The transport rates increased linearly with concentration from 5 to 15 μM , after which the increase in transport rate accelerated (as signified by the rates above the straight line (Fig.3A)). In contrast, the basolateral excretion rates of both conjugates increased with concentration but gradually plateaued at higher concentrations (Fig.3B). On the apical membrane, excretion rates of glucuronic acid increased but excretion rates of sulfate did not change significantly with concentration (Fig.3C). In general, higher percentages of conjugates were excreted as sulfates at lower concentrations of apigenin but as glucuronic acid at higher concentrations (Fig.3B and Fig.3C).

Effect of Concentrations on the Metabolism of Apigenin in Caco-2 Cell Lysate.

Conjugated metabolism of apigenin was measured as a function of concentrations (2.5-50 μM) for 30 min (for sulfation) or 60 min (for glucuronidation) after preliminary studies demonstrated that amounts of metabolites were linear with time for at least 90 min (not shown). The results indicated that glucuronidation rates increased with concentration and started to approach but did not reach saturation (Fig.4A). In contrast, sulfation appeared to have reached saturation and started to show some level of substrate inhibition at high concentrations (≥ 50 μM) (Fig.4B). When the conjugation reaction rates were converted

JPET #53496

to nmol/hr per Caco-2 cell monolayer, these conjugation formation rates in cell lysate were substantially higher (2-6 folds) than the total excretion rates of conjugates from the Caco-2 cells (Fig.4).

Effects of MRP Inhibitors on the Disposition of Apigenin in Caco-2 Cell Lysate.

We were interested in the determination of effects of MRP inhibitors leukotriene C₄ and MK-571 on the efflux of conjugated apigenin since MRPs are known to transport glucuronic and sulfated conjugates in the liver by the hepatocytes (Konig et al, 1999). Therefore, we first determined if these inhibitors would affect the conjugation of apigenin via the UGT and sulfotransferase pathways. To our surprise, MK-571 inhibited glucuronidation of apigenin at concentration (25-50 μM) previously used to study the efflux mechanisms of these conjugates (Vaidyanathan and Walle 2001; O'Leary et al, 2003) (Fig.5). On the other hand, leukotriene C₄; did not inhibit the glucuronidation of apigenin. Lastly, neither MK-571 (up to 100 μM) nor leukotriene C₄ (up to 1.6 μM) inhibited the sulfation of apigenin.

Effects of MRP Inhibitors on the Disposition of Apigenin in Intact Caco-2 Cells.

MRP inhibitors leukotriene C₄ and MK-571 (the latter also inhibits glucuronidation at high concentrations) were used to determine the mechanisms by which conjugated apigenin is transported out of the Caco-2 cells. The results indicated that 0.1 μM leukotriene C₄ significantly (p<0.05) decreased the apical efflux rate of apigenin sulfate but significantly increased (p<0.05) its basolateral efflux rates when it is present on the basolateral side. Presence of this inhibitor at the apical side only or at both sides only slightly increased its basolateral efflux (p>0.05). Effects of leukotriene C₄ on excretion

JPET #53496

of glucuronidated metabolites were more complex, but the effects were generally moderate. For apical efflux of glucuronidated apigenin, presence of leukotriene C₄ at both sides of the monolayers decreased its efflux ($p < 0.05$) but the effect was moderate (32%). The presence of leukotriene C₄ at the basolateral side, however, significantly increased (up to 51%, $p < 0.05$) the basolateral efflux rate of apigenin glucuronic acid (Fig.6).

Unlike leukotriene C₄, 10 μ M MK-571, which is not expected to inhibit the formation of glucuronidated metabolites because of the low concentration, significantly ($p < 0.05$) inhibited efflux of glucuronidated conjugates (31% for basolateral and 44% for apical). MK-571 at 10 μ M also inhibited apical efflux of sulfated metabolites by 61% ($p < 0.05$) but only slightly decreased basolateral efflux of sulfated metabolites (20%, $p > 0.05$). When concentration of MK-571 was increased to 50 μ M, polarized excretion of both conjugates further decreased. Use of MK-571 and leukotriene C₄ greatly further reduced ($p < 0.05$) apical efflux of these conjugates when compared to using these inhibitors alone, but the combination did not enhanced the extent of basolateral excretion.

Effects of OAT Inhibitors on the Transport and Metabolism of Apigenin.

Oestrone sulfate has been shown to be the substrate of many prototypical anion transporters in the kidney and to a lesser extent in the liver cannicular membrane (Sekine et al, 2000). We found that the presence of the oestrone sulfate (10 μ M) in the apical and basolateral media did not significant affect excretion of apigenin glucuronic acid but significantly decreased apical and basolateral excretion of apigenin sulfate by 76% and 46% ($p < 0.01$), respectively (Fig.6).

JPET #53496

DISCUSSION

Apigenin is extensively metabolized in vivo and in cell culture models, and it is metabolized via the dual recycling scheme involving both enteric recycling and enterohepatic recycling (Chen et al, 2003; Liu and Hu, 2002; Galijatovic et al, 1999; Walle and Walle, 2002). Here we determined intestinal disposition of apigenin and the mechanisms by which its metabolites are excreted in the Caco-2 cells to gain better understanding of processes responsible for the intestinal flavonoid disposition.

We first defined a time window for the proposed studies by determining how transport and metabolism change as a function of days-post-seeding. Changes in days-post-seeding were shown to affect peptide transporter function and prolidase activities (Hu et al, 1995). The results indicated that most of absorbed apigenin were conjugated, regardless of days-post-seeding (Fig.1). The structures of the conjugates appeared to be the same too at different days-post-seeding because of similarities in: (1) metabolites' retention times, (2) the retention times of metabolites' hydrolysis products, and (3) the UV spectra of the metabolites (not shown). The types of flavonoid metabolites found here are similar to types of flavonoid conjugates found by Walle and his-coworkers (e.g., Galijatovic et al, 1999). Since the excretion rates were similar from 21 to 24 days-post-seeding (Fig.2), the later experiments were performed using cell monolayers of 19-22 days-post-seeding.

We also determined the effects of concentration on the disposition of apigenin, and found that cellular conjugation was saturable (a plot of concentration versus passive absorption

JPET #53496

rates deviated positively) (Fig. 3A). In intact cell monolayers, types of metabolites formed and how they were effluxed depended on the concentration (Fig.3B and Fig.3C). At low concentrations, the primary metabolite is apigenin sulfate, which is mainly excreted apically (Fig.3B and Fig.3C). At high concentrations, the primary metabolites became apigenin glucuronic acid, which was preferentially excreted basolaterally (Fig.3B and Fig.3C). This suggests that apical efflux transporter has high affinity and low capacity for apigenin sulfate, because apical efflux of apigenin sulfate was constant even though sulfate formation rates increased with concentrations (Fig.3B and Fig.3C, Fig.4B).

To explain the apparent lack of correlations between cellular sulfate formation rates and its excretion rates, we hypothesized that conjugate excretion could be rate-limited by their efflux rates. We found that the excretion rates of apigenin sulfate are much slower than its formation rates, especially at higher concentrations (Fig.4B). This important finding supports our hypothesis, which is novel since the current theory of metabolite production asserts that enzyme level is the most important factor in determining how much metabolite is produced. Therefore, when changes in phase II metabolite levels are observed in vivo or in vitro, the most likely experiments to do at the present are the determination of phase II enzyme expression levels. Our finding suggests that it will be necessary to determine the expression levels of efflux transporters as well.

Our hypothesis that cellular excretion of conjugates is rate-limited by their efflux will have significant impacts on drug induction and drug-drug interaction studies, if it is

JPET #53496

proven *in vivo*. Traditionally, phase II metabolism-related interaction studies always focus on the induction or the inhibition of enzymes. The results presented here suggest that it will be valuable to determine the induction or inhibition of the efflux transporters in drug-drug interaction studies involving hydrophilic phase II conjugates.

We believed that multidrug resistance related proteins (or MRPs) and organic anion transporters (OATs) are involved in the efflux of hydrophilic conjugates, based on what are known about their transport in kidney and liver (Konig et al, 1999; Sekine et al, 2001). We found that leukotriene C₄ inhibited apical efflux of both apigenin conjugates, but mildly increased basolateral efflux of apigenin conjugates. We also found that MK-571 (10 μM) was a highly effective inhibitor of apical efflux of apigenin conjugates, but was only mildly effective against basolateral efflux of apigenin conjugates. When used at a higher concentration, MK-571 (50 μM) was more effective against the excretion of conjugated metabolites, but its inhibition of cellular excretion of glucuronic acid is somewhat complicated by the fact that MK-571 also inhibited glucuronidation at this concentration. Taken together, these results support the hypothesis that the apical efflux is mediated by MRP and possibly other OATs, whereas basolateral efflux of these conjugates are probably mediated by an OAT since leukotriene C₄ was capable of stimulating efflux of both conjugates. The nature of this basolateral efflux transporter is currently unknown, but it may be similar to an organic anion transporter called rOAT1 because rOAT1 could interact with both oestrone sulfate (inhibition) and leukotriene C₄ (stimulation) (Sekine et al, 2000).

JPET #53496

The finding that intestinal Caco-2 cells secreted glucuronidated and sulfated metabolites via MRPs and oestrone sulfate sensitive organic transporter is novel. In general, mechanisms responsible for intestinal efflux of phase II conjugates of xenobiotics are poorly understood. We have hypothesized that these conjugates will be effluxed from the intestinal cells via MRPs based on the fact that cannicular MRPs transported many conjugates into the bile (Konig et al, 1999). Previously, only efflux of glutathione conjugates from the intestinal Caco-2 cells via MRP was shown (Gotoh et al, 2000; Terlouw et al, 2001). Now we have shown that MRPs, probably MRP2 based on the fact that MRP2 is located apically (Konig et al, 1999), are capable of excreting sulfated and glucuronidated conjugates of flavonoids. We have also shown that efflux of sulfates but not glucuronides may be mediated by the organic anion transporter similar to rOAT1. Based on previous observations that conjugates of many flavonoid are excreted into the lumen (Crespy et al 1999; Andlauer et al, 2000; Liu and Hu, 2002; Chen et al, 2003), MRP2 and an oestrogen sulfate sensitive transporter may be responsible for apical excretion of many flavonoid conjugates. Further studies are underway to define the structural characteristics of suitable flavonoid conjugates.

In summary, we have shown for the first time that intestinal excretion of hydrophilic phase II conjugates is rate-limited by the efflux transporters. We are the first to demonstrate that intestinal MRP and possibly oestrone sulfate sensitive OATs are involved in the apical and basolateral efflux of hydrophilic phase II conjugates of apigenin.

JPET #53496

REFERENCES

Andlauer W, Kolb J, Furst P (2000) Isoflavones from tofu are absorbed and metabolized in the isolated rat small intestine. *J Nutr.* **130**:3021-7.

Birt DF, Hendrich S, Wang W (2001) Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Ther.* **90**:157-77.

Chen J, Lin H, Hu M (2003) Metabolism of Flavonoids via Enteric Recycling: Role of Intestinal Disposition. *J Pharmacol Exp Ther.* **304**:1228-1235.

Crespy V, Morand C, Manach C, Besson C, Demigne C, Remesy C (1999) Part of quercetin absorbed in the small intestine is conjugated and further secreted in the intestinal lumen. *Am J Physiol.* **277**:G120-6.

Galijatovic A, Otake Y, Walle UK, Walle T (1999) Extensive metabolism of the flavonoid chrysin by human Caco-2 and Hep G2 cells. *Xenobiotica.* **29**:1241-56.

Gotoh Y, Suzuki H, Kinoshita S, Hirohashi T, Kato Y, Sugiyama Y. (2000) Involvement of an organic anion transporter (canalicular multispecific organic anion transporter/multidrug resistance-associated protein 2) in gastrointestinal secretion of glutathione conjugates in rats. *J Pharmacol Exp Ther.* **292**:433-9.

Gupta S, Afaq F, Mukhtar H (2001) Selective growth-inhibitory, cell-cycle deregulatory and apoptotic response of apigenin in normal versus human prostate carcinoma cells. *Biochem Biophys Res Commun.* **287**:914-20.

Gupta S, Afaq F, Mukhtar H. (2002) Involvement of nuclear factor-kappa B, Bax and Bcl-2 in induction of cell cycle arrest and apoptosis by apigenin in human prostate carcinoma cells. *Oncogene* **21**:3727-38.

JPET #53496

Harborne JB, Williams CA. (2000) Advances in flavonoid research since 1992.

Phytochemistry. **55**:481-504.

Hu M, Chen J, Tran D, Zhu Y, Leonardo G. (1994a) The Caco-2 cell monolayers as an intestinal metabolism model: metabolism of dipeptide Phe-Pro. *J Drug Target*. **2**:79-89.

Hu M, Chen J, Zhu Y, Dantzig A H, Stratford RE Jr, Kuhfeld MT (1994b) Mechanism and kinetics of transcellular transport of a new β -lactam antibiotic loracarbef across a human intestinal epithelial model system (Caco-2). *Pharm Res* **11**:1405-1413.

Hu M, Zheng L, Chen J, Liu L, Li Y, Dantzig AH, Stratford RE Jr. (1995) Peptide transporter function and prolidase activities in Caco-2 cells: a lack of coordinated expression. *J Drug Target*. **3**:291-300.

Knowles LM, Zigrossi DA, Tauber RA, Hightower C, Milner JA. (2000) Flavonoids suppress androgen-independent human prostate tumor proliferation. *Nutr Cancer* **38**:116-22.

Kobayashi T, Nakata T, Kuzumaki T. (2002) Effect of flavonoids on cell cycle progression in prostate cancer cells. *Cancer Lett* **176**:17-23.

Konig J, Nies AT, Cui Y, Leier I, Keppler D. (1999) Conjugate export pumps of the multidrug resistance protein (MRP) family: localization, substrate specificity, and MRP2-mediated drug resistance. *Biochim Biophys Acta* **1461**:377-94.

Lee SC, Kuan CY, Yang CC, Yang SD. (1998) Bioflavonoids commonly and potently induce tyrosine dephosphorylation/inactivation of oncogenic proline-directed protein kinase FA in human prostate carcinoma cells. *Anticancer Res*. **18**:1117-21.

Li B, Pinch H, Birt DF (1996) Influence of vehicle, distant topical delivery, and biotransformation on the chemopreventive activity of apigenin, a plant flavonoid, in mouse skin. *Pharm Res* **13**:1530-4.

JPET #53496

Liu Y and Hu M (2002) Absorption and metabolism of flavonoids in the Caco-2 cell culture model and a perfused rat intestinal model. *Drug Metab Dispos* **30**:370-7.

Makela S, Poutanen M, Kostian ML, Lehtimaki N, Strauss L, Santti R, Vihko R. (1998) Inhibition of 17beta-hydroxysteroid oxidoreductase by flavonoids in breast and prostate cancer cells. *Proc Soc Exp Biol Med* **217**:310-6.

O'Leary KA, Day AJ, Needs PW, Mellon FA, O'Brien NM, Williamson G. (2003) Metabolism of quercetin-7- and quercetin-3-glucuronides by an in vitro hepatic model: the role of human beta-glucuronidase, sulfotransferase, catechol-O-methyltransferase and multi-resistant protein 2 (MRP2) in flavonoid metabolism. *Biochem Pharmacol.* **65**:479-91.

Ross JA, Kasum CM. (2002) Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annu Rev Nutr* **22**:19-34.

Sekine T, Cha SH, Endou H. (2000) The multispecific organic anion transporter (OAT) family. *Pflugers Arch.* **440**:337-50.

Terlouw SA, Masereeuw R, van den Broek PH, Notenboom S, Russel FG. (2001) Role of multidrug resistance protein 2 (MRP2) in glutathione-bimane efflux from Caco-2 and rat renal proximal tubule cells. *Br J Pharmacol.* **134**:931-8.

Vaidyanathan JB, Walle T. (2001) Transport and metabolism of the tea flavonoid (-)-epicatechin by the human intestinal cell line Caco-2. *Pharm Res.* **18**:1420-5.

Walle UK, Walle T. (2002) Induction of human UDP-glucuronosyltransferase UGT1A1 by flavonoids-structural requirements. *Drug Metab Dispos.* **30**:564-9.

JPET #53496

Wang W, Heideman L, Chung CS, Pelling JC, Koehler KJ, Birt DF. (2000) Cell-cycle arrest at G2/M and growth inhibition by apigenin in human colon carcinoma cell lines. *Mol Carcinog.* **28**:102-10.

Yang CS, Landau JM, Huang MT, Newmark HL. (2001) Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annu Rev Nutr.* **21**:381-406.

JPET #53496

Footnote

*This work was supported by NIH AT00182 and CA87779.

JPET #53496

Figure Legends:

Fig.1. Effects of Days-Post-Seeding (DPS) on the Transport and Metabolism of Apigenin in the Caco-2 Cell Culture Model. Amounts of apigenin (or aglycone) (Panel A and D), glucuronidated metabolites (Panel B and E), and sulfated metabolites (Panel C and F) in the apical (AP, left three panels, or Panel A,B and C) or basolateral (BL, right three panels, or Panels D, E, and F) side were followed as a function of time (9 days, diamonds; 15 days, squares; 21 days, triangles; 24 days, hollow circles; 28 days, solid circles). Additional experimental details were provided in the “Materials and Methods” section. Each bar represents the average of three determinations, and the error bar is the standard deviation of the mean.

Fig.2. Effects of Days-Post-Seeding (DPS) on the Disposition of Apigenin in Intact Caco-2 Cells. Total amounts of apigenin aglycone transported (slashed bars), total amounts of sulfated (brick bars) and total amounts of glucuronic acid excreted in two hours were determined (Fig.2A) as a function of days-post-seeding. Total rates of conjugate excretion from both sides (Fig.2B) or from one side of the Caco-2 cell monolayers (Fig.2C) were also measured as a function of days-post-seeding. Additional experimental details were provided in the “Materials and Methods” section. Each bar represents the average of three determinations, and the error bar is the standard deviation of the mean. The symbols in Fig.2A indicated that days-post-seeding had a statistically significant effect amounts of transcellular transport of apigenin (*), cellular excretion of glucuronic acid (♣) and sulfate (♠). The arrows in Fig.2B and Fig.2C indicate

JPET #53496

statistically significant differences in the excretion of sulfate versus glucuronic metabolites.

Fig.3. Effects of Concentration on Transport and Metabolism of Apigenin. As apical apigenin concentration changed from 5 to 40 μ M, A apical to basolateral transport of apigenin (circles) (Fig.3A), basolateral excretion of apigenin sulfate (diamonds with solid curves) and glucuronic acids (squares with dashed curves) (Fig.3B), and apical excretion of same apigenin conjugates (Fig.3C) were measured. Each data point represents the average of three determinations, and the error bar is the standard deviation of the mean. The star symbol (*) represents statistically significant differences in the excretion rates of sulfate versus glucuronic acid. The regression lines represent the trend of the data and are not based on a kinetic model of metabolism or transport.

Fig.4. Effects of Concentration on the Formation and Excretion Rates of Apigenin Glucuronic Acid (Panel A) and Apigenin Sulfate (Panel B). Formation rates (hollow symbols with solid curves) of apigenin sulfate (diamonds) and apigenin glucuronic acid (squares) were measured in Caco-2 cell lysate, obtained from cell monolayers (21-22 days-post-seeding). Excretion rates (solid symbols with dashed curves) of apigenin conjugates were measured separately from the formation rates using cell monolayers of the same ages. To facilitate comparison, both rates were normalized to nmol/hr per cell monolayer or nmol/hr in the figure. We assumed that the activities of various enzymes in the cell lysate were the same as those present in intact cells, which created a slight

JPET #53496

negative bias since some enzyme activities are expected to be lost during lysate preparations. Each point represents the average of three determinations, and the error bar is the standard deviation of the mean. The regression lines represent the trend of the data and are not based on a kinetic model of metabolism or transport.

Fig.5. Effects of Different Concentrations of MK-571 and leukotriene C₄ (LTC₄) on the Formation Rates of Apigenin (10 μM) Conjugates in Caco-2 Cell Lysate. Effects of MK-571 concentrations on the formation rates of sulfate (diamonds) and glucuronic acid (squares) were determined as a function of MK-571 (Fig.5A) or leukotriene C₄ (Fig.5B) in Caco-2 cell lysate. The maximal MRP inhibitor concentrations used for cell lysate experiments were at least 2 times the maximal concentrations used in the intact cell studies. Each point (bar) represents the average of three determinations, and the error bar is the standard deviation of the mean. The symbol (*) indicates that the rate of metabolism is significantly less than the control.

Fig.6. Effects of MRP Inhibitors on Transport and Metabolism of Apigenin (10 μM). Control experiments in this figure are the experiments performed in the absence of any MRP inhibitors. MRP inhibitor leukotriene C₄ (LTC₄) or MK-571 was used alone or in combination with one another. We also used different concentrations of MK-571. We measured the apical to basolateral transport rate of apigenin (Panel A), basolateral excretion rate (Panel B) and apical excretion (Panel C) rates of apigenin glucuronic acid (slash bars) and apigenin sulfate (brick bars). Each bar represents the average of three determinations, and the error bar is the standard deviation of the mean. The symbol (*) indicates that the rate of metabolism is significantly different from the control.




Fig.1




Fig.2




Fig.3




Fig.4




Fig.5




Fig.6