

JPET #96792

Pharmacological characterization of (*E*)-N-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropane (LBT-999) as a highly promising fluorinated ligand for the dopamine transporter

Sylvie Chalon, Hakan Hall, Wadad Saba, Lucette Garreau, Frédéric Dollé, Christer Halldin, Patrick Emond, Michel Bottlaender, Jean-Bernard Deloye, Julie Helfenbein, Jean-Claude Madelmont, Sylvie Bodard, Zoïa Mincheva, Jean-Claude Besnard and Denis Guilloteau

Inserm, U619, Tours, France; Université François-Rabelais de Tours, France (S.C., L.G., P.E., S.B., Z.M., J.-C.B., D.G.); Karolinska Institutet, Stockholm, Sweden (H.H., C.H.); SHFJ, CEA/DSV, Orsay, France (W.S., F.D., M.B.); Laboratoires Cyclopharma, Saint-Beauzire, France (J.-B.D.); Orphachem, Clermont-Ferrand, France (J.H.); Inserm U484, Clermont-Ferrand, France (J.-C.M.)

JPET #96792

**Running title:** Pharmacological characterization of LBT-999

**Corresponding author:** Sylvie Chalon, Inserm U619; Université François-Rabelais de Tours,  
UFR Pharmacie, 31 avenue Monge, 37200, Tours, France

Tel: 33 2 47 36 72 43

Fax: 33 2 47 36 72 24

e-mail: [chalon@univ-tours.fr](mailto:chalon@univ-tours.fr)

Number of text pages: 23

Number of tables: 1

Number of figures: 6

Number of references: 39

Number of words in the Abstract: 224

Number of words in the Introduction: 508

Number of words in the Discussion: 840

**Abbreviations:** DAT, dopamine transporter;  $\beta$ -CIT, 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-iodophenyl)tropane; CBT, 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-bromophenyl)nortropane; CFT, 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-fluorophenyl)tropane; FPCBT, N-(3-fluoropropyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-bromophenyl)nortropane; FPCIT, N-3-fluoropropyl-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-iodophenyl)nortropane; FECNT, N-2-fluoroethyl-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-chlorophenyl)nortropane; IPT, N-(3'-iodopropen-2'-yl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-chlorophenyl)tropane; MADAM, N,N-dimethyl-2-(2-amino-4-methylphenyl thio)benzylamine; NET, norepinephrine transporter; PE2I, (*E*)-N-(3-iodoprop-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-methylphenyl)nortropane; PET, positron emission tomography ; SERT, serotonin transporter; SPECT, single photon computerized emission tomography.

**Section:** Neuropharmacology

JPET #96792

## ABSTRACT

In the aim to develop an efficient fluorinated probe for PET exploration of the dopamine transporter (DAT), we studied several in vitro and in vivo characteristics of the phenyltropane derivative (*E*)-N-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropane (LBT-999). In vitro on rat striatal membrane, [<sup>3</sup>H]LBT-999 bound to a single site with a K<sub>d</sub> of 9 nM, B<sub>max</sub> of 17 pmol/mg protein, and a very high selectivity for the DAT (IC<sub>50</sub> for GBR 12909 and PE2I: 2.4 and 18 nM, respectively; IC<sub>50</sub> for paroxetine, citalopram, MADAM, nisoxetine and desipramine > 1  $\mu$ M). In vitro on post-mortem human brain section, LBT-999 bound with high intensity to the caudate-putamen, weakly to the thalamus and not in the neocortex and cerebellum. This binding was totally abolished in the presence of PE2I. Ex vivo cerebral biodistribution of [<sup>11</sup>C]LBT-999 in rats showed a striatum/cerebellum radioactivity ratio of 18 and 25 at 30 and 60 min post-injection, respectively. This accumulation was strongly prevented by a pre-injection of GBR 12909 whereas paroxetine and nisoxetine had no effect. In vivo kinetic PET study in a Baboon showed a fast and very high uptake in the striatum, with a plateau at 30 min post-injection and a maximal putamen/cerebellum ratio of 30. Taken together, these findings demonstrate that LBT-999 is a highly promising agent for in vivo exploration of the DAT. This probe is currently labeled with <sup>18</sup>F for further characterizations.

## Introduction

As the membrane dopamine transporter (DAT) is very involved in a number of brain functions, in vivo exploration of this molecular target could become a relevant tool for the knowledge of pathophysiological mechanisms, diagnosis and follow-up of several neurological and psychiatric affections. In this field, scintigraphic exploration of the DAT could be very useful for the early diagnosis and follow-up of Parkinson's disease. Most of these studies have been performed using single photon emission computerized tomography (SPECT) radioligands chemically derived from the cocaine structure and labelled with  $^{123}\text{I}$  such as 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-iodophenyl)tropane ( $\beta$ -CIT) (Innis et al., 1993; Seibyl et al., 1998; Schwarz et al., 2000), N-3-fluoropropyl-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-iodophenyl)nortropane (FPCIT) (Booij et al., 1999), altropane (Fischmann et al., 1998), N-(3'-iodopropen-2'-yl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-chlorophenyl)tropane (IPT) (Schwarz et al., 2000) or with  $^{99\text{m}}\text{Tc}$  such as TRODAT (Mozley et al., 2000). A major challenge of these investigations is to approach the most accurate in vivo quantification of the DAT density. For this specific aim, it seems that positron emission tomography (PET) imaging has better performances due to a higher resolution than SPECT. For PET imaging, several DAT radioligands labeled with  $^{11}\text{C}$  are now proposed (Leenders et al., 1990; Guttman et al., 1997; Ilgin et al., 1999). However, as  $^{18}\text{F}$  can be more largely used for clinical purpose than  $^{11}\text{C}$  due to its longer physical period (110 versus 20 min), this is an increasing need to make available high-performance radioligands labeled with  $^{18}\text{F}$ . To date several cocaine derivatives labelled with  $^{18}\text{F}$  have been developed such as FPCIT (Chaly et al., 1996; Kazumata et al., 1998; Ma et al., 2002), N-(3-fluoropropyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-bromophenyl)nortropane (FPCBT) (Chaly et al., 2004), 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-fluorophenyl)tropane (CFT) (Haaparanta et al., 1996; Laakso et al., 1998; Nurmi et al., 2000), FPCT (Goodman et al., 1997) and FECNT (Goodman et al., 2000; Deterding et al., 2001; Davis et al., 2003). Although they are interesting potential tools for PET exploration of the DAT, most of them require a prolonged period (>100 min or more) to reach specific binding equilibrium with the DAT and/or have a moderate selectivity for the DAT over serotonin transporter (SERT). The most adapted available ligand seems to be N-2-fluoroethyl-2 $\beta$ -carbomethoxy-3 $\beta$ -(4-chlorophenyl)-nortropane (FECNT), with the most attractive kinetic characteristics by achieving peak striatal uptake at 60-75 min post-injection. As we previously developed the tropane derivative (*E*)-N-(3-iodoprop-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-

JPET #96792

methylphenyl)nortropine (PE2I) characterized by the presence of a methyl group on the phenyl ring (Emond et al., 1997) which is highly efficient in terms of DAT selectivity and in vivo kinetics (Chalon et al., 1999; Pinborg et al., 2002; Halldin et al., 2003), we recently developed a fluorinated compound derived from the PE2I structure (Dollé et al., in press). We present here the pharmacological properties of this new compound, (*E*)-N-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropine (LBT-999). For this characterization, we labeled LBT-999 with  $^3\text{H}$  for in vitro studies on brain tissue and with  $^{11}\text{C}$  for in vivo studies in the rat and monkey brain. As the present results show that it could be a very efficient tool for in vivo DAT quantification, LBT-999 is currently labeled with  $^{18}\text{F}$  for a detailed study of its in vivo properties in the primate.

## Materials and Methods

### Preparation of non-labeled LBT-999 and [<sup>3</sup>H/<sup>11</sup>C]LBT-999

(*E*)-*N*-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropine (LBT-999) and its *N*-desmethylated precursor for labeling (Nor-LBT-999, (*E*)-*N*-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropine) were synthesized as described elsewhere (Dollé et al., in press).

The labeling of LBT-999 with tritium at its methyl ester function was performed as follows: 0.7 mg of nor-LBT-999 diluted in 300  $\mu$ l of *N,N*-dimethyl formamide was mixed with 150  $\mu$ l of [<sup>3</sup>H]methyl iodide (370 MBq/ml, specific activity 3.1 TBq/mmol, Amersham, Sweden) and heated for 15 min at 90°C. After cooling to room temperature, 300  $\mu$ l of acetonitrile was added, and the [<sup>3</sup>H]LBT-999 was purified by HPLC using a C18 reverse phase column ( $\mu$ Bondapak, Waters) and CH<sub>3</sub>CN/H<sub>2</sub>O/TFA (35/65/0.1) as mobile phase. [<sup>3</sup>H]LBT-999 was easily separated from the precursor and non-reacted [<sup>3</sup>H]methyl iodide, and obtained with a specific activity of 3.1 TBq/mmol.

The labeling of LBT-999 with carbon-11 ( $T_{1/2}$ : 20.4 min) at its methyl ester function was performed using standard conditions that have so far been used for the routine radiosynthesis of several radiotracers (Langer et al., 1999; Sandell et al., 2000; Dollé et al., 2000, 2002) and the efficient methylation reagent [<sup>11</sup>C]methyl triflate. The conditions used were the following: (1) trapping at room temperature of [<sup>11</sup>C]methyl triflate in 300  $\mu$ L of acetone containing 0.7 mg of nor-LBT-999 and 3  $\mu$ L of a 3 M solution of NaOH in water; (2) concentration to dryness of the reaction mixture (at 110°C, using a helium stream for 1 min); (3) taking up the residue with 0.5 mL of the HPLC mobile phase and (4) HPLC purification using semi-preparative C18 reverse phase column (Symmetry C-18, Waters) and CH<sub>3</sub>CN/H<sub>2</sub>O/TFA (35/65/0.1) as mobile phase. Typically, 5.55 to 9.25 GBq of [<sup>11</sup>C]LBT-999 were routinely obtained within 30 min of radiosynthesis (including HPLC purification) with specific radioactivity ranging from 29.6 to 44.4 GBq/ $\mu$ mol.

### Animals and drugs

All animal use procedures were in accordance with the recommendations of the European Community (86/609/CEE) and the French National Committee (décret 87/848) for the care and use of laboratory animals. Experiments in rats were conducted on male Wistar animals

JPET #96792

weighing 250-300 g (Centre d'Elevage R. Janvier, Le Genest St Isle, France). PET experiment was conducted in 3 male *Papio anubis* baboon weighing 10-15 kg.

Desipramine and nisoxetine were obtained from RBI Bioblock (Illkirch, France); paroxetine was a gift from Smith-Kline-Beecham (Nanterre, France); citalopram was a gift from Lundbeck (Copenhagen, Denmark); (*E*)-*N*-(3-iodoprop-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-methylphenyl)nortropane (PE2I) and *N,N*-dimethyl-2-(2-amino-4-methylphenylthio)benzylamine (MADAM) were synthesized according to previously published data (Emond et al., 1997, 2002).

### **In vitro binding assays on rat striatal membranes**

#### *Tissue preparation*

Male rats were killed by decapitation on the day of the assay and both striata of each animal were removed on ice and weighed (2 rats were used for each experiment). The tissue was homogenized in 10 vol of 0.32 M sucrose using an Ultraturrax T25. After 1000  $\times$  g centrifugation at 4°C for 10 min, the supernatant was kept and the pellet was treated as described above. Both supernatants were then pooled and centrifuged at 17,500  $\times$  g for 30 min at 4°C. Twenty vol of the incubation buffer was added to the pellet, and the mixture was homogenized and centrifuged at 50,000  $\times$  g for 10 min at 4°C. The final pellet was suspended in a minimum volume of the assay buffer and the protein concentration was measured according to Bradford (1976) using bovine serum albumin as standard.

#### *Saturation studies*

[<sup>3</sup>H]LBT-999 (3 KBq, 10 pmoles) was incubated with unlabeled LBT-999 at concentrations varying between 0.2 and 50 nM with 30  $\mu$ g protein in a total volume of 1 ml in a pH 7.4 Tris-HCl buffer (50 mM Tris-HCl, 120 mM NaCl, 5 mM KCl) for 90 min at 22°C. Nonspecific binding was determined in the presence of 1  $\mu$ M PE2I. Samples were then rapidly filtered through Whatman GF/C fiber filters soaked with 0.05% polyethylenimine (Sigma, St Quentin-Fallavier, France). The filters were washed twice with 4 ml of cold buffer, and the residual radioactivity was measured in a  $\beta$  counter (LKB 1215 Rackbeta) in the presence of 6 ml scintillator (LKB Optiphase). Results were analyzed with the EBDA RADLIG program (Biosoft, Cambridge, UK).

JPET #96792

### *Competition studies*

For these studies, [<sup>3</sup>H]LBT-999 (3 KBq, 10 pmoles) together with 10 nM of unlabeled LBT-999 were incubated with 30 µg of protein in a total volume of 1 ml in a pH 7.4 Tris-HCl buffer (50 mM Tris-HCl, 120 mM NaCl, 5 mM KCl) for 90 min at 22°C, in the presence of different drugs: PE2I or GBR12935 at concentrations of 10<sup>-8</sup> to 10<sup>-11</sup> M, paroxetine, citalopram, MADAM, desipramine or nisoxetine at concentrations of 10<sup>-5</sup> to 10<sup>-10</sup> M. Samples were then treated as described above. Total binding was determined in the absence of any drug, and nonspecific binding was measured in the presence of 1 µM PE2I. The IC<sub>50</sub> values were determined graphically for each compound.

### **In vitro autoradiographic studies on human post-mortem brains**

The human brain used was obtained from clinical autopsy at the National Institute of Forensic Medicine (Karolinska Institutet, Stockholm, Sweden), and handled as previously described (Hall et al., 1998, 2001). The study was approved by the Ethics Committee at Karolinska Institutet and the Swedish Board of Social Welfare. Cryosectioning on whole hemisphere sections was performed as previously described (Hall et al., 1998, 2001). Experiments were performed on 100-µm horizontal sections.

For the autoradiography studies, sections were incubated for 60 min at 22°C with 2 nM of [<sup>3</sup>H]LBT-999 in 10 ml of a phosphate buffer (10.14 mM NaH<sub>2</sub>PO<sub>4</sub>, 137 mM NaCl, 2.7 mM KCl, 1.76 mM KH<sub>2</sub>PO<sub>4</sub>), pH 7.4. Non-specific binding was defined on adjacent sections incubated in the presence of excess (10 µM) of PE2I. Sections were then washed twice for 10 min in cold phosphate buffer and rinsed for 1 sec in distilled water. After drying, the sections were exposed to film (Kodak Biomax MR, Amersham) for 12 weeks before development (developer: Kodak D19, fixation: Kodak Fixer 3000). The autoradiograms were digitized using a Scan Maker E6 high-resolution scanner (Microtek) and Adobe PhotoShop 6.5.

### **Ex vivo cerebral biodistribution in rats**

#### *Kinetic study*

Twelve rats received an i.v. injection of [<sup>14</sup>C]LBT-999 (2.22 MBq) and were sacrificed 10, 30 or 60 min post-injection (4 animals per group). Samples of blood and cerebral regions (cerebellum, frontal cortex and striatum) were removed, rapidly wiped (brain samples) and weighed, and radioactivity was measured in a γ counter (Cobra Quantun, Packard). Uptake was expressed as the percentage of injected dose per gram of tissue (%ID/g tissue).

JPET #96792

### *Competition study*

Six rats per group received an i.v. injection of [ $^{11}\text{C}$ ]LBT-999 (0.74 MBq) without (control) or with a pre-injection (3 min) of drug (GBR 12909, paroxetine or nisoxetine) at a dose of 5 mg/kg i.v. They were sacrificed 30 min after injection of [ $^{11}\text{C}$ ]LBT-999. Samples of blood and cerebral regions (cerebellum, frontal cortex and striatum) were removed, rapidly wiped (brain samples) and weighed, and radioactivity was measured in a  $\gamma$  counter (Cobra Quantun, Packard). Uptake was expressed as the percentage of injected dose per gram of tissue (%ID/g tissue).

### **In vivo Positron Emission Tomography studies in Baboons**

PET study was performed with a CTI HR + Exact positron tomograph (CTI PET Systems, Knoxville, TN, USA). This scanner allowed simultaneous acquisition of 63 slices every 2.2 mm with a spatial and axial resolutions of 4.5 mm. Each animal was anesthetized using 1% isoflurane and 33%/66%  $\text{O}_2/\text{N}_2\text{O}$ , controlled by an Ohmeda ventilator (Ohmeda OAV 7710, Madison, WI). The tidal volume was adjusted to achieve stable end-tidal carbon dioxide tension between 38 to 40 mm Hg. The baboon's head was fixed in a head-holder and positioned in the scanner gantry for axial plane acquisition. A venous catheter was inserted for [ $^{11}\text{C}$ ]LBT-999 administration and a femoral arterial catheter was used for blood sampling. A transmission scan ( $^{68}\text{Ge}$  rods) was acquired for 15 min and used for subsequent correction attenuation of emission scans. The image acquisition started (T0 min) at the intravenous injection of [ $^{11}\text{C}$ ]LBT-999 (419 MBq; specific radioactivity, 14 GBq/ $\mu\text{mol}$ ) and lasted 80 min (T80 min). Twenty-four images were acquired with scan duration starting from 30 s and increasing up to 10 min during each experiment.

### *PET data analysis*

The regions of interest (ROIs) were identified and drawn on 1.5 T MRI scans images (SIGNA system, General Electric, Milwaukee, WI, USA), which were matched with PET images (anatomist) and resliced according to the PET images. The ROIs were then copied onto the PET images. The average of the radioactivity concentration in the different ROIs were then calculated and expressed as Bq/mL. The means of the corresponding left and right hemisphere ROI values were averaged. Tissue-time activity curves were constructed by plotting the percent injected dose per 100 mL tissue (%ID/100mL). The cerebellum was used as reference in the analysis. The ROIs to cerebellum ratios was at 110 minutes post-injection.

JPET #96792

### **Statistical analysis**

For ex vivo cerebral biodistribution studies in rats, comparisons between groups (competition study) were performed using Student's test.

## Results

### **In vitro binding on rat striatal membranes**

#### *Saturation studies*

The affinity and density of specific [<sup>3</sup>H]LBT-999 binding sites were measured by saturation experiments on rat striatal membranes. In our experimental conditions, the nonspecific binding, determined in the presence of 1 μM PE2I was around 1.5%. The Scatchard transformation of the resulting data (Fig.1) revealed a linear curve suggesting a one site model (Hill coefficient = 1) with a *K<sub>d</sub>* value of  $9.15 \pm 2.8$  nM (mean  $\pm$  SD of 3 independent determinations, each performed in triplicate) and a *B<sub>max</sub>* value of  $17.47 \pm 2.73$  pmol/mg protein (mean  $\pm$  SD).

#### *Competition studies*

The pharmacological profile of specific [<sup>3</sup>H]LBT-999 binding in the striatum was consistent with binding to the DAT (Fig.2), as the rank order of potency of membrane transporter-drugs was GBR 12909 = PE2I > nisooxetine > desipramine = citalopram = paroxetine = MADAM (Table 1).

### **Autoradiographic studies in human post-mortem brain sections**

As shown on Fig.3, [<sup>3</sup>H]LBT-999 bound with high intensity to the basal ganglia (putamen, nucleus caudatus) of the human brain section. Weak binding could be seen in the thalamus, whereas there was virtually no or very low [<sup>3</sup>H]LBT-999 binding in the cortical regions. The labelling of DAT with [<sup>3</sup>H]LBT-999 was totally abolished in the presence of 10 μM PE2I.

### **Ex vivo cerebral biodistribution in rats**

#### *Kinetic study*

Whole-blood radioactivity showed stable values 10 to 60 min post-injection ( $0.25 \pm 0.02$ ,  $0.22 \pm 0.02$ ,  $0.15 \pm 0.01$  %ID/g, respectively). Whatever the time post-injection, the highest accumulation of [<sup>11</sup>C]LBT-999 was observed in the striatum compared to the cortex and cerebellum (Fig.4). Radioactivity in the striatum was stable from 10 to 30 min post-injection ( $4.90 \pm 0.36$  and  $4.95 \pm 0.25$  %ID/g, respectively) and was slightly decreased at 60 min ( $3.37 \pm 0.61$  %ID/g). The cortex and cerebellum displayed a weak degree of radioactivity from 10 to 60 min post-injection. The radioactivity ratio striatum/cerebellum reached 7, 18 and 25 at 10, 30 and 60 min post-injection, respectively.

JPET #96792

### *Competition study*

As shown in Fig.5, in vivo cerebral uptake of [ $^{11}\text{C}$ ]LBT-999 in control rats reached  $5.03 \pm 0.7$  %ID/g in the striatum, with a ratio striatum/cerebellum of 15 at 30 min post-injection. Pre-injection of GBR 12909 (5 mg/kg i.v.) strongly decreased the uptake of [ $^{11}\text{C}$ ]LBT-999 (around 70%) in the striatum ( $1.16 \pm 0.3$  %ID/g,  $p < 0.01$ ), whereas no effect was observed consecutively to a paroxetine pre-injection ( $3.99 \pm 0.3$  %ID/g), and a slight effect was observed with a nisoxetine pre-injection ( $3.44 \pm 0.5$  %ID/g,  $p < 0.05$ ).

In the cerebellum, the radioactivity level remained low and similar in the four groups of animals ( $0.29 \pm 0.07$ ,  $0.27 \pm 0.06$ ,  $0.23 \pm 0.06$ ,  $0.26 \pm 0.07$  %ID/g for the control, GBR 12909, paroxetine and nisoxetine groups, respectively).

In the cortex, a 30% decrease in the accumulation of [ $^{11}\text{C}$ ]LBT-999 was observed in the GBR 12909 compared to the control group ( $0.36 \pm 0.04$  vs  $0.58 \pm 0.18$  %ID/g,  $p < 0.05$ ), whereas no significant difference was observed for the paroxetine ( $0.51 \pm 0.10$  ID/g) and nisoxetine ( $0.64 \pm 0.07$  ID/g) groups.

### **In vivo PET study in Baboon**

Time activity curves (Fig.6) of the concentration of [ $^{11}\text{C}$ ]LBT-999 in different regions of the brain show that the highest accumulation was observed in the striatum with a peak uptake at 40 min ( $16.3$  and  $13.3$  %ID/100mL in putamen and caudate, respectively), the washout was then very slow from 30 to 80 min. Radioactivity uptake in the midbrain peaked at 12 min ( $6.9$ %ID/100mL) and decreased with time. The lowest uptake was observed in the cortex and cerebellum ( $1.37$  and  $1.09$  %ID/100mL, respectively).

The ROIs to cerebellum radioactivity ratio were calculated at 80 min post injection. This ratio was very high in the striatum (30 and 24.6 in the putamen and caudate, respectively), intermediate in the midbrain (5.9), low in the thalamus (2.1) and cortical structure (1.2).

## Discussion

In the aim to develop at term an efficient DAT probe for PET exploration with  $^{18}\text{F}$ , we studied several in vitro and in vivo characteristics of the phenyltropane derivative (*E*)-*N*-(4-fluorobut-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-tolyl)nortropine (LBT-999) labelled in the present study with  $^3\text{H}$  or  $^{11}\text{C}$ , and demonstrated that this ligand is a highly promising imaging agent. Its in vitro pharmacological properties on rat striatal membranes are very close to those of PE2I (Chalon et al., 1999). It binds to a single high affinity site with a moderate affinity (9 nM) close related to that of other fluorinated tropane derivatives such as FPCIT (Neumeyer et al., 1996), FPCBT (Chaly et al., 2004), CFT (Meltzer et al., 1993), FPCT (Goodman et al., 1997), FECNT (Goodman et al., 2000) and PE2I (Chalon et al., 1999), and has a very high selectivity for the DAT. The  $\text{IC}_{50}$  values of GBR 12935 and PE2I vs LBT-999 were 2.4 and 18 nM, respectively, and we did not find competition ( $\text{IC}_{50} > 1 \mu\text{M}$ ) between LBT-999 and several SERT ligands (citalopram, paroxetine, *N,N*-dimethyl-2-(2-amino-4-methylphenyl thio)benzylamine) as well as norepinephrine transporter (NET) ligands (nisoxetine, desipramine). This selectivity is higher than that observed in vitro for other DAT ligands from the chemical tropane family such as FPCIT (Neumeyer et al., 1996), FPCBT (Chaly et al., 2004), CFT (Meltzer et al., 1993), FPCT (Goodman et al., 1997), FECNT (Goodman et al., 2000). The selectivity of LBT-999 as indicated from the in vitro binding data on rat brain was clearly confirmed in the post-mortem human autoradiographic study where it bound highly to the caudate-putamen and only weakly to the thalamus and neocortical areas, as already observed for PE2I (Hall et al., 1999). This binding was achieved at concentration (2 nM) that was near to the  $K_d$ , and was totally abolished in the presence of a saturating concentration of PE2I.

These homogenate and autoradiographic assessments were thus very useful as an initial view of the affinity/selectivity properties of LBT-999 as a DAT probe, but did not totally predict its in vivo characteristics. In a first step, ex vivo cerebral biodistribution in rats demonstrated that the ligand had a rapid and high entry into the brain where its distribution was in agreement with a specific binding to the DAT. The brain entrance of LBT-999 in rats (mean value of 4.95 % ID/g in the striatum at 30 min post-injection) was higher than for PE2I (1.06% in same experimental design, Guilloteau et al., 1998), as well as the striatum/cerebellum ratio (18 for LBT-999 vs 10 for PE2I). The striatal accumulation of LBT-999 was around 70% decreased in the presence of a saturating dose of GBR12909, as already observed for PE2I. In contrast, no significant effect was observed with a pre-injection of paroxetine and a low significant

JPET #96792

18% decrease was observed with nisoxetine. It can be hypothesized that this latter corresponds to a non-specific binding rather than a binding of LBT-999 to the norepinephrine transporter which is practically absent from the striatum (Sanders et al., 2005).

In vivo kinetic study in baboon confirmed that LBT-999 brain uptake was fast, high, and mainly localized in the putamen and caudate. The peak uptake in these regions reached 17.1 and 12.8%ID/100 ml, respectively, at 30 min post-injection. This peak appeared earlier than that observed for [<sup>11</sup>C]PE2I in human brain (55-65 min p.i.; Halldin et al., 2003) and for [<sup>18</sup>F]-labelled DAT tracers such as CFT (225 min p.i.; Laakso et al., 1998), FPCT (60 min p.i.; Goodman et al., 1997), FECNT (90 min p.i.; Davis et al., 2003), FPCIT (40 min p.i.; Kazumata et al., 1998). An undesired property accompanying DAT binding by the fluorinated tracers available to date is the need for a prolonged time required to reach binding equilibrium, and it can be hypothesised from these preliminary data that LBT-999 would not have this drawback. The regional distribution of LBT-999 in baboon brain was in agreement with the known DAT localization with highest density in the putamen and caudate, lower in the midbrain, much lower in the thalamus and very low in the cortex and cerebellum. The maximum striatum/cerebellum ratio reached the very high values of 34.9 and 26.5 in the putamen and caudate, respectively. This ratio was higher than the ratios previously obtained for all previously cited tracers, thus showing that LBT-999 is the in vivo DAT probe which possesses the highest signal-to-noise ratio. The specificity of the LBT-999 in vivo binding on the DAT in the baboon brain is currently studied using competition experiments. From the present data obtained in vitro on rat striatal membranes and post-mortem human brain as well as ex vivo in the rat, it can be strongly hypothesized that LBT-999 will selectively bind to the DAT in primate brain. Taken together, these findings suggest that LBT-999 could be a very efficient tool for in vivo detection of DAT variations, thus allowing e.g. to early diagnose, monitor progression or assess efficacy of therapies of Parkinson's disease; as well as to improve the aetiology and pathophysiology of less known CNS affections such as attention-deficit/hyperactivity disorders (Jucaite et al., 2005).

## References

Booij J, Hemelaar TG, Speelman JD, de Bruin K, Janssen AG, van Royen EA (1999) One-day protocol for imaging of the nigrostriatal dopaminergic pathway in Parkinson's disease by [<sup>123</sup>I]FPCIT SPECT. *J Nucl Med* **40**: 753-761.

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254.

Chaly T, Dhawan V, Kazumata K, Antonini A, Margouleff C, Dahl R, Belakhlef A, Margouleff D, Yee A, Wang S, Tamagnan G, Neumeyer JL, Eidelberg D (1996) Radiosynthesis of [<sup>18</sup>F]N-3-fluoropropyl-2- $\beta$ -carbomethoxy-3- $\beta$ -(4-iodophenyl)nortropine and the first human study with Positron Emission Tomography. *Nucl Med Biol* **23**: 999-1004.

Chaly T, Baldwin RM, Neumeyer JL, Hellman MJ, Dhawan V, Garg PK, Tamagnan G, Staley JK, Al-Tikriti MS, Hou Y, Zoghbi SS, Gu X-H, Zong R and Eidelberg D (2004) Radiosynthesis of [<sup>18</sup>F]N-(3-fluoropropyl)-2- $\beta$ -carbomethoxy-3- $\beta$ -(4-bromophenyl)nortropine and the regional brain uptake in non human primate using PET. *Nucl Med Biol* **31**: 125-131.

Chalon S, Garreau L, Emond P, Zimmer L, Vilar M P, Besnard JC, Guilloteau D (1999) Pharmacological characterization of (E)-N-(3-iodoprop-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-methylphenyl)nortropine as a selective and potent inhibitor of the neuronal dopamine transporter. *J. Pharmacol. Exp. Ther.* **291**, 648-654.

Davis MR, Votaw JR, Bremner JD, Byas-Smith MG, Faber TL, Voll RJ, Hoffman JM, Grafton ST, Kilts CD, Goodman MM (2003) Initial human PET imaging studies with the dopamine transporter ligand [<sup>18</sup>F]FECNT. *J Nucl Med* **44**: 855-861.

Deterding TA, Votaw J, Wang CK, Eshima D, Eshima L, Keil R, Malveaux E, Kilts CD, Goodman MM, Hoffman JM (2001) Biodistribution and radiation dosimetry of the dopamine transporter ligand [<sup>18</sup>F]FECNT. *J Nucl Med* **42**: 376-381.

JPET #96792

Dollé F, Bottlaender M, Demphel S, Emond P, Fuseau C, Coulon C, Ottaviani M, Valette H, Loc'h C, Halldin C, Mauclaire L, Guilloteau D, Mazière B, Crouzel C (2000) Highly efficient synthesis of [<sup>11</sup>C]PE2I, a selective radioligand for the quantification of the dopamine transporter using PET. *J Label Compounds Radiopharm* **43**: 997-1004.

Dollé F, Bramoullé Y, Hinnen F, Fowler JS (2002) Efficient synthesis and formulation of (R)-(-)-[<sup>11</sup>C]Deprenyl, a selective radioligand for the quantification of MAO-B activity using PET. *J Label Compounds Radiopharm* **45**: 803-811.

Dollé F, Emond P, Mavel S, Demphel S, Hinnen F, Mincheva Z, Saba W, Valette H, Chalon S, Halldin C, Helfenbein J, Legaillard J, Madelmont JC, Deloye JB, Bottlaender M, Guilloteau D. Synthesis, radiosynthesis and in vivo preliminary evaluation of [<sup>11</sup>C]LBT-999, a selective radioligand for the visualization of the dopamine transporter with PET. *Bioorg Med Chem*, in press.

Emond P, Garreau L, Chalon S, Boazi M, Caillet M, Bricard J, Frangin Y, Mauclaire L, Besnard J-C and Guilloteau D (1997) Synthesis and ligand binding of nortropine derivatives: N-substituted 2β-carbomethoxy-3β-(4'-iodophenyl) nortropine and N-(3-iodoprop-2E-enyl)-2β-carbomethoxy-3β-(3',4'-disubstitutedphenyl) nortropine. New high-affinity and selective compounds for the dopamine transporter. *J Med Chem* **40**: 1366-1372.

Emond P, Vercouillie J, Innis R, Chalon S, Mavel S, Frangin Y, Halldin C, Besnard J-C, Guilloteau D (2002) Substituted diphenyl sulfides as selective serotonin transporter ligands: synthesis and in vitro evaluation. *J Med Chem* **45**: 1253-1258.

Fischman AJ, Bonab AA, Babich JW, Palmer EP, Alpert NM, Elmaleh DR, Callahan RJ, Barrow SA, Graham W, Meltzer PC, Hanson RN, Madras BK (1998) Rapid detection of Parkinson's disease by SPECT with altropine: a selective ligand for dopamine transporters. *Synapse* **29**: 128-141.

Goodman MM, Keil R, Shoup T, Eshima D, Eshima L, Kilts CD, Votaw J, Camp VM, Votaw D, Smith E, Kung MP, Malveaux E, Watts R, Huerkamp M, Garcia E, Hoffman JM (1997)

JPET #96792

Fluorine-18 labeled 2 $\beta$ -carbomethoxy-3 $\beta$ -(4-chlorophenyl)-8-(3-fluoropropyl)nortropane (FPCT): a PET radiotracer for imaging dopamine transporters. *J Nucl Med* **38**: 119-126.

Goodman MM, Kilts CD, Keil R, Shi B, Martarello L, Xing D, Votaw J, Ely TD, Lambert P, Owens MJ, Camp VM, Malveaux E, Hoffman JM (2000) <sup>18</sup>F-labeled FECNT: a selective radioligand for PET imaging of brain dopamine transporter. *Nucl Med Biol* **27**: 1-12.

Guilloteau D, Emond P, Baulieu J-L, Garreau L, Frangin Y, Pourcelot L, Mauclaire L, Besnard J-C, Chalon S (1998) Exploration of the dopamine transporter: in vitro and in vivo characterization of a high-affinity and high-specificity iodinated tropane derivative (E)-N-(3-iodoprop-2-enyl)-2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-methylphenyl)nortropane (PE2I). *Nucl Med Biol* **25**: 331-337.

Guttman M, Burkholder J, Kish SJ, Hussey D, Wilson A, DaSilva J, Houle S (1997) [<sup>11</sup>C]RTI-32 PET studies of the dopamine transporter in early dopa-naïve Parkinson's disease: implications for the symptomatic threshold. *Neurology* **48**: 1578-1583.

Haaparanta M, Bergman J, Laakso A, Hietala J, Solin O (1996) [<sup>18</sup>F]CFT ([<sup>18</sup>F]WIN 35,428), a radioligand to study the dopamine transporter with PET: biodistribution in rats. *Synapse* **23**: 321-327.

Hall H, Halldin C, Farde L, Sedvall G (1998) Whole hemisphere autoradiography of the postmortem human brain. *Nucl Med Biol* **25**: 715-719.

Hall H, Hurd Y, Pauli S, Halldin C, Sedvall G (2001) Human brain imaging post-mortem – whole hemisphere technologies. *International Review of Psychiatry on Research Methods and Biological Psychiatry, Special Issue*, **13**:12-17.

JPET #96792

Hall H, Halldin C, Guilloteau D, Chalon S, Emond P, Besnard JC, Farde L, Sedvall G (1999) Visualization of the dopamine transporter in the human brain post-mortem with the new selective ligand [<sup>125</sup>I]PE2I. *NeuroImage* **9**: 108-116

Halldin C, Erixon-Lindroth N, Pauli S, Chou YH, Okubo Y, Karlsson P, Lundkvist C, Olsson H, Guilloteau D, Emond P, Farde L (2003) [(11)C]PE2I: a highly selective radioligand for PET examination of the dopamine transporter in monkey and human brain. *Eur J Nucl Med Mol Imaging* **30**: 1220-1230.

Ilgin N, Zubieta J, Reich SG, Dannals RF, Ravert HT, Frost JJ (1999) PET imaging of the dopamine transporter in progressive supranuclear palsy and Parkinson's disease. *Neurology* **52**: 1221-1226.

Innis RB, Seibyl JP, Scanley BE, Laruelle M, Abi-Dargham A, Wallace E, Baldwin RM, Zea-Ponce Y, Zoghbi S, Wang S, Gao Y, Neumeyer JL, Charney DS, Hoffer PB, Marek KL (1993) Single photon emission computed tomographic imaging demonstrates loss of striatal dopamine transporters in Parkinson disease. *Proc Natl Acad Sci USA* **90**: 11965-11969.

Jucaite A, Fernell E, Halldin C, Forsberg H, Farde L (2005) Reduced midbrain dopamine transporter binding in male adolescent with attention-deficit/hyperactivity disorder: association between striatal dopamine markers and motor hyperactivity. *Biol Psychiatry* **57**: 229-238.

Kazumata K, Dhawan V, Chaly T, Antonini A, Margouleff C, Belakhlef A, Margouleff D, Neumeyer JL, Eidelberg D (1998) Dopamine transporter imaging with fluorine-18-FPCIT and PET. *J Nucl Med* **39**: 1521-1530.

Laakso A, Bergman J, Haaparanta M, Vilkmann H, Solin O, Hietala J (1998) [18F]CFT [(<sup>18</sup>F)WIN 35,428], a radioligand to study the dopamine transporter with PET: characterization in human subjects. *Synapse* **28**: 244-250.

Langer O, Någren K, Dollé F, Lundkvist C, Sandell J, Swahn C-G, Vaufrey F, Crouzel C, Mazière B, Halldin C (1999) Precursor synthesis and radiolabelling of the dopamine D<sub>2</sub>

JPET #96792

receptor ligand [<sup>11</sup>C]Raclopride from [<sup>11</sup>C]Methyl Triflate. *J Label Compounds Radiopharm* **42**: 1183-1193.

Leenders KL, Salmon EP, Tyrrell P, Perani D, Brooks DJ, Sager H, Jones T, Marsden CD, Frackowiak RS (1990) The nigrostriatal dopaminergic system assessed in vivo by positron emission tomography in healthy volunteer subjects and patients with Parkinson's disease. *Arch Neurol* **47**: 1290-1298.

Ma Y, Dhawan V, Mentis T, Chaly T, Spetsieris PG, Eidelberg D (2002) Parametric mapping of [<sup>18</sup>F]FPCIT binding in early stage Parkinson's disease: a PET study. *Synapse* **45**: 125-133.

Meltzer PC, Liang AY, Brownell A-L, Elmaleh DR, Madras BK (1993) Substituted 3-phenyltropane analogs of cocaine: synthesis, inhibition of binding at cocaine recognition sites, and positron emission tomography imaging. *J Med Chem* **36**: 855-862.

Mozley PD, Schneider JS, Acton PD, Plossl K, Stern MB, Siderowf A, Leopold NA, Li PY, Alavi A, Kung HF (2000) Binding of [<sup>99m</sup>Tc]TRODAT-1 to dopamine transporters in patients with Parkinson's disease and healthy volunteers. *J Nucl Med* **41**: 584-589.

Neumeyer JL, Tamahgnan G, Wang S, Gao Y, Milius RA, Kula NS, Baldessarini RJ (1996) N-substituted analogs of 2 $\beta$ -carbomethoxy-3 $\beta$ -(4'-iodophenyl)tropane ( $\beta$ -CIT) with selective affinity to dopamine or serotonin transporters in rat forebrain. *J Med Chem* **39**: 543-548.

Nurmi E, Ruottinen HM, Kaasinen V, Bergman J, Haaparanta M, Solin O, Rinne JO (2000) Progression in Parkinson's disease: a positron emission tomography study with a dopamine transporter ligand [<sup>18</sup>F]CFT. *Ann Neurol* **47**: 804-808.

Paxinos G, Watson C (1986) The rat brain in stereotaxic coordinates. Academic Press, New-York.

Pinborg LH, Videbaek C, Svarer C, Ynggaard S, Paulson OB, Knudsen GM (2002) Quantification of [<sup>123</sup>I]PE2I binding to dopamine transporters with SPET. *Eur J Nucl Med* **29**: 623-631.

JPET #96792

Sandell J, Langer O, Larsen P, Dollé F, Vaufrey F, Demphel S, Crouzel C, Halldin C (2000) Improved radioactivity of the PET radioligand [<sup>11</sup>C]FLB 457 by use of the GE medical systems PETtrace MeI MicroLab. *J Label Compounds Radiopharm* **43**: 331-338.

Sanders JD, Happe HK, Bylund DB, Murrin LC (2005) Development of the norepinephrine transporter in the rat CNS. *Neuroscience* **130**: 107-117.

Schwarz J, Linke R, Kerner M, Mozley PD, Trenkwalder C, Gasser T, Tatsch K (2000) Striatal dopamine transporter binding assessed by [I-123]IPT and single photon emission computed tomography in patients with early Parkinson's disease: implications for a preclinical diagnosis. *Arch Neurol* **57**: 205-208.

Seibyl JP, Marek K, Sheff K, Zoghbi S, Baldwin RM, Charney DS, van Dyck CH, Innis RB (1998) Iodine-123-beta-CIT and iodine-123-FPCIT SPECT measurement of dopamine transporters in healthy subjects and Parkinson's patients. *J Nucl Med* **39**: 1500-1508.

JPET #96792

### Footnotes

This work was supported by the French “Réseau National de Technologies pour la Santé” under the grant RNTS 03B243 FLUOPARK programme and by a grant from Biotechnocentre (Région Centre, France).

Partial results were presented as abstracts:

- Dollé F, Emond P, Saba W, Chalon S, Demphel S, Halldin C, Mavel S, Garreau L, Coulon C, Ottaviani M, Mazière B, Valette H, Bottlaender M, Guilloteau D. Radiosynthesis of [<sup>11</sup>C]LBT-999, a selective radioligand for the visualisation of the dopamine transporter with PET, *J. Label. Compounds Radiopharm.* 2003, *46*, S145.
- Hassoun W, Chalon S, Valette H, Dollé F, Garreau L, Emond P, Halldin C, Deloye J-B, Bottlaender M, Guilloteau D. [<sup>11</sup>C]LBT-999, a new radioligand to study the dopamine transporter with PET: Preclinical characterisation. *Eur. J. Nucl. Med.* 2004, *31*, 28, S208.

Correspondence to: Sylvie Chalon, Inserm U619; Université François-Rabelais de Tours, UFR Pharmacie, 31 avenue Monge, 37200, Tours, France

JPET #96792

### Legends for figures

**Fig.1.** Saturation experiments of [<sup>3</sup>H]LBT-999 binding in the rat striatum. Striatal membranes (30 μg protein/assay) were incubated with a constant concentration of [<sup>3</sup>H]LBT-999 (10 pmoles) and increasing concentrations of unlabeled LBT-999 (0.2 to 50 nM) in pH 7.4 Tris-HCl buffer for 90 min at 22°C. Nonspecific binding was determined in the presence of 1 μM PE2I.

A: one representative saturation curve. B: Scatchard transformation of the data.

**Fig.2.** Displacement of [<sup>3</sup>H]LBT-999 binding in rat striatum by various drugs. Striatal membranes were incubated with 10 pmoles of [<sup>3</sup>H]LBT-999 together with 10 nM of LBT-999 and increasing quantities of drugs (10<sup>-11</sup> to 10<sup>-5</sup> M).

Each curve is representative of 3 independent experiments, each performed in triplicate. The inhibitory constants of the drugs are given in Table 1.

**Fig.3.** Distribution of dopamine transporters in the human brain (horizontal sections) as visualized using [<sup>3</sup>H]LBT-999 (2 nM). Nonspecific binding was determined in the presence of 10 μM PE2I. Numbers in lower right corner represent brain number and distance from vertex in mm.

**Fig.4.** Time course of [<sup>11</sup>C]LBT-999 concentration in blood and cerebral areas (cerebellum, striatum, cortex) at 10, 30 and 60 min after i.v. injection in the rat. n = 4 per time.

**Fig.5.** Cerebral biodistribution of [<sup>11</sup>C]LBT-999 in rats: competition study.

Animals received a pre-injection of GBR 12909, paroxetine or nisoxetine at the dose of 5 mg/kg 3 min before [<sup>11</sup>C]LBT-999 i.v. injection and were sacrificed 30 min later. n = 6 per group. \*\*, p<0.01; \*, p<0.05.

**Fig.6.** Time-activity curves in various brain structures in baboon after intravenous injection of [<sup>11</sup>C]LBT-999.

JPET #96792

TABLE 1

**Inhibition of specific [<sup>3</sup>H]LBT-999 binding in rat striatal membranes**

A mixture of 10 pmoles of [<sup>3</sup>H]LBT-999 and 10 nM of unlabeled LBT-999 was incubated with 30 µg of protein in Tris-HCl buffer pH 7.4 for 90 min at 22°C in the presence of different drugs at concentrations of 10<sup>-11</sup> to 10<sup>-5</sup> M. Total binding was determined in the absence of drugs and nonspecific binding in the presence of 1 µM PE2I. IC<sub>50</sub> values were determined graphically. Values are mean ± SD of 3 independent experiments each performed in triplicate.

Drug	IC <sub>50</sub> (nM)
GBR 12935	2.4 ± 0.5
PE2I	18.0 ± 3.0
Paroxetine	> 1000
Citalopram	> 1000
MADAM	> 1000
Nisoxetine	> 1000
Desipramine	> 1000

Fig.1





Fig.2

## LBT 999 competition studies

% B





Fig.3

**Total binding**

Putamen  
Caudate

An autoradiograph of a brain slice showing total binding. The image displays a dark, dense region on the left side, which is labeled as the Putamen and Caudate. The rest of the brain slice shows a lighter, less dense appearance. The labels 'Putamen' and 'Caudate' are written in red text over the dark region.

HBB001R / 92.5

**+ PE2I (10  $\mu$ M)**

An autoradiograph of a brain slice showing total binding in the presence of 10 μM PE2I. The image displays a much lighter, less dense region on the left side compared to the total binding image, indicating a significant reduction in binding. The labels 'Putamen' and 'Caudate' are not visible in this image.

HBB001R / 92.6

Fig.4

**[11C]LBT-999**




Fig.5

[<sup>11</sup>C]LBT-999




Fig.6

