

Review Article

Role(s) of G3BPs in human pathogenesis

Chandrani Mukhopadhyay and Pengbo Zhou*

Department of Pathology and Laboratory Medicine, Weill Medical College of Cornell University, New York, NY 10065, USA.

Running Title: Role(s) of G3BPs in human pathogenesis

Corresponding author: Pengbo Zhou, Department of Pathology and Laboratory Medicine, Weill Medical College of Cornell University, New York, NY 10065, USA. e-mail address: pez2001@med.cornell.edu

Number of text pages: 23

Number of tables: 0

Number of figures: 3

Number of references: 163

Number of words in the Abstract: 130

Number of words in the Introduction: 164

Number of words in the Discussion: 6885

Abbreviations

G3BP1 [Ras-GTPase-activating protein (SH3 domain)-binding protein], SPOP (speckle type BTB/POZ protein), SG (Stress Granules)

Abstract

Ras-GTPase-activating protein (SH3 domain)-binding proteins (G3BP) are RNA binding proteins that plays a critical role in stress granule (SG) formation. SGs protect critical mRNAs from various environmental stress conditions by regulating mRNA stability and translation to maintain regulated gene expression. Recent evidence suggests that G3BPs can also regulate mRNA expression through interactions with RNA outside of SGs. G3BPs have been associated with a number of disease states, including cancer progression, invasion, metastasis, and viral infections, and may be useful as a cancer therapeutic target. This review summarizes the biology of G3BP including their structure, function, localization, role in cancer progression, virus replication, mRNA stability, and SGs formation. We will also discuss the potential of G3BPs as a therapeutic target.

KEYWORDS

G3BP1, G3BP2, cancer, SPOP, rasputin, stress granule, virus replication

Significance Statement

In this present review, we will discuss the molecular mechanism(s) and functional role(s) of G3BPs in the context of stress granule formation, interaction with viruses, stability of RNA, and tumorigenesis.

1. Introduction

G3BPs [also known as Rasputin (Rin) in *Drosophila*] are ubiquitously expressed 68 kDa proteins associated with RasGAP's SH3 domain, an essential Ras signaling component (Parker *et al*, 1996). Recent studies have raised doubts about the role of G3BP1 as a genuine binding partner of RasGAP (Annibaldi *et al*, 2011). Additional research is needed to confirm the

interplay between G3BPs and RasGAP. Moreover, G3BPs play a pivotal role in RNA processing, decay, and stress granule (SG) formation, and have been implicated in several disease pathologies such as neurodegenerative disorders, cancer advancement, and viral infections (Alam & Kennedy, 2019). Growing evidence indicates that G3BPs could be promising targets for cancer treatments (Annibaldi *et al.*, 2011; Dou *et al.*, 2016; Wei *et al.*, 2015). In this article, we will discuss the latest developments in the realm of G3BP research, encompassing the functions of G3BPs in cancer advancement, SG formation, RNA stabilization, translational regulation, and viral infections.

2. Structure and function of G3BPs in different cellular processes

In mammals, there are two homologs of G3BP. They are G3BP1 and G3BP2. G3BP2 exhibits two splice variants G3BP2a and G3BP2b. Distinct genes on human chromosomes 5 and 4 encode for G3BP1 and G3BP2, respectively. Structurally, all G3BPs exhibit a similar structure with distinct domains including a nuclear transport factor 2 (NTF2)-like domain at the N-terminus (Suyama *et al.*, 2000), an acidic and proline-rich region (PxxP) (Kang *et al.*, 2021), an RNA recognition motif (RRM) (Nagai *et al.*, 1995) and a C-terminal arginine and glycine rich (RGG) region (Burd & Dreyfuss, 1994) (Figure 1).

2.1 NTF2-like domain (amino acids 1-138): The NTF2-like domain of G3BPs is an important structural and functional element that contributes to the diverse functions of G3BPs including stress response, RNA metabolism, and nucleocytoplasmic transport. This domain exhibits both structural and functional similarity with the small NTF2 protein, which facilitates the transportation of molecules across the nuclear membrane *via* nuclear pores. (French *et al.*, 2002; Vognsen *et al.*, 2013). Both G3BP1 and G3BP2 are present in the nucleus during serum stimulation, supporting the role of the NTF2-like domain in nuclear transport (Barnes *et al.*, 2002). On the other hand, a separate research, conducted through targeted mutations in G3BP2 revealed that the NTF2-like domain primarily functions in guiding G3BP2 towards the

nuclear envelope rather than impacting its movement within the nucleus. (Prigent *et al*, 2000). In addition, it has been demonstrated that the NTF2-like domain aids in protein-protein interactions (Kennedy *et al*, 2001) such as interaction with Caprin1 (Reineke *et al*, 2015), which can mediate the dimerization of G3BPs (Tourriere *et al*, 2003), and also the assembly of SG (Yang *et al*, 2020a). NTF2-like domains also contribute to viral replication by binding to viral motifs and being enlisted by viral replication complexes (Schulte *et al*, 2016).

2.2 Acidic domain (amino acids 139-220): The acidic domain is a central region of G3BPs that is rich in negatively charged amino acid residues such as aspartic acid and glutamic acid. The domain is a crucial region that plays a key role in mediating protein-protein interactions and regulating various cellular processes, including SG formation. This region can also negatively affect the phase separation of membrane less compartments and regulate biochemical reactions. (Guillen-Boixet *et al*, 2020).

2.3 PxxP domain (amino acids 221-339): The PxxP domain is a proline-rich region found in many signaling proteins, including G3BPs, and is involved in protein-protein interactions and SG formation. Furthermore, this region is linked to protein interactions and has the ability to target SH3 domains through binding to aromatic amino acids (Booker *et al*, 1993). G3BP1's ability to interact with partner proteins is limited due to the existence of three PxxP motifs (Kay *et al*, 2000), in contrast to G3BP2a and G3BP2b, which possess five and six PxxP motifs respectively (Kennedy *et al.*, 2001). Moreover, according to Reineke and colleagues, the PxxP domain in G3BP is responsible for initiating eukaryotic initiation factor 2a (eIF2a) phosphorylation and SG formation. They also observed that this motif is essential for antiviral activity (Reineke *et al.*, 2015; Reineke & Lloyd, 2015).

2.4 RRM domain (amino acids 340-407): The RRM domain of G3BPs plays a critical role in mediating RNA binding and protein-protein interactions, which are essential for SG formation and other cellular processes involving RNA regulation. The region contains two conserved sequences, RNP1 and RNP2, that engage in interactions with target RNA sequences ranging

from 2-8 nucleotides by utilizing a binding platform formed by beta sheets (Kennedy *et al.*, 2001; Nagai *et al.*, 1995). This RNA recognition and binding domain regulates tumor-related mRNA stability. In addition to interacting with RNA, the RRM can also bind to other proteins (Clery *et al.*, 2008). The variation in RRM between G3BP1 and G3BP2 is the replacement of valines with isoleucines in the RNP-2 region (Kennedy *et al.*, 2001).

2.5 RGG boxes (amino acids 408-465): The RGG domain consists of clusters of arginine-glycine-glycine repeats. This structure enables post-transcriptional modifications and affects the interactions with proteins or RNA. Protein-mRNA interactions may be affected by methylation of arginine residues on RGG box of G3BPs and is important for several cellular process as discussed below. (1) When G3BP1 is methylated at Arg-433, it regulates the mRNA of Ctnnb1 (β -catenin) in a Wnt-dependent manner (Bikkavilli & Malbon, 2011). (2) The methylation of G3BP2 in a Wnt-dependent manner recruits methylated G3BP2 and leads to the formation of supermolecular complexes with disheveled protein 3 (Dvl3-complexes), thereby enabling the phosphorylation of lipoprotein receptor 6 (LRP6) at Ser-149 through GSK3 β (Glycogen Synthase Kinase 3 Beta) (Bikkavilli & Malbon, 2011). (3) G3BP2 methylation is also implicated in the regulation of β -catenin mRNA; however, it does not affect the protein (Bikkavilli & Malbon, 2011). (4) The oxidative stress triggered by sodium arsenite prompts the demethylation of G3BP1 at Arg-447, which regulates the assembly of SGs (Tsai *et al.*, 2016). (5) PRMT1 (Protein Arginine Methyltransferase 1) enters SGs and methylates G3BP1 at Arg-477 and stimulates SG disassembly whereas methylation of Arg-435 and Arg-460 by PRMT1 and PRMT5 has no effect during SGs formation (Tsai *et al.*, 2016). (6) During SG formation, Jumonji C domain-containing protein 6 (JMJD6), a constituent of SGs, interacts with G3BP1 and catalyzes its demethylation (Tsai *et al.*, 2016). (7) A recent report suggests that LRP6, a canonical Wnt receptor, suppresses the arginine methylation of several proteins, including G3BP1 (Ramachandran *et al.*, 2018). In addition, the RG-rich domain is also involved in nucleocytoplasmic shuttling (Isabelle *et al.*, 2012) and is essential for virus replication in the host (Gotte *et al.*, 2019).

3. Expression and subcellular localization of G3BPs

Although G3BPs are universally expressed in normal cells, there is some isoform-specific tissue restriction in their expression. For instance, G3BP1 is primarily expressed in the lung and kidney, G3BP2a in the brain, and G3BP2b in the small intestine (Kennedy *et al.*, 2001). Although G3BPs are mainly cytoplasmic proteins, they can translocate to the nucleus by virtue of the NTF2-like domain, which is responsible for their nuclear localization (Smith *et al.*, 1998). Costa and colleagues isolated G3BP1 from nuclear extract of HeLa cell as a functional DNA and RNA helicase indicating the nuclear localization of G3BP1 (Costa *et al.*, 1999). We have also observed, in multiple experiments and models, that G3BP1 is present in both the cytosol and nucleus, suggesting shuttling between these compartments (Mukhopadhyay *et al.*, 2021). As a result of its phosphorylation at Ser-149, G3BP1 can move to nuclei in quiescent cells (Gallouzi *et al.*, 1998; Tourriere *et al.*, 2001) whereas G3BP2 can move to nuclei in response to serum stimulation (French *et al.*, 2002).

4. Regulation of G3BPs at transcriptional, translational and post-translational levels

It is known that G3BP1 is subject to transcriptional, translational, and post-translational regulation. The regulation of G3BP1 has a wide range of downstream effects, including the assembly of SGs (Zhang *et al.*, 2019a) and the promotion of tumors (Alam & Kennedy, 2019). Studies investigating the transcriptional regulation of G3BP1 are limited. Using gene set enrichment analyses (GSEA), Ali and colleagues identified YY1 (Yin Yang 1 transcription factor), SYK (Spleen Associated Tyrosine Kinase), E2F-1 (E2F transcription factor 1), FOXA1 (Forkhead Box A1) and TGFBR2 (Transforming Growth Factor Beta Receptor 2) as activators of G3BP1 gene transcription, while SRF (Serum Response Factor), SIN3A (SIN3 Transcription

Regulator Family Member A), and AKT-1 (AKT Serine/Threonine Kinase 1) as suppressors (Ali *et al*, 2021).

Several proteins and miRs regulate G3BP1 at the post-transcriptional level. Many of these proteins interact with 5'-untranslated region (5'-UTR) (Hoftman *et al*, 2008; Lee *et al*, 2020; Somasekharan *et al*, 2015) of G3BP1, than with 3'-untranslated region (3'-UTR) (Deng *et al*, 2015; Liu *et al*, 2021; Lv *et al*, 2017). In contrast, many miRs have been identified to bind the 3'-UTR of G3BP1 and subsequently downregulate its expression. Y-box binding protein 1 (YB-1) is a DNA/RNA binding protein that binds to the 5'-UTR of G3BP1 transcripts and regulates translation (Somasekharan *et al.*, 2015). Thus, YB-1 regulates G3BP1 availability for SG assembly (Zhang *et al.*, 2019a) and there is a strong association between YB-1 and G3BP1 expression in human sarcomas (Somasekharan *et al.*, 2015). Lee and colleagues showed that Melanoma-associated antigen gene B2 (MAGE-B2), which is normally found in testicular tissue but is abnormally expressed in tumors (Hoftman *et al.*, 2008), can interact with the 5'-UTR of G3BP1. This interaction results in increased translation of G3BP1 (Lee *et al.*, 2020). The TAR DNA-binding protein 43 (TDP-43), which binds directly to highly conserved cis-regulatory elements found in the 3'-UTR of G3BP1, stabilizes the mRNA (Sidibe *et al*, 2021). Recent studies have identified G3BP1 interacting miRs, such as miR-193a-3p, miR-129-5p, miR-622, miR-132, and miR-362-3p (Deng *et al.*, 2015; Liu *et al.*, 2021; Lv *et al.*, 2017).

G3BP1 undergoes various post-translational modifications, including but not limited to methylation (Tsai *et al*, 2017), phosphorylation (Reineke *et al*, 2017), acetylation (Gal *et al*, 2019), polyADP-ribosylation (pADPr) (Leung *et al*, 2011), and ubiquitination (Gwon *et al*, 2021). These post-translational modifications of G3BP1 are often necessary for the disassembly of SGs rather than their assembly (Gwon *et al.*, 2021). One example is the modification of G3BP1 by the PRMT, arginine methyltransferase, which can methylate residues Arg-435, Arg-447, and Arg-460. (Tsai *et al.*, 2017). In the same vein, the acetylation of lysine 376 of G3BP1 by CBP/P300, which is a transcription co-factor for various nuclear proteins, results in the

disruption of RNA binding by G3BP1. This disruption leads to the disassembly of SGs (Gal *et al.*, 2019). Similarly, G3BP1 Ser-149 is phosphorylated by casein kinase 2 (CK2) both *in vivo* and *in vitro*, which inhibits SG assembly (Reineke *et al.*, 2017). As per Tsai *et al.*'s findings, JMJD6 augments the formation of SGs by reducing the monomethylation of G3BP1 and the asymmetric dimethylation of its three arginine residues (Tsai *et al.*, 2017). Moreover, Gal and colleagues confirmed that the same residue of G3BP1 is targeted for deacetylation by HDAC6 (histone deacetylase 6) as is targeted for acetylation by CBP/P300. G3BP1 is excessively acetylated when HDAC6 is lost or inhibited, resulting in impaired SG induction (Gal *et al.*, 2019). Additionally, poly-ubiquitinated G3BP1 at Lys-63 regulates the disassembly of SGs (Gwon *et al.*, 2021). Conversely, specific post-translational modifications of G3BP1 aid in the formation of SGs. One example of a post-translational modification that promotes the assembly of SGs is the methylation of G3BP1 by PRMT5, which preferentially targets Arg-460 to enhance G3BP1 signaling and SG formation (Tsai *et al.*, 2016). Just like the previous instances, the modification of G3BP1 by pADPr polymerase (PARP) can trigger the assembly of SGs (Leung *et al.*, 2011). In addition to its role in regulating SG assembly, post-translational modification of G3BP1 is also involved in a variety of other physiological functions. The PRMT1 mediates methylation of G3BP1 Arg-435, releases β -catenin mRNA and activates Wnt signaling (Bikkavilli & Malbon, 2011). Additionally, dexamethasone or ginsenoside-Rg1 activates the glucocorticoid receptor (GR), resulting in the rapid phosphorylation of G3BP1, facilitating its transfer from the cytoplasm into the nucleus (Kwok *et al.*, 2017).

Regulation of G3BP2 also occurs at the transcriptional, translational, and posttranslational modification levels. At transcriptional level, AR (androgen receptor) and FOXD1 (Human Forkhead-box D1) can bind to the G3BP2 promoter and modulate its transcription (Ashikari *et al.*, 2017; Hannenhalli & Kaestner, 2009; Heinlein & Chang, 2004; Kanda *et al.*, 2014; Lin *et al.*, 2020). At translational level, miRNAs targeting the 3'UTR of G3BP2 generally decrease its translation (Carney, 2016; Guo *et al.*, 2021; Zhao *et al.*, 2016; Zhao & He, 2021) while binding to

the 5'UTR of G3BP2 enhances its expression (Bezzi *et al*, 2017; Liu *et al*, 2018, 2020). There are various post-translational modifications of G3BP2. PRMTs target the RGG domain of G3BP2 in response to Wnt3a, leading to its methylation. The methylated form of G3BP2 serves as a positive regulator of Wnt signaling (Bikkavilli & Malbon, 2011; Jamieson *et al*, 2014; Katoh & Katoh, 2009; Li *et al*, 2012). The stability of G3BP2 protein is preserved through deubiquitination by the USP10 (Matsuki *et al*, 2013; Soncini *et al*, 2001; Takayama *et al*, 2018).

5. Roles of G3BPs in stress granule formation

When cells are exposed to cellular stresses such as oxidative stress, hypoxia, or viral infections, SG are formed in the cytoplasm. These SGs are primarily composed of complexes of mRNA and proteins that have been halted in their translation (Anderson *et al*, 2015; Buchan & Parker, 2009; White & Lloyd, 2012). Under stress conditions, SGs protect mRNA (Kedersha & Anderson, 2002) although their precise role in protein retrieval is not fully understood, studies have indicated that SGs also recruit proteins associated with diverse signaling pathways that can influence cellular metabolism and survival. Thus, SGs likely have additional functions beyond their role in mRNA storage and translational control. (Kim *et al*, 2005; Li *et al*, 2004). Under conditions of stress-induced eIF2a activation, both G3BP1 and G3BP2 localize within SGs (Tourriere *et al.*, 2003), but they are also capable of forming SGs independently (Matsuki *et al.*, 2013). G3BPs are phosphorylated under normal conditions and promote mRNA degradation. When cellular stress occurs, such as treatment with arsenite, unphosphorylated G3BPs oligomerize and deliver mRNA to SGs. When mammalian cells are treated with arsenite, G3BPs are found to be unphosphorylated at Ser-149, which results in the formation of SGs (Tourriere *et al.*, 2003) (Figure 2). The findings of these studies imply that the formation of SGs is a finely controlled process that is regulated in response to cellular stress. Besides stress-inducing agents, overexpression of several RNA-binding proteins such as TIA-1 (TIA1 Cytotoxic Granule Associated RNA Binding Protein), CPEB1 (Cytoplasmic Polyadenylation Element

Binding Protein 1) and G3BPs in combination with inhibition of translational initiation complex components (Dang *et al*, 2006; Mazroui *et al*, 2006) can induce the assembly of SGs (Gilks *et al*, 2004; Tourriere *et al.*, 2001; Wilczynska *et al*, 2005). Unlike other stress-inducing agents, the induction of SGs by G3BPs was found to be independent of eIF2 α phosphorylation but dependent on PKR-mediated phosphorylation of eIF2 α (Reineke *et al*, 2012). However, Taniuchi and colleagues demonstrated that the C-terminal region of G3BPs can induce phosphorylation of eIF2 α (Taniuchi *et al*, 2011a). Therefore, it remains to be elucidated if specific cellular contexts determine the dependence of eIF2 α phosphorylation by G3BPs. The formation of SGs in response to different stress stimuli is influenced by the interplay between G3BP's own transcript and/or protein and other proteins. McDonald and colleagues demonstrated that TDP-43 regulates SG by controlling the aggregation of TIA-1 and the mRNA levels of G3BP1. Specifically, when TDP-43 was silenced, it prevented the aggregation of TIA-1 and reduced the mRNA levels of G3BP1 (McDonald *et al*, 2011). Furthermore, YB-1 regulated the expression levels of G3BP1 through its interaction with the 5'-UTR of G3BP1, resulting in the formation of SGs during stress (Somasekharan *et al.*, 2015) (Figure 2a). Caprin1 can form a complex with G3BP1, and this complex is found in SGs (Figure 2a). However, it is not essential for the sequestration or formation of Caprin1 and G3BP1 into SGs, as they can still accumulate in SGs even in the absence of their association. (Solomon *et al*, 2007). However, Kedersha and colleagues reported that the interaction of G3BP1 with Caprin1 or USP10 (Ubiquitin Specific Peptidase 10) determined the fate of SGs as the former induced SGs formation while the later induced SGs inhibition (Kedersha *et al*, 2016). So far, based on published literature, it is considered that G3BPs are the basic components of SGs necessary for SG formation. Under stress conditions, G3BP2 may undergo dephosphorylation at Ser-149, leading to enhanced homo- and hetero-multimerization with G3BP1 and initiation of SG formation. The acidic region of both G3BP1 and G3BP2 has been shown to inhibit SG formation (Matsuki *et al.*, 2013;

Tourriere *et al.*, 2003). While G3BP2 can induce SG formation even in the absence of stress, its activity is lower compared to that of G3BP1 (Matsuki *et al.*, 2013).

It has recently become clearer how G3BPs contribute to the formation of SGs. It has been reported that SGs form in response to stress through liquid-liquid phase separation (LLPS), creating dynamic and membrane-less condensates (Maruri-Lopez *et al.*, 2021). Recently, Taylor and colleagues reported that NTF2 domain of G3BP1 dimerizes under normal conditions and generates a "locked conformation". However, under stressed conditions unfolded RNA interacts with the acidic and PxxP domains of G3BPs and generates an "open conformation" that allows interaction with other RNAs. This allows the G3BPs-RNA complex, which then recruits other stress proteins and forms a multimeric complex "SG" (Guillen-Boixet *et al.*, 2020; Sanders *et al.*, 2020; Yang *et al.*, 2020a) (Figure 2a). In line with this, according to Taylor and colleagues, the main structural difference between G3BP1 and G3BP2 is in their internal regions, which are not essential for SG formation (Yang *et al.*, 2020a). G3BP2 has predicted phosphorylation sites at Ser-149 and Ser-232, suggesting that it can also shift between open and closed conformational states like G3BP1. However, Franzmann and colleagues found that the phase separation properties of G3BP2 are slightly different from those of G3BP1. This group showed that G3BP2 has a higher tendency to phase separate with proteins and RNA compared to G3BP1 (Guillen-Boixet *et al.*, 2020). This may be due to G3BP2's longer intrinsically disordered regions and RNA-binding domain compared to G3BP1. Overall, it is not clear yet how these SGs contribute to disease progression, such as oncogenesis or drug resistance. Even though few articles mentioned SGs are involved in potential drug resistance mechanism (Gong *et al.*, 2013; Sabile *et al.*, 2013; Shi *et al.*, 2019; Somasekharan *et al.*, 2022), but future studies may explore detail mechanism within this perspective.

6. Contribution of G3BPs in cancer

G3BPs promotes tumor cell proliferation by attenuating the function of tumor suppressor genes. A number of human cancers overexpress G3BP1 and/or G3BP2, including breast cancer (Barnes *et al.*, 2002; Guitard *et al.*, 2001; Marx, 1989), pancreatic cancer (Taniuchi *et al.*, 2011b), colon, head and neck, lung (French *et al.*, 2002; Guitard *et al.*, 2001), and prostate cancer (Ashikari *et al.*, 2017; Mukhopadhyay *et al.*, 2021; Mukhopadhyay C, 2019; Takayama *et al.*, 2018).

6.1 G3BP1 driven molecular mechanism of cancer progression

Parker and colleagues reported that G3BP1 promotes tumor cell proliferation by activating the Ras signaling pathway (Parker *et al.*, 1996). However, a recent study found no direct interaction between G3BP1 and Ras-GAP, which suggests that G3BP1 may not dysregulate RAS signaling by directly targeting Ras-GAP (Annibaldi *et al.*, 2011). Numerous studies have indicated that G3BPs play a role in regulating cancer cells and act as accessory genes that support the survival of cancer cells Figure 2. Huang and colleagues observed that PTEN has the ability to downregulate the expression of G3BP1 protein *via* the PI3K pathway by virtue of its phosphatase activity (Huang *et al.*, 2005). Therefore, it is conceivable that loss of PTEN function resulting in the overexpression of G3BP1 leads to induce cellular proliferation and dysregulated promotion of the cell cycle (Figure 2a). Binding of G3BPs to the tumor suppressor p53 leads to its translocation from the nucleus to the cytoplasm, resulting in the inactivation of p53 (Figure 2a). This step is considered crucial in human tumorigenesis (Kim *et al.*, 2007). G3BP2 interacts with its negative regulator MDM2 (murine double minute 2), and exports p53 from the nucleus to the cytoplasm (Kim *et al.*, 2007). In a separate study, Mao and colleagues observed that a cytosolic long intervening/intergenic noncoding RNAs, P53RRA, functions as a tumor suppressor by activating the p53 pathway. Their findings showed that P53RRA interacts with the RRM domain of G3BP1 in the cytoplasm through nucleotides 1 and 871. This interaction displaced p53 from the G3BP1 complex, thereby keeping p53 in the nucleus, which led to cell cycle arrest and cell death (Mao *et al.*, 2018). Additionally, G3BP2 is involved in regulating the

translocation of p53 from the nucleus to the cytoplasm by interacting with the SUMO-E3 ligase RanBP2 when androgen is induced resulting in tumor formation (Ashikari *et al.*, 2017).

G3BP1 has been found to play a role in the progression of breast cancer and esophageal cancer by inhibiting the degradation of beta-catenin and enhancing its stability, thereby promoting cell proliferation (Zhang *et al.*, 2021a; Zhang *et al.*, 2019b). However, some studies suggest that G3BP1 actually has a negative regulatory effect on the Wnt/ β -catenin signaling pathway, which leads to a decrease in β -catenin levels and an increase in G3BP1 expression (Bikkavilli & Malbon, 2011). In addition, G3BP1 has been shown to bind to the tuberous sclerosis complex (TSC) and inhibit mTORC1 signaling, which can otherwise drive the growth and migration of breast cancer cells (Prentzell *et al.*, 2021).

G3BP1 has been shown to contribute to the proliferation, migration, and invasion of renal cell carcinoma (RCC), and knockdown of G3BP1 has been found to block the IL-6/STAT3 signaling pathway and reduce the metastatic ability of RCC (Wang *et al.*, 2018). In addition, G3BP1 has been found to interact with Y-box binding protein 1 (YBX1) to upregulate the expression of secreted phosphoprotein 1 (SPP1) and activate nuclear factor- κ B (NF- κ B), ultimately promoting RCC metastasis (Wang *et al.*, 2018). Moreover, G3BP1 has been implicated in promoting resistance to sunitinib in RCC (Sonnhof *et al.*, 1975). These findings suggest that targeting G3BP1 may be a promising therapeutic strategy for the treatment of RCC (Cho *et al.*, 2019).

In human lung cancer (H1299) cells, knockdown of G3BP has been found to inhibit the activation of Src, focal adhesion kinase (FAK), and ERK1/2, while also reducing the expression of NF- κ B. This leads to a decrease in the expression levels of matrix metalloproteinase-2 (MMP-2), matrix metalloproteinase-9 (MMP-9), and urokinase-type plasminogen activator (uPA), ultimately inhibiting the proliferation, invasion, and migration of lung cancer cells (Zhang *et al.*, 2013). Additionally, G3BP1 has been shown to be necessary for the activation of the senescence-associated secretory phenotype (SASP) (Omer *et al.*, 2020). The impact of SASP

on the proliferation and migration of cancer cells is complex and not well understood (Rao & Jackson, 2016).

According to reports, the strong cytoplasmic localization of p53 is associated with an increase in G3BP2 expression in prostate cancer (Ashikari *et al.*, 2017). This correlation predicts a poor prognosis and a higher likelihood of progression to hormone-refractory disease. These findings suggest that G3BP2 plays a role in the progression of prostate cancer (Ashikari *et al.*, 2017). A recent study showed that TRIM25 functions through interaction with G3BP2 to downregulate p53 activity resulting in prostate cancer cell proliferation (Takayama *et al.*, 2018). Moreover, Takayama and colleagues observed that the deubiquitinase USP10 regulates androgen-mediated signaling. It has been observed that the interaction between USP10 and G3BP2 leads to the suppression of p53 signaling, correlating to a poor prostate cancer prognosis. These findings emphasize the oncogenic function of USP10 in prostate cancer through its interaction with G3BP2 (Takayama *et al.*, 2018).

Recently, we observed that G3BP1 as an interactor and upstream regulator of CUL3^{SPOP}, functioning as an inhibitor of CUL3^{SPOP} ubiquitin ligase. This reveals a distinct function of G3BP1 that abrogates CUL3^{SPOP} tumor suppressive role of CUL3^{SPOP} and promotes prostate tumorigenesis (Mukhopadhyay *et al.*, 2021) (Figure 2a). Mechanistically, G3BP1 competitively inhibits substrate binding to SPOP and accelerates prostate epithelial cell proliferation, migration and invasion. As such, silencing of G3BP1 by CRISPER-Cas9 significantly reduced organoid formation of mPECs (murine prostate epithelial cells) and tumorigenicity in a prostate cell line-derived xenograft tumor model. Moreover, we showed that G3BP1 is a bona fide target of an AR that upregulates G3BP1 transcription and sets up a feed-forward loop of AR-mediated adverse signaling that further exacerbates PCa progression (Mukhopadhyay *et al.*, 2021; Mukhopadhyay C, 2019). This study shed light on new mode of SPOP inactivation other than SPOP mutation.

6.2 Role of G3BPs in cancer metastasis, invasion and Epithelial to mesenchymal transition (EMT) phenotype

The degradation of Binder of Arl Two (BART) transcripts is facilitated by G3BP1, while CD24 interacts with G3BP1 to inhibit its endoribonuclease activity on BART within SGs (Figure 2a). This mechanism prevents the metastasis and invasion of pancreatic cancer cells (Taniuchi *et al.*, 2011a). The dominant-negative overexpression of the N-terminal domain of G3BP1 suppresses the binding of G3BP1 and BART, thereby influencing the posttranscriptional control of cell invasiveness and metastasis in pancreatic cancer cells (Taniuchi *et al.*, 2011b). In addition, silencing of G3BP1 by RNAi led to a decrease in the spread and infiltration of lung cancer cells in humans. This was achieved by reducing the levels of Src, FAK, ERK, and NF- κ B through the MEK/ERK pathway, which in turn led to a decrease in the activation of matrix metalloprotease (MMP) 2 and 9, as well as plasminogen activator (uPA) (Zhang *et al.*, 2013) (Figure 2a). Therefore, silencing of G3BPs may attenuate the signaling cascades responsible for invasion and metastasis. In esophageal squamous cell carcinoma, Wang and colleagues observed that G3BP1 may have a crucial function in promoting lymph node metastasis and invasiveness. This is thought to occur through the activation of both the PI3K/AKT/mTOR and Wnt signaling pathways (Zhang *et al.*, 2019b). G3BP1 was found to inhibit lung metastasis by inhibiting SG formation in a lung cancer mouse model (Somasekharan *et al.*, 2015).

Epithelial to mesenchymal transition (EMT) is one of the hallmarks of cancer (Hanahan & Weinberg, 2011). The SMAD signaling pathway appears to be involved in the promotion of EMT in breast cancer cells as a result of G3BP1 overexpression (Zhang *et al.*, 2015). Knockdown of SMADs by siRNA completely inhibited G3BP1-induced EMT, indicating that the SMAD signaling pathway participates in this process (Figure 2a). Studies have indicated that reducing G3BP1 expression can prevent MDA-MB-231 cells from displaying mesenchymal characteristics *in vitro*, as well as impede tumor growth and metastasis in 4T1 cells *in vivo*. These findings suggest that targeting G3BP1 could be a promising therapeutic strategy for treating metastatic

breast cancer. G3BP1 activates EMT factors through phosphorylation of SMAD (Zhang *et al.*, 2015). In both *in vitro* and *in vivo* settings, researchers have demonstrated that the reduction of G3BP1 leads to decreased cell migration and metastasis in hepatocellular carcinomas (HCCs) (Gupta *et al.*, 2017). Research has indicated that G3BP2 plays a role in the initiation of breast cancer by stabilizing mRNA transcripts of the squamous cell carcinoma antigen recognized by T cells 3 (SART3). The SART3 gene increases the expression of the pluripotent transcription factors octamer binding protein 4 (Oct-4) and Nanog Homeobox (Nanog) (Gupta *et al.*, 2017) (Figure 2b). An anticancer compound, C108, interacts with G3BP2 *via* its RRM domain and inhibits tumor growth by degrading SART3 mRNA (Gupta *et al.*, 2017). According to a recent study, G3BP2 acts as both a positive regulator in the initiation of breast cancer and a negative regulator in its metastasis (Gupta *et al.*, 2017). One possible explanation for the opposing roles of G3BP2 is that cancer cells need to shed their EMT-phenotype to acquire cancer-initiating properties necessary for metastatic colonization (Ocana *et al.*, 2012). Recently, Shuang and colleagues observed that G3BP2 is significantly overexpressed in osteosarcoma cells and that FGA5-AS1 plays a role in promoting osteosarcoma cell proliferation and invasion by acting as a competing endogenous RNA (ceRNA) to sequester miR-124-3p and upregulate the expression of G3BP2 (Shuang *et al.*, 2022).

6.3 G3BP1 affects tumor growth by regulating mRNA stability

Controlling mRNA stability is crucial in regulating gene expression during cell proliferation, differentiation, and development. G3BPs can interact with RNA and determine its stability. It has been reported that G3BP1 interacts with c-Myc mRNA and, through its endoribonuclease activity, it cleaves the 3'-UTR of the c-Myc mRNA in a phosphorylation-dependent manner *in vitro*. However, this finding was not supported by other studies (Gallouzi *et al.*, 1998; Tourriere *et al.*, 2001; Zekri *et al.*, 2005) where they used MEF cells isolated from G3BP1 knock out mice. There is a possibility that G3BP2 is still present in these cells, and it is uncertain whether there are functional redundancies between these proteins in regulating c-Myc mRNA. Additionally, it is

possible that under these conditions, the endoribonuclease activity of G3BP1 is incapable of degrading short-lived mRNAs like the c-Myc transcript. Altogether, stability of c-Myc mRNA by G3BP1 is still ambiguous and requires further investigation. However, the endoribonuclease activity of G3BP1 is further supported by the stability of other transcripts such as BART (Taniuchi *et al.*, 2011a), CTNNB1 (Bikkavilli & Malbon, 2011), PMP22 (Winslow *et al.*, 2013), IGF-II and GAS5 (Zekri *et al.*, 2005) TAU (Atlas *et al.*, 2004) and CDK7 (Lypowy *et al.*, 2005) (Figure 2a). Ortega and colleagues reported that G3BP1 interacts with beta-F1 mRNA, mitochondrial H(+)-ATP synthase subunit and inhibits its translation, which may be involved in cancer progression (Ortega *et al.*, 2010). In a recent study by Fisher and colleagues, it was discovered that the RNA decay machinery proteins Up-frameshift 1 (UPF1) and G3BP1 play essential roles in the degradation of highly structured mRNAs with long 3'UTRs. (Fischer *et al.*, 2020). The regulation of mRNA transcription by G3BP1 is intricately controlled and relies on post-transcriptional modifications as well as its interaction with other proteins. For example, G3BP1 interacts with CDK7 and CDK9 mRNA transcripts in the presence of filamin for their stabilization (Lypowy *et al.*, 2005). G3BP1 exhibits its endoribonuclease activity in a phosphorylation-dependent manner (Irvine *et al.*, 2004). So far, there is only one report suggesting a role for G3BP2 in stabilizing mRNA. Gupta and colleagues observed that SART3 mRNA was stabilized by G3BP2 resulting in increased expression of the pluripotent transcription factors Oct-4 and Nanog which are involved in tumor initiation in breast cancer cells (Gupta *et al.*, 2017) (Figure 2b).

7. Contributions of G3BPs in other diseases

G3BP1's most acknowledged function is its contribution to the initiation of SGs. Additionally, it has been associated with neurodegenerative disease through various proposed mechanisms (Li *et al.*, 2013). Firstly, the failure of SGs to assemble correctly may lead to neurodegeneration

(Wolozin & Ivanov, 2019). Secondly, the persistence of SGs can lead to the development of cytoplasmic aggregates, a common feature observed in neurodegenerative disorders (Baron *et al*, 2013). Finally, the inability to disassemble SGs may sequester vital translation factors, transport proteins, or other critical cellular chaperones (Wolozin & Ivanov, 2019). Newly published articles have highlighted the significant role of G3BP1 in the regulation of cardiac hypertrophy, atrial fibrillation, and coronary heart disease (Dong *et al*, 2019; He *et al*, 2015; Xia *et al*, 2020). It has been reported that G3BP1/RIG-I/MAVS relay pathway is a component of the Wnt signaling pathway and targeting Wnt signaling may help reduce atherosclerosis, a disease of the arteries due to the deposition of fatty material on the inner walls (Ramachandran *et al*, 2018).

8. Roles of G3BPs in viral infection

The specific function of G3BPs in the context of viral infection has yet to be fully understood. Some studies have demonstrated that viruses exploit G3BPs to aid in their replication or transcription. On the other hand, other research has shown that G3BPs activate the innate immune system in response to viral infection to safeguard host cells against virus replication. In the following sections, each of these processes will be discussed (Figure 2a).

8.1 G3BPs promote virus replication

Vaccinia virus (VV), a DNA virus, utilizes G3BP1 to promote its replication in host cells (Katsafanas & Moss, 2004). During the initial stages of infection, the virus becomes enveloped by the rough endoplasmic reticulum, which attracts essential translation-initiating factors (including G3BP1 and Caprin-1). This enhances the activity of viral RNA polymerase and transcription factors, ultimately promoting viral replication (Katsafanas & Moss, 2004). G3BP1 also promotes the replication of different RNA viruses. Negative strand synthesis of the +SSRNA chikungunya viral genome is regulated by G3BP1 and/or G3BP2 (Scholte *et al*, 2015).

The study reveals that the interaction between G3BPs and nsP2 can facilitate CHIKV RNA replication and recover the infectivity of viruses that do not contain G3BP-binding motifs in the HVD of nsP3 (Wang & Merits, 2022). To promote the efficient translation of Sindbis virus (SINV) mRNA, G3BP1 interacts with both the nsP3 (non-structural protein 3) and the 40S ribosomal subunit (Gotte *et al.*, 2019). The RNA-binding protein farnesoid X receptor (FXR) from Eastern equine encephalitis virus (EEEV) and Venezuelan equine encephalitis virus (VEEV) interacts with G3BP1 and/or G3BP2 and nsP3 proteins to construct the viral replication complex (vRC) (Frolov *et al.*, 2017; Kim *et al.*, 2016). The binding sites of the G3BP and FXR protein families in the nsP3 HVD (hypervariable Domain) of EEEV are deleted, the virus becomes unable to cause disease in mice (Meshram *et al.*, 2020). Hepatitis C virus (HCV), dengue virus (DENV), rubella virus (RUBV), and Newcastle disease virus (NDV) interact with G3BP1 and/or G3BP2 to facilitate either their replication or their transcription (Bidet *et al.*, 2017; Matthews & Frey, 2012; Sun *et al.*, 2017; Xia *et al.*, 2015; Yi *et al.*, 2006). The bursal disease virus (IBDV) exploits G3BPs to trigger the development of multiple SGs, facilitating the virus's replication within the host cell (Zhao *et al.*, 2020). Based on published reports, roles of SGs differs in virus-infected cells. As an illustration, the formation of SG enhances viral replication in the case of respiratory syncytial virus (RSV) infection (Lindquist *et al.*, 2010). However, the assembly of SGs inhibits Zika virus (ZIKV) infection (Hou *et al.*, 2017), while the creation of SGs has no impact on viral replication during rubella virus (RUBV) infection (Matthews & Frey, 2012). Recently, it has been reported that G3BP1 and G3BP2 promote the viral growth of Mayaro virus (MAYV), which is belong to the group of Old World alphaviruses (Neyret *et al.*, 2022).

The current pandemic of COVID-19 (Coronavirus Disease-2019), a respiratory disease, has led to over millions of deaths in over 188 countries. COVID-19 is caused by a novel virus strain, SARS-CoV-2, an enveloped, positive-sense, single-stranded RNA beta-coronavirus of the family Coronaviridae (Wang *et al.*, 2020; Zhu *et al.*, 2020). In 2020, the interactome study by Gordon *et al.*, identified SARS-CoV-2 nucleocapsid (N) protein, which is critically involved in

viral RNA replication, binds to G3BP1, which is later supported by many other groups (Biswal *et al*, 2022; Gordon *et al*, 2020; Liu *et al*, 2022). Recently, it has been reported that SG assembly, which is a defense mechanism and protects cells in any stress condition (Dolliver *et al*, 2022; Liu *et al*, 2022), was significantly reduced upon SARS-CoV-2 infection (Liu *et al*, 2022). They also observed that N protein of SARS-CoV-2 inhibit G3BP1-RIG-I-mediated interferon (IFN) production to facilitate viral replication (Liu *et al*, 2022). According to these studies, G3BP1 can turn as a "promiscuous" regulator of SARS-CoV-2 infection and thus targeting G3BP1 may be used as a new therapeutic strategy for SARS-CoV-2 virus infection.

8.2 G3BPs inhibit virus replication

Both DNA and RNA virus replications are inhibited by G3BPs. Liu and colleagues reported that G3BP1 facilitated the binding between cGAS (Cyclic GMP-AMP Synthase) and DNA to form a large cGAS complex that induced cGAS-dependent IFN production. Therefore, G3BP1 plays a critical role in combating DNA virus infections by engaging in the cGAS-mediated antiviral response (Liu *et al*, 2019). G3BPs also antagonize RNA virus growth. For instance, G3BP interacts with various viral proteins of porcine epidemic diarrhea virus (PEDV), foot-and-mouth disease virus (FMDV), mammalian reovirus (MRV), and porcine reproductive and respiratory syndrome virus (PRRSV), thus inhibiting their propagation in the host cell (Catanzaro & Meng, 2019; Choudhury *et al*, 2017; Galan *et al*, 2017; Pandey *et al*, 2019). It has also been reported that FMDV suppresses the host's stress and antiviral responses, and highlights the significance of G3BP1 dephosphorylation and its breakdown by viral protease in hindering the host's ability to defend against FMDV infection (Ye *et al*, 2018). G3BP1 is an antiviral protein (Reineke & Lloyd, 2015), as such, viral proteins cleave G3BP to initiate antiviral responses. For an example, 3Cpro of coxsackie virus type B3 (CVB3) (Fung *et al*, 2013), human enterovirus D68 (EV-D68)(Cheng *et al*, 2020) and poliovirus (PV) can cleave G3BP1 at Gln-325, resulting in the disassembly of SGs, which aids in viral translation. G3BP1 promotes the expression of RIG-I and positively regulates type-1 interferon (IFN1) response to generate antiviral responses (Yang

et al, 2020b). Furthermore, G3BP1 facilitates the generation of cytokines through activation of the NF- κ B and c-Jun N terminal kinase (JNK) signaling pathways (Reineke & Lloyd, 2015), to generate an antiviral response (Wen *et al*, 2020; Zhou *et al*, 2021). G3BP inhibits the replication of retroviruses. As an illustration, G3BP1 seizes HIV-1 RNA transcripts, which restricts viral protein production, mRNA translation, and virus particle formation (Cobos Jimenez *et al*, 2015). Nevertheless, the capsid domain of HIV-1 Gag protein can also interact with G3BP1, leading to the disassembly of preformed SGs by replacing eEF2. This response may facilitate virus replication (Valiente-Echeverria *et al*, 2014).

9. G3BPs as potential therapeutic targets

There is increasing evidence that G3BPs plays a role in oncogenesis, autoimmune diseases, and viral infection. In the following sections, we summarize the latest articles describing G3BPs as a potential therapeutic targets.

9.1 Cancer-related therapeutic treatments

Currently a combination of resveratrol, epigallocatechin gallate (EGCG) and GAP161 has been shown to target G3BP1 and inhibit its carcinogenic effects (Oi *et al*, 2015; Shim *et al*, 2010; Zhang *et al.*, 2013). By interacting with USP10, a deubiquitinating enzyme of p53, G3BP1 negatively regulates the expression of p53, thus controlling melanoma cell proliferation and colon cancer cell proliferation (Oi *et al.*, 2015). Resveratrol directly binds to G3BP1, which leads to the disruption of G3BP1/USP10 interaction. As a result, USP10 is released, allowing for the deubiquitination of p53 and triggering of apoptosis in tumor cells (Oi *et al.*, 2015). Therefore, resveratrol, which is found in natural products, is considered a promising anticancer agent that specifically targets G3BP1.

EGCG, a polyphenol compounds present in green tea, was found to interact with G3BP1 with high binding affinity and thus inhibiting MAPK-mediated signaling to suppress growth of H1299

and CL13 lung cancer cells, where G3BP1 expression is high (Shim *et al.*, 2010). Cho and colleagues reported that lung cancer cell lines become more susceptible to radiation therapy when G3BP1 is depleted. This impact is associated with the production of reactive oxygen species (ROS) (Cho *et al.*, 2019). In colon cancer cells, a novel peptide, GAP161, inhibited MAPK signaling by interfering with the G3BP1 and RasGAP interaction to suppress Ras signaling pathways (Zhang *et al.*, 2012). Furthermore, reducing the levels of G3BP enhances the effectiveness of cisplatin-induced apoptosis and growth inhibition in HCT116 cells (Zhang *et al.*, 2012), indicating that GAP161 could be a promising therapeutic agent for colon cancer. G3BP1's ability to regulate the assembly of SGs in cancer cells can affect the efficacy of commonly used clinical drugs like Bortezomib. Bortezomib-induced SG assembly can lead to resistance to Bortezomib-mediated apoptosis, highlighting the importance of G3BP1 in cancer progression (Fournier *et al.*, 2010). Bittencourt and colleagues found that G3BP1 knockdown in U87 glioblastoma cells diminished SG formation and stimulated Bortezomib-induced apoptosis *in vitro*, in addition to inhibiting glioblastoma-induced angiogenesis *in vivo* (Bittencourt *et al.*, 2019).

There are few reports on the therapeutic treatment of G3BP2 alone. As G3BP1 and G3BP2 share common domains and most of the drugs target the NTF2 domain of G3BPs, which is similar in both G3BP1 and G3BP2. As Such, compounds/drugs used to treat G3BP1 can also be used to treat G3BP2. In the following section we discuss the therapeutic treatment options for G3BP2. There are two peptides (P109 and P110) that can inhibit the interaction between NTF2 domain of G3BPs and Ras-GAP and biologically involve in cisplatin-induced cellular death (Cui *et al.*, 2010). Additionally, it has been reported that compound C108 inhibits G3BP2, delays stress response, suppresses cancer cell metastasis, and promotes the proliferation and infiltration of CD8 T-cells (Gupta *et al.*, 2017; Zhang *et al.*, 2021b; Zheng *et al.*, 2022).

9.2 Therapeutic treatments for other diseases

G3BP1's participation in signaling pathways associated with innate immunity indicates its potential involvement in autoimmune diseases. G3BP1 is recognized for its involvement in the identification of foreign DNA by cGAS. However, when G3BP1 is excessively activated, it can trigger the overproduction of IFN *via* cGAS, which can result in autoimmune diseases (Liu *et al.*, 2019). Efficient activation of cGAS, a key component of DNA sensing, relies on the critical involvement of G3BP1. G3BP1 assists in the formation of significantly large cGAS complexes, which amplifies cGAS' capability to bind to DNA. Conversely, deficiency in G3BP1 leads to inefficient DNA binding by cGAS, resulting in reduced cGAS-dependent IFN production. In this regard, EGCG has been discovered to interrupt the G3BP1-cGAS complex, thereby obstructing DNA-triggered activation of cGAS. This leads to the inhibition of IFN production both *in vivo* and *in vitro* (Liu *et al.*, 2019). Resveratrol inhibits the production of IFN triggered by intracellular DNA and RNA by targeting G3BP1 (Cai *et al.*, 2021). Given their ability to disrupt the G3BP1-cGAS complex and inhibit IFN production, both EGCG and resveratrol hold promise as potential treatments for autoimmune diseases related to cGAS or caused by abnormal nucleic acid sensing.

10. Concluding Remarks and Further Perspectives

G3BPs have emerged as important molecules in various pathological conditions like oncogenesis and viral infection. Current evidence suggest that G3BPs are highly expressed in cancer, and thus, could be extremely useful in either detection or prognosis of cancer patients. Furthermore, G3BPs, particularly G3BP1, have emerged as promising targets for drug development and cancer therapy. G3BPs engage in numerous interactions with various proteins to regulate a wide range of biological processes. Targeting these protein interactions may present a promising opportunity for developing novel therapeutic strategies in various disease states. While the crystal structure of G3BP1's NTF2 domain has been determined, the complete molecular architecture of G3BPs remains elusive. Given the broad functional involvement of

different domains within G3BPs across various biological processes, it is imperative that future investigations prioritize unraveling the molecular structure of G3BPs to gain insights into their protein interactions at the molecular level. A thorough understanding of G3BPs' structural features will be critical in the rational design of drugs or inhibitors that can specifically target their protein interactions. This can be used to treat not only cancer patients but also reduce viral infection and neurological diseases.

Author Contributions

Wrote and contributed to the writing of the manuscript: Mukhopadhyay, C. and Zhou, P

References

- Alam U, Kennedy D (2019) Rasputin a decade on and more promiscuous than ever? A review of G3BPs. *Biochim Biophys Acta Mol Cell Res* 1866: 360-370
- Ali N, Prasad K, AlAsmari AF, Alharbi M, Rashid S, Kumar V (2021) Genomics-guided targeting of stress granule proteins G3BP1/2 to inhibit SARS-CoV-2 propagation. *Int J Biol Macromol* 190: 636-648
- Anderson P, Kedersha N, Ivanov P (2015) Stress granules, P-bodies and cancer. *Biochim Biophys Acta* 1849: 861-870
- Annibaldi A, Dousse A, Martin S, Tazi J, Widmann C (2011) Revisiting G3BP1 as a RasGAP binding protein: sensitization of tumor cells to chemotherapy by the RasGAP 317-326 sequence does not involve G3BP1. *PLoS One* 6: e29024
- Ashikari D, Takayama K, Tanaka T, Suzuki Y, Obinata D, Fujimura T, Urano T, Takahashi S, Inoue S (2017) Androgen induces G3BP2 and SUMO-mediated p53 nuclear export in prostate cancer. *Oncogene* 36: 6272-6281

- Atlas R, Behar L, Elliott E, Ginzburg I (2004) The insulin-like growth factor mRNA binding-protein IMP-1 and the Ras-regulatory protein G3BP associate with tau mRNA and HuD protein in differentiated P19 neuronal cells. *J Neurochem* 89: 613-626
- Barnes CJ, Li F, Mandal M, Yang Z, Sahin AA, Kumar R (2002) Heregulin induces expression, ATPase activity, and nuclear localization of G3BP, a Ras signaling component, in human breast tumors. *Cancer Res* 62: 1251-1255
- Baron DM, Kaushansky LJ, Ward CL, Sama RR, Chian RJ, Boggio KJ, Quaresma AJ, Nickerson JA, Bosco DA (2013) Amyotrophic lateral sclerosis-linked FUS/TLS alters stress granule assembly and dynamics. *Mol Neurodegener* 8: 30
- Bezzi M, Guarnerio J, Pandolfi PP (2017) A circular twist on microRNA regulation. *Cell Res* 27: 1401-1402
- Bidet K, Dadlani D, Garcia-Blanco MA (2017) Correction: G3BP1, G3BP2 and CAPRIN1 Are Required for Translation of Interferon Stimulated mRNAs and Are Targeted by a Dengue Virus Non-coding RNA. *PLoS Pathog* 13: e1006295
- Bikkavilli RK, Malbon CC (2011) Arginine methylation of G3BP1 in response to Wnt3a regulates beta-catenin mRNA. *J Cell Sci* 124: 2310-2320
- Biswal M, Lu J, Song J (2022) SARS-CoV-2 Nucleocapsid Protein Targets a Conserved Surface Groove of the NTF2-like Domain of G3BP1. *J Mol Biol* 434: 167516
- Bittencourt LFF, Negreiros-Lima GL, Sousa LP, Silva AG, Souza IBS, Ribeiro R, Dutra MF, Silva RF, Dias ACF, Soriani FM *et al* (2019) G3BP1 knockdown sensitizes U87 glioblastoma cell line to Bortezomib by inhibiting stress granules assembly and potentializing apoptosis. *J Neurooncol* 144: 463-473
- Booker GW, Gout I, Downing AK, Driscoll PC, Boyd J, Waterfield MD, Campbell ID (1993) Solution structure and ligand-binding site of the SH3 domain of the p85 alpha subunit of phosphatidylinositol 3-kinase. *Cell* 73: 813-822
- Buchan JR, Parker R (2009) Eukaryotic stress granules: the ins and outs of translation. *Mol Cell* 36: 932-941

- Burd CG, Dreyfuss G (1994) Conserved structures and diversity of functions of RNA-binding proteins. *Science* 265: 615-621
- Cai H, Liu X, Zhang F, Han QY, Liu ZS, Xue W, Guo ZL, Zhao JM, Sun LM, Wang N *et al* (2021) G3BP1 Inhibition Alleviates Intracellular Nucleic Acid-Induced Autoimmune Responses. *J Immunol* 206: 2453-2467
- Carney EF (2016) Diabetic nephropathy: MiR-23b protects against fibrosis in diabetic nephropathy. *Nat Rev Nephrol* 12: 197
- Catanzaro N, Meng XJ (2019) Porcine reproductive and respiratory syndrome virus (PRRSV)-induced stress granules are associated with viral replication complexes and suppression of host translation. *Virus Res* 265: 47-56
- Cheng J, Gao S, Zhu C, Liu S, Li J, Kang J, Wang Z, Wang T (2020) Typical Stress Granule Proteins Interact with the 3' Untranslated Region of Enterovirus D68 To Inhibit Viral Replication. *J Virol* 94
- Cho E, Than TT, Kim SH, Park ER, Kim MY, Lee KH, Shin HJ (2019) G3BP1 Depletion Increases Radiosensitisation by Inducing Oxidative Stress in Response to DNA Damage. *Anticancer Res* 39: 6087-6095
- Choudhury P, Bussiere LD, Miller CL (2017) Mammalian Orthoreovirus Factories Modulate Stress Granule Protein Localization by Interaction with G3BP1. *J Virol* 91
- Clery A, Blatter M, Allain FH (2008) RNA recognition motifs: boring? Not quite. *Curr Opin Struct Biol* 18: 290-298
- Cobos Jimenez V, Martinez FO, Booiman T, van Dort KA, van de Klundert MA, Gordon S, Geijtenbeek TB, Kootstra NA (2015) G3BP1 restricts HIV-1 replication in macrophages and T-cells by sequestering viral RNA. *Virology* 486: 94-104
- Costa M, Ochem A, Staub A, Falaschi A (1999) Human DNA helicase VIII: a DNA and RNA helicase corresponding to the G3BP protein, an element of the ras transduction pathway. *Nucleic Acids Res* 27: 817-821
- Cui W, Wei Z, Chen Q, Cheng Y, Geng L, Zhang J, Chen J, Hou T, Ji M (2010) Structure-based design of peptides against G3BP with cytotoxicity on tumor cells. *J Chem Inf Model* 50: 380-387

- Dang Y, Kedersha N, Low WK, Romo D, Gorospe M, Kaufman R, Anderson P, Liu JO (2006) Eukaryotic initiation factor 2alpha-independent pathway of stress granule induction by the natural product pateamine A. *J Biol Chem* 281: 32870-32878
- Deng W, Yan M, Yu T, Ge H, Lin H, Li J, Liu Y, Geng Q, Zhu M, Liu L *et al* (2015) Quantitative proteomic analysis of the metastasis-inhibitory mechanism of miR-193a-3p in non-small cell lung cancer. *Cell Physiol Biochem* 35: 1677-1688
- Dolliver SM, Kleer M, Bui-Marinis MP, Ying S, Corcoran JA, Khapersky DA (2022) Nsp1 proteins of human coronaviruses HCoV-OC43 and SARS-CoV2 inhibit stress granule formation. *PLoS Pathog* 18: e1011041
- Dong G, Liang F, Sun B, Wang C, Liu Y, Guan X, Yang B, Xiu C, Yang N, Liu F *et al* (2019) Presence and function of stress granules in atrial fibrillation. *PLoS One* 14: e0213769
- Dou N, Chen J, Yu S, Gao Y, Li Y (2016) G3BP1 contributes to tumor metastasis via upregulation of Slug expression in hepatocellular carcinoma. *Am J Cancer Res* 6: 2641-2650
- Fischer JW, Busa VF, Shao Y, Leung AKL (2020) Structure-Mediated RNA Decay by UPF1 and G3BP1. *Mol Cell* 78: 70-84 e76
- Fournier MJ, Gareau C, Mazroui R (2010) The chemotherapeutic agent bortezomib induces the formation of stress granules. *Cancer Cell Int* 10: 12
- French J, Stirling R, Walsh M, Kennedy HD (2002) The expression of Ras-GTPase activating protein SH3 domain-binding proteins, G3BPs, in human breast cancers. *Histochem J* 34: 223-231
- Frolov I, Kim DY, Akhrymuk M, Mobley JA, Frolova EI (2017) Hypervariable Domain of Eastern Equine Encephalitis Virus nsP3 Redundantly Utilizes Multiple Cellular Proteins for Replication Complex Assembly. *J Virol* 91
- Fung G, Ng CS, Zhang J, Shi J, Wong J, Piesik P, Han L, Chu F, Jagdeo J, Jan E *et al* (2013) Production of a dominant-negative fragment due to G3BP1 cleavage contributes to the disruption of mitochondria-associated protective stress granules during CVB3 infection. *PLoS One* 8: e79546
- Gal J, Chen J, Na DY, Tichacek L, Barnett KR, Zhu H (2019) The Acetylation of Lysine-376 of G3BP1 Regulates RNA Binding and Stress Granule Dynamics. *Mol Cell Biol* 39

- Galan A, Lozano G, Pineiro D, Martinez-Salas E (2017) G3BP1 interacts directly with the FMDV IRES and negatively regulates translation. *FEBS J* 284: 3202-3217
- Gallouzi IE, Parker F, Chebli K, Maurier F, Labourier E, Barlat I, Capony JP, Tocque B, Tazi J (1998) A novel phosphorylation-dependent RNase activity of GAP-SH3 binding protein: a potential link between signal transduction and RNA stability. *Mol Cell Biol* 18: 3956-3965
- Gilks N, Kedersha N, Ayodele M, Shen L, Stoecklin G, Dember LM, Anderson P (2004) Stress granule assembly is mediated by prion-like aggregation of TIA-1. *Mol Biol Cell* 15: 5383-5398
- Gong B, Hu H, Chen J, Cao S, Yu J, Xue J, Chen F, Cai Y, He H, Zhang L (2013) Caprin-1 is a novel microRNA-223 target for regulating the proliferation and invasion of human breast cancer cells. *Biomed Pharmacother* 67: 629-636
- Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, O'Meara MJ, Rezelj VV, Guo JZ, Swaney DL *et al* (2020) A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature*
- Gotte B, Panas MD, Hellstrom K, Liu L, Samreen B, Larsson O, Ahola T, McInerney GM (2019) Separate domains of G3BP promote efficient clustering of alphavirus replication complexes and recruitment of the translation initiation machinery. *PLoS Pathog* 15: e1007842
- Guillen-Boixet J, Kopach A, Holehouse AS, Wittmann S, Jahnel M, Schlussler R, Kim K, Trussina I, Wang J, Mateju D *et al* (2020) RNA-Induced Conformational Switching and Clustering of G3BP Drive Stress Granule Assembly by Condensation. *Cell* 181: 346-361 e317
- Guitard E, Parker F, Millon R, Abecassis J, Tocque B (2001) G3BP is overexpressed in human tumors and promotes S phase entry. *Cancer Lett* 162: 213-221
- Guo YX, Wang N, Wu WC, Li CQ, Chen RH, Zhang Y, Li X (2021) The Role of miR-23b in Cancer and Autoimmune Disease. *J Oncol* 2021: 6473038
- Gupta N, Badeaux M, Liu Y, Naxerova K, Sgroi D, Munn LL, Jain RK, Garkavtsev I (2017) Stress granule-associated protein G3BP2 regulates breast tumor initiation. *Proc Natl Acad Sci U S A* 114: 1033-1038
- Gwon Y, Maxwell BA, Kolaitis RM, Zhang P, Kim HJ, Taylor JP (2021) Ubiquitination of G3BP1 mediates stress granule disassembly in a context-specific manner. *Science* 372: eabf6548

- Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* 144: 646-674
- Hannenhalli S, Kaestner KH (2009) The evolution of Fox genes and their role in development and disease. *Nat Rev Genet* 10: 233-240
- He M, Yang Z, Abdellatif M, Sayed D (2015) GTPase Activating Protein (Sh3 Domain) Binding Protein 1 Regulates the Processing of MicroRNA-1 during Cardiac Hypertrophy. *PLoS One* 10: e0145112
- Heinlein CA, Chang C (2004) Androgen receptor in prostate cancer. *Endocr Rev* 25: 276-308
- Hoftman AD, Tai LQ, Tze S, Seligson D, Gatti RA, McCurdy DK (2008) MAGE-B2 autoantibody: a new biomarker for pediatric systemic lupus erythematosus. *J Rheumatol* 35: 2430-2438
- Hou S, Kumar A, Xu Z, Airo AM, Stryapunina I, Wong CP, Branton W, Tchesnokov E, Gotte M, Power C *et al* (2017) Zika Virus Hijacks Stress Granule Proteins and Modulates the Host Stress Response. *J Virol* 91
- Huang Y, Wernyj RP, Norton DD, Precht P, Seminario MC, Wange RL (2005) Modulation of specific protein expression levels by PTEN: identification of AKAP121, DHFR, G3BP, Rap1, and RCC1 as potential targets of PTEN. *Oncogene* 24: 3819-3829
- Irvine K, Stirling R, Hume D, Kennedy D (2004) Rasputin, more promiscuous than ever: a review of G3BP. *Int J Dev Biol* 48: 1065-1077
- Isabelle M, Gagne JP, Gallouzi IE, Poirier GG (2012) Quantitative proteomics and dynamic imaging reveal that G3BP-mediated stress granule assembly is poly(ADP-ribose)-dependent following exposure to MNNG-induced DNA alkylation. *J Cell Sci* 125: 4555-4566
- Jamieson C, Sharma M, Henderson BR (2014) Targeting the beta-catenin nuclear transport pathway in cancer. *Semin Cancer Biol* 27: 20-29
- Kanda T, Jiang X, Yokosuka O (2014) Androgen receptor signaling in hepatocellular carcinoma and pancreatic cancers. *World J Gastroenterol* 20: 9229-9236
- Kang W, Wang Y, Yang W, Zhang J, Zheng H, Li D (2021) Research Progress on the Structure and Function of G3BP. *Front Immunol* 12: 718548
- Kato M, Kato M (2009) Transcriptional mechanisms of WNT5A based on NF-kappaB, Hedgehog, TGFbeta, and Notch signaling cascades. *Int J Mol Med* 23: 763-769

- Katsafanas GC, Moss B (2004) Vaccinia virus intermediate stage transcription is complemented by Ras-GTPase-activating protein SH3 domain-binding protein (G3BP) and cytoplasmic activation/proliferation-associated protein (p137) individually or as a heterodimer. *J Biol Chem* 279: 52210-52217
- Kay BK, Williamson MP, Sudol M (2000) The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *FASEB J* 14: 231-241
- Kedersha N, Anderson P (2002) Stress granules: sites of mRNA triage that regulate mRNA stability and translatability. *Biochem Soc Trans* 30: 963-969
- Kedersha N, Panas MD, Achorn CA, Lyons S, Tisdale S, Hickman T, Thomas M, Lieberman J, McInerney GM, Ivanov P *et al* (2016) G3BP-Caprin1-USP10 complexes mediate stress granule condensation and associate with 40S subunits. *J Cell Biol* 212: 845-860
- Kennedy D, French J, Guitard E, Ru K, Tocque B, Mattick J (2001) Characterization of G3BPs: tissue specific expression, chromosomal localisation and rasGAP(120) binding studies. *J Cell Biochem* 84: 173-187
- Kim DY, Reynaud JM, Rasaloukaya A, Akhrymuk I, Mobley JA, Frolov I, Frolova EI (2016) New World and Old World Alphaviruses Have Evolved to Exploit Different Components of Stress Granules, FXR and G3BP Proteins, for Assembly of Viral Replication Complexes. *PLoS Pathog* 12: e1005810
- Kim MM, Wiederschain D, Kennedy D, Hansen E, Yuan ZM (2007) Modulation of p53 and MDM2 activity by novel interaction with Ras-GAP binding proteins (G3BP). *Oncogene* 26: 4209-4215
- Kim WJ, Back SH, Kim V, Ryu I, Jang SK (2005) Sequestration of TRAF2 into stress granules interrupts tumor necrosis factor signaling under stress conditions. *Mol Cell Biol* 25: 2450-2462
- Kwok HH, Poon PY, Mak KH, Zhang LY, Liu P, Zhang H, Mak NK, Yue PY, Wong RN (2017) Role of G3BP1 in glucocorticoid receptor-mediated microRNA-15b and microRNA-23a biogenesis in endothelial cells. *Cell Mol Life Sci* 74: 3613-3630
- Lee AK, Klein J, Fon Tacer K, Lord T, Oatley MJ, Oatley JM, Porter SN, Prueitt-Miller SM, Tikhonova EB, Karamyshev AL *et al* (2020) Translational Repression of G3BP in Cancer and Germ Cells Suppresses Stress Granules and Enhances Stress Tolerance. *Mol Cell* 79: 645-659 e649

- Leung AK, Vyas S, Rood JE, Bhutkar A, Sharp PA, Chang P (2011) Poly(ADP-ribose) regulates stress responses and microRNA activity in the cytoplasm. *Mol Cell* 42: 489-499
- Li VS, Ng SS, Boersema PJ, Low TY, Karthaus WR, Gerlach JP, Mohammed S, Heck AJ, Maurice MM, Mahmoudi T *et al* (2012) Wnt signaling through inhibition of beta-catenin degradation in an intact Axin1 complex. *Cell* 149: 1245-1256
- Li W, Simarro M, Kedersha N, Anderson P (2004) FAST is a survival protein that senses mitochondrial stress and modulates TIA-1-regulated changes in protein expression. *Mol Cell Biol* 24: 10718-10732
- Li YR, King OD, Shorter J, Gitler AD (2013) Stress granules as crucibles of ALS pathogenesis. *J Cell Biol* 201: 361-372
- Lin CH, Lee HH, Chang WM, Lee FP, Chen LC, Lu LS, Lin YF (2020) FOXD1 Repression Potentiates Radiation Effectiveness by Downregulating G3BP2 Expression and Promoting the Activation of TXNIP-Related Pathways in Oral Cancer. *Cancers (Basel)* 12
- Lindquist ME, Lifland AW, Utley TJ, Santangelo PJ, Crowe JE, Jr. (2010) Respiratory syncytial virus induces host RNA stress granules to facilitate viral replication. *J Virol* 84: 12274-12284
- Liu H, Bai Y, Zhang X, Gao T, Liu Y, Li E, Wang X, Cao Z, Zhu L, Dong Q *et al* (2022) SARS-CoV-2 N Protein Antagonizes Stress Granule Assembly and IFN Production by Interacting with G3BPs to Facilitate Viral Replication. *J Virol* 96: e0041222
- Liu H, Bi J, Dong W, Yang M, Shi J, Jiang N, Lin T, Huang J (2018) Invasion-related circular RNA circFNDC3B inhibits bladder cancer progression through the miR-1178-3p/G3BP2/SRC/FAK axis. *Mol Cancer* 17: 161
- Liu H, Bi J, Dong W, Yang M, Shi J, Jiang N, Lin T, Huang J (2020) Correction to: Invasion-related circular RNA circFNDC3B inhibits bladder cancer progression through the miR-1178-3p/G3BP2/SRC/FAK axis. *Mol Cancer* 19: 124
- Liu S, Chen L, Chen H, Xu K, Peng X, Zhang M (2021) Circ_0119872 promotes uveal melanoma development by regulating the miR-622/G3BP1 axis and downstream signalling pathways. *J Exp Clin Cancer Res* 40: 66

- Liu ZS, Cai H, Xue W, Wang M, Xia T, Li WJ, Xing JQ, Zhao M, Huang YJ, Chen S *et al* (2019) G3BP1 promotes DNA binding and activation of cGAS. *Nat Immunol* 20: 18-28
- Lv S, Ma M, Sun Y, Wang X, Qimuge N, Qin J, Pang W (2017) MicroRNA-129-5p inhibits 3T3-L1 preadipocyte proliferation by targeting G3BP1. *Anim Cells Syst (Seoul)* 21: 269-277
- Lypowy J, Chen IY, Abdellatif M (2005) An alliance between Ras GTPase-activating protein, filamin C, and Ras GTPase-activating protein SH3 domain-binding protein regulates myocyte growth. *J Biol Chem* 280: 25717-25728
- Mao C, Wang X, Liu Y, Wang M, Yan B, Jiang Y, Shi Y, Shen Y, Liu X, Lai W *et al* (2018) A G3BP1-Interacting lncRNA Promotes Ferroptosis and Apoptosis in Cancer via Nuclear Sequestration of p53. *Cancer Res* 78: 3484-3496
- Maruri-Lopez I, Figueroa NE, Hernandez-Sanchez IE, Chodasiewicz M (2021) Plant Stress Granules: Trends and Beyond. *Front Plant Sci* 12: 722643
- Marx J (1989) Science gives ivory a sense of identity. *Science* 246: 1120-1121
- Matsuki H, Takahashi M, Higuchi M, Makokha GN, Oie M, Fujii M (2013) Both G3BP1 and G3BP2 contribute to stress granule formation. *Genes Cells* 18: 135-146
- Matthews JD, Frey TK (2012) Analysis of subcellular G3BP redistribution during rubella virus infection. *J Gen Virol* 93: 267-274
- Mazroui R, Sukarieh R, Bordeleau ME, Kaufman RJ, Northcote P, Tanaka J, Gallouzi I, Pelletier J (2006) Inhibition of ribosome recruitment induces stress granule formation independently of eukaryotic initiation factor 2alpha phosphorylation. *Mol Biol Cell* 17: 4212-4219
- McDonald KK, Aulas A, Destroismaisons L, Pickles S, Beleac E, Camu W, Rouleau GA, Vande Velde C (2011) TAR DNA-binding protein 43 (TDP-43) regulates stress granule dynamics via differential regulation of G3BP and TIA-1. *Hum Mol Genet* 20: 1400-1410
- Meshram CD, Shiliaev N, Frolova EI, Frolov I (2020) Hypervariable Domain of nsP3 of Eastern Equine Encephalitis Virus Is a Critical Determinant of Viral Virulence. *J Virol* 94
- Mukhopadhyay C XL, Liu D, Yang C, Hao A, Deonaraine LD, Cyrta J, Robinson B, Rubin M, Barbieri C, Zhou P. , 2019. Regulation of the tumor suppressive SPOP ubiquitin ligase in prostate cancer, 12th Annual Multi-institutional Prostate Cancer Program Retreat. W Hotel, Fort Lauderdale, Florida.

- Mukhopadhyay C, Yang C, Xu L, Liu D, Wang Y, Huang D, Deonarine LD, Cyrta J, Davicioni E, Sboner A *et al* (2021) G3BP1 inhibits Cul3(SPOP) to amplify AR signaling and promote prostate cancer. *Nat Commun* 12: 6662
- Nagai K, Oubridge C, Ito N, Avis J, Evans P (1995) The RNP domain: a sequence-specific RNA-binding domain involved in processing and transport of RNA. *Trends Biochem Sci* 20: 235-240
- Neyret A, Bernard E, Aiqui-Reboul-Paviet O, Bakhache W, Eldin P, Chaloin L, Briant L (2022) Identification of a non-canonical G3BP-binding sequence in a Mayaro virus nsP3 hypervariable domain. *Front Cell Infect Microbiol* 12: 958176
- Ocana OH, Corcoles R, Fabra A, Moreno-Bueno G, Acloque H, Vega S, Barrallo-Gimeno A, Cano A, Nieto MA (2012) Metastatic colonization requires the repression of the epithelial-mesenchymal transition inducer Prrx1. *Cancer Cell* 22: 709-724
- Oi N, Yuan J, Malakhova M, Luo K, Li Y, Ryu J, Zhang L, Bode AM, Xu Z, Li Y *et al* (2015) Resveratrol induces apoptosis by directly targeting Ras-GTPase-activating protein SH3 domain-binding protein 1. *Oncogene* 34: 2660-2671
- Omer A, Barrera MC, Moran JL, Lian XJ, Di Marco S, Beausejour C, Gallouzi IE (2020) G3BP1 controls the senescence-associated secretome and its impact on cancer progression. *Nat Commun* 11: 4979
- Ortega AD, Willers IM, Sala S, Cuezva JM (2010) Human G3BP1 interacts with beta-F1-ATPase mRNA and inhibits its translation. *J Cell Sci* 123: 2685-2696
- Pandey K, Zhong S, Diel DG, Hou Y, Wang Q, Nelson E, Wang X (2019) GTPase-activating protein-binding protein 1 (G3BP1) plays an antiviral role against porcine epidemic diarrhea virus. *Vet Microbiol* 236: 108392
- Parker F, Maurier F, Delumeau I, Duchesne M, Faucher D, Debussche L, Dugue A, Schweighoffer F, Tocque B (1996) A Ras-GTPase-activating protein SH3-domain-binding protein. *Mol Cell Biol* 16: 2561-2569
- Prentzell MT, Rehbein U, Cadena Sandoval M, De Meulemeester AS, Baumeister R, Brohee L, Berdel B, Bockwoldt M, Carroll B, Chowdhury SR *et al* (2021) G3BPs tether the TSC complex to lysosomes and suppress mTORC1 signaling. *Cell* 184: 655-674 e627

- Prigent M, Barlat I, Langen H, Dargemont C (2000) I κ B and NF- κ B complexes are retained in the cytoplasm through interaction with a novel partner, RasGAP SH3-binding protein 2. *J Biol Chem* 275: 36441-36449
- Ramachandran B, Stabley JN, Cheng SL, Behrmann AS, Gay A, Li L, Mead M, Kozlitina J, Lemoff A, Mirzaei H *et al* (2018) A GTPase-activating protein-binding protein (G3BP1)/antiviral protein relay conveys arteriosclerotic Wnt signals in aortic smooth muscle cells. *J Biol Chem* 293: 7942-7968
- Rao SG, Jackson JG (2016) SASP: Tumor Suppressor or Promoter? Yes! *Trends Cancer* 2: 676-687
- Reineke LC, Dougherty JD, Pierre P, Lloyd RE (2012) Large G3BP-induced granules trigger eIF2 α phosphorylation. *Mol Biol Cell* 23: 3499-3510
- Reineke LC, Kedersha N, Langereis MA, van Kuppeveld FJ, Lloyd RE (2015) Stress granules regulate double-stranded RNA-dependent protein kinase activation through a complex containing G3BP1 and Caprin1. *mBio* 6: e02486
- Reineke LC, Lloyd RE (2015) The stress granule protein G3BP1 recruits protein kinase R to promote multiple innate immune antiviral responses. *J Virol* 89: 2575-2589
- Reineke LC, Tsai WC, Jain A, Kaelber JT, Jung SY, Lloyd RE (2017) Casein Kinase 2 Is Linked to Stress Granule Dynamics through Phosphorylation of the Stress Granule Nucleating Protein G3BP1. *Mol Cell Biol* 37
- Sabile AA, Arlt MJ, Muff R, Husmann K, Hess D, Bertz J, Langsam B, Aemisegger C, Ziegler U, Born W *et al* (2013) Caprin-1, a novel Cyr61-interacting protein, promotes osteosarcoma tumor growth and lung metastasis in mice. *Biochim Biophys Acta* 1832: 1173-1182
- Sanders DW, Kedersha N, Lee DSW, Strom AR, Drake V, Riback JA, Bracha D, Eeftens JM, Iwanicki A, Wang A *et al* (2020) Competing Protein-RNA Interaction Networks Control Multiphase Intracellular Organization. *Cell* 181: 306-324 e328
- Scholte FE, Tas A, Albulescu IC, Zusinaite E, Merits A, Snijder EJ, van Hemert MJ (2015) Stress granule components G3BP1 and G3BP2 play a proviral role early in Chikungunya virus replication. *J Virol* 89: 4457-4469

- Schulte T, Liu L, Panas MD, Thaa B, Dickson N, Gotte B, Achour A, McInerney GM (2016) Combined structural, biochemical and cellular evidence demonstrates that both FGDF motifs in alphavirus nsP3 are required for efficient replication. *Open Biol* 6
- Shi Q, Zhu Y, Ma J, Chang K, Ding D, Bai Y, Gao K, Zhang P, Mo R, Feng K *et al* (2019) Prostate Cancer-associated SPOP mutations enhance cancer cell survival and docetaxel resistance by upregulating Caprin1-dependent stress granule assembly. *Mol Cancer* 18: 170
- Shim JH, Su ZY, Chae JI, Kim DJ, Zhu F, Ma WY, Bode AM, Yang CS, Dong Z (2010) Epigallocatechin gallate suppresses lung cancer cell growth through Ras-GTPase-activating protein SH3 domain-binding protein 1. *Cancer Prev Res (Phila)* 3: 670-679
- Shuang O, Zhou J, Cai Z, Liao L, Wang Y, Wang W, Xu M (2022) EBF1-mediated up-regulation of lncRNA FGD5-AS1 facilitates osteosarcoma progression by regulating miR-124-3p/G3BP2 axis as a ceRNA. *J Orthop Surg Res* 17: 332
- Sidibe H, Khalfallah Y, Xiao S, Gomez NB, Fakim H, Tank EMH, Di Tomasso G, Bareke E, Aulas A, McKeever PM *et al* (2021) TDP-43 stabilizes G3BP1 mRNA: relevance to amyotrophic lateral sclerosis/frontotemporal dementia. *Brain* 144: 3461-3476
- Smith A, Brownawell A, Macara IG (1998) Nuclear import of Ran is mediated by the transport factor NTF2. *Curr Biol* 8: 1403-1406
- Solomon S, Xu Y, Wang B, David MD, Schubert P, Kennedy D, Schrader JW (2007) Distinct structural features of caprin-1 mediate its interaction with G3BP-1 and its induction of phosphorylation of eukaryotic translation initiation factor 2alpha, entry to cytoplasmic stress granules, and selective interaction with a subset of mRNAs. *Mol Cell Biol* 27: 2324-2342
- Somasekharan SP, El-Naggar A, Leprivier G, Cheng H, Hajee S, Grunewald TG, Zhang F, Ng T, Delattre O, Evdokimova V *et al* (2015) YB-1 regulates stress granule formation and tumor progression by translationally activating G3BP1. *J Cell Biol* 208: 913-929
- Somasekharan SP, Saxena N, Zhang F, Beraldi E, Huang JN, Gentle C, Fazli L, Thi M, Sorensen PH, Gleave M (2022) Regulation of AR mRNA translation in response to acute AR pathway inhibition. *Nucleic Acids Res* 50: 1069-1091

- Soncini C, Berdo I, Draetta G (2001) Ras-GAP SH3 domain binding protein (G3BP) is a modulator of USP10, a novel human ubiquitin specific protease. *Oncogene* 20: 3869-3879
- Sonnhof U, Grafe P, Krumnikl J, Linder M, Schindler L (1975) Inhibitory postsynaptic actions of taurine, GABA and other amino acids on motoneurons of the isolated frog spinal cord. *Brain Res* 100: 327-341
- Sun Y, Dong L, Yu S, Wang X, Zheng H, Zhang P, Meng C, Zhan Y, Tan L, Song C *et al* (2017) Newcastle disease virus induces stable formation of bona fide stress granules to facilitate viral replication through manipulating host protein translation. *FASEB J* 31: 1337-1353
- Suyama M, Doerks T, Braun IC, Sattler M, Izaurralde E, Bork P (2000) Prediction of structural domains of TAP reveals details of its interaction with p15 and nucleoporins. *EMBO Rep* 1: 53-58
- Takayama KI, Suzuki T, Tanaka T, Fujimura T, Takahashi S, Urano T, Ikeda K, Inoue S (2018) TRIM25 enhances cell growth and cell survival by modulating p53 signals via interaction with G3BP2 in prostate cancer. *Oncogene* 37: 2165-2180
- Taniuchi K, Nishimori I, Hollingsworth MA (2011a) Intracellular CD24 inhibits cell invasion by posttranscriptional regulation of BART through interaction with G3BP. *Cancer Res* 71: 895-905
- Taniuchi K, Nishimori I, Hollingsworth MA (2011b) The N-terminal domain of G3BP enhances cell motility and invasion by posttranscriptional regulation of BART. *Mol Cancer Res* 9: 856-866
- Tourriere H, Chebli K, Zekri L, Courselaud B, Blanchard JM, Bertrand E, Tazi J (2003) The RasGAP-associated endoribonuclease G3BP assembles stress granules. *J Cell Biol* 160: 823-831
- Tourriere H, Gallouzi IE, Chebli K, Capony JP, Mouaikel J, van der Geer P, Tazi J (2001) RasGAP-associated endoribonuclease G3Bp: selective RNA degradation and phosphorylation-dependent localization. *Mol Cell Biol* 21: 7747-7760
- Tsai WC, Gayatri S, Reineke LC, Sbardella G, Bedford MT, Lloyd RE (2016) Arginine Demethylation of G3BP1 Promotes Stress Granule Assembly. *J Biol Chem* 291: 22671-22685
- Tsai WC, Reineke LC, Jain A, Jung SY, Lloyd RE (2017) Histone arginine demethylase JMJD6 is linked to stress granule assembly through demethylation of the stress granule-nucleating protein G3BP1. *J Biol Chem* 292: 18886-18896

- Valiente-Echeverria F, Melnychuk L, Vyboh K, Ajamian L, Gallouzi IE, Bernard N, Mouland AJ (2014) eEF2 and Ras-GAP SH3 domain-binding protein (G3BP1) modulate stress granule assembly during HIV-1 infection. *Nat Commun* 5: 4819
- Vognsen T, Moller IR, Kristensen O (2013) Crystal structures of the human G3BP1 NTF2-like domain visualize FxFG Nup repeat specificity. *PLoS One* 8: e80947
- Wang C, Horby PW, Hayden FG, Gao GF (2020) A novel coronavirus outbreak of global health concern. *Lancet* 395: 470-473
- Wang S, Merits A (2022) G3BP/Rin-Binding Motifs Inserted into Flexible Regions of nsP2 Support RNA Replication of Chikungunya Virus. *J Virol* 96: e0127822
- Wang Y, Fu D, Chen Y, Su J, Li X, Zhai W, Niu Y, Yue D, Geng H (2018) G3BP1 promotes tumor progression and metastasis through IL-6/G3BP1/STAT3 signaling axis in renal cell carcinomas. *Cell Death Dis* 9: 501
- Wei SC, Fattet L, Tsai JH, Guo Y, Pai VH, Majeski HE, Chen AC, Sah RL, Taylor SS, Engler AJ *et al* (2015) Matrix stiffness drives epithelial-mesenchymal transition and tumour metastasis through a TWIST1-G3BP2 mechanotransduction pathway. *Nat Cell Biol* 17: 678-688
- Wen W, Zhao Q, Yin M, Qin L, Hu J, Chen H, Li X, Qian P (2020) Seneca Valley Virus 3C Protease Inhibits Stress Granule Formation by Disrupting eIF4GI-G3BP1 Interaction. *Front Immunol* 11: 577838
- White JP, Lloyd RE (2012) Regulation of stress granules in virus systems. *Trends Microbiol* 20: 175-183
- Wilczynska A, Aigueperse C, Kress M, Dautry F, Weil D (2005) The translational regulator CPEB1 provides a link between dcp1 bodies and stress granules. *J Cell Sci* 118: 981-992
- Winslow S, Leandersson K, Larsson C (2013) Regulation of PMP22 mRNA by G3BP1 affects cell proliferation in breast cancer cells. *Mol Cancer* 12: 156
- Wolozin B, Ivanov P (2019) Stress granules and neurodegeneration. *Nat Rev Neurosci* 20: 649-666
- Xia J, Chen X, Xu F, Wang Y, Shi Y, Li Y, He J, Zhang P (2015) Dengue virus infection induces formation of G3BP1 granules in human lung epithelial cells. *Arch Virol* 160: 2991-2999

- Xia J, Zhang J, Chang J, Tian Y, Li J, Zhang B, Zeng X, Yin C (2020) The effects of glycaemic variability on intimal hyperplasia and plaque stability after stenting via autophagy-mediated G3BP1/NLRP3 inflammasome. *Ann Transl Med* 8: 1388
- Yang P, Mathieu C, Kolaitis RM, Zhang P, Messing J, Yurtsever U, Yang Z, Wu J, Li Y, Pan Q *et al* (2020a) G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble Stress Granules. *Cell* 181: 325-345 e328
- Yang W, Li D, Ru Y, Bai J, Ren J, Zhang J, Li L, Liu X, Zheng H (2020b) Foot-and-Mouth Disease Virus 3A Protein Causes Upregulation of Autophagy-Related Protein LRRC25 To Inhibit the G3BP1-Mediated RIG-Like Helicase-Signaling Pathway. *J Virol* 94
- Ye X, Pan T, Wang D, Fang L, Ma J, Zhu X, Shi Y, Zhang K, Zheng H, Chen H *et al* (2018) Foot-and-Mouth Disease Virus Counteracts on Internal Ribosome Entry Site Suppression by G3BP1 and Inhibits G3BP1-Mediated Stress Granule Assembly via Post-Translational Mechanisms. *Front Immunol* 9: 1142
- Yi Z, Fang C, Pan T, Wang J, Yang P, Yuan Z (2006) Subproteomic study of hepatitis C virus replicon reveals Ras-GTPase-activating protein binding protein 1 as potential HCV RC component. *Biochem Biophys Res Commun* 350: 174-178
- Zekri L, Chebli K, Tourriere H, Nielsen FC, Hansen TV, Rami A, Tazi J (2005) Control of fetal growth and neonatal survival by the RasGAP-associated endoribonuclease G3BP. *Mol Cell Biol* 25: 8703-8716
- Zhang CH, Liu H, Zhao WL, Zhao WX, Zhou HM, Shao RG (2021a) G3BP1 promotes human breast cancer cell proliferation through coordinating with GSK-3beta and stabilizing beta-catenin. *Acta Pharmacol Sin* 42: 1900-1912
- Zhang CH, Wang JX, Cai ML, Shao R, Liu H, Zhao WL (2019a) The roles and mechanisms of G3BP1 in tumour promotion. *J Drug Target* 27: 300-305
- Zhang H, Ma Y, Zhang S, Liu H, He H, Li N, Gong Y, Zhao S, Jiang JD, Shao RG (2015) Involvement of Ras GTPase-activating protein SH3 domain-binding protein 1 in the epithelial-to-mesenchymal transition-induced metastasis of breast cancer cells via the Smad signaling pathway. *Oncotarget* 6: 17039-17053

- Zhang H, Zhang S, He H, Zhao W, Chen J, Shao RG (2012) GAP161 targets and downregulates G3BP to suppress cell growth and potentiate cisplatin-mediated cytotoxicity to colon carcinoma HCT116 cells. *Cancer Sci* 103: 1848-1856
- Zhang H, Zhang SH, He HW, Zhang CX, Yu DK, Shao RG (2013) Downregulation of G3BPs inhibits the growth, migration and invasion of human lung carcinoma H1299 cells by suppressing the Src/FAK-associated signaling pathway. *Cancer Gene Ther* 20: 622-629
- Zhang LN, Zhao L, Yan XL, Huang YH (2019b) Loss of G3BP1 suppresses proliferation, migration, and invasion of esophageal cancer cells via Wnt/beta-catenin and PI3K/AKT signaling pathways. *J Cell Physiol* 234: 20469-20484
- Zhang Y, Yue C, Krichevsky AM, Garkavtsev I (2021b) Repression of the stress granule protein G3BP2 inhibits immune checkpoint molecule PD-L1. *Mol Oncol*
- Zhao B, Li H, Liu J, Han P, Zhang C, Bai H, Yuan X, Wang X, Li L, Ma H *et al* (2016) MicroRNA-23b Targets Ras GTPase-Activating Protein SH3 Domain-Binding Protein 2 to Alleviate Fibrosis and Albuminuria in Diabetic Nephropathy. *J Am Soc Nephrol* 27: 2597-2608
- Zhao D, Li J, Wang Y, Li X, Gao L, Cao H, Zheng SJ (2020) Critical role for G3BP1 in infectious bursal disease virus (IBDV)-induced stress granule formation and viral replication. *Vet Microbiol* 248: 108806
- Zhao H, He Y (2021) MiR-124-3p Suppresses the Dysfunction of High Glucose-Stimulated Endothelial Cells by Targeting G3BP2. *Front Genet* 12: 723625
- Zheng Y, Wu J, Deng R, Lin C, Huang Y, Yang X, Wang C, Yang M, He Y, Lu J *et al* (2022) G3BP2 regulated by the lncRNA LINC01554 facilitates esophageal squamous cell carcinoma metastasis through stabilizing HDGF transcript. *Oncogene* 41: 515-526
- Zhou Y, Guan J, Gao F, Li Z, Lan Y, Lu H, Song D, Lv L, Lv P, Xu M *et al* (2021) Orf Virus ORF120 Protein Positively Regulates the NF-kappaB Pathway by Interacting with G3BP1. *J Virol* 95: e0015321
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R *et al* (2020) A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 382: 727-733

Footnotes

This work was supported in part by R01 CA221152, CTSC (1 UL1 TR002384) Pilot Award and the Weill Cornell Medicine Prostate SPORE's Developmental Research Award to P.Z. and US Department of Defense (W81XWH-17-1-0176) to C.M. We thank Dr. Sarah See for critical reading and editing of the manuscript and constructive comments and suggestions.

The authors declare that they have no conflict of interest with any company that could benefit from the findings of this manuscript.

Figure Legends

Figure 1. Schematic representation of different domain structure of G3BP1 and G3BP2.

Both G3BP1 and G3BP2 have an NTF2 domain at the N-terminus and an RRM/RGG domain at the C-terminus. However, G3BP1 has only one PxxP domain, while G3BP2a and G3BP2b contain four and five PxxP domains, respectively.

Figure 2. Summary of well-characterized and emerging roles of (a) G3BP1 and (b) G3BP2

(a) Different functions of G3BP1 in cancer, stress granule formation, RNA stabilization, and viral infections. **(b)** Different functions of G3BP2 in cancer, stress granule formation, and RNA stabilization.





Figure 1

(a)



(b)




Figure 2